

Lectures on Mathematical Cosmology

Hans-Jürgen Schmidt

Universität Potsdam, Institut für Mathematik, Am Neuen Palais 10

D-14469 Potsdam, Germany, E-mail: hjschmi@rz.uni-potsdam.de

<http://www.physik.fu-berlin.de/~hjschmi>

©2004, H.-J. Schmidt

Contents

1	Preface	7
2	Changes of the Bianchi type	9
2.1	Introduction to generalized Bianchi models	9
2.2	Spaces possessing homogeneous slices	11
2.3	Continuous changes of the Bianchi type	13
2.4	Physical conditions	16
3	Inhomogeneous models with flat slices	19
3.1	Models with flat slices	20
3.2	Energy inequalities and perfect fluid models	22
3.3	A simple singularity theorem	26
4	Properties of curvature invariants	31
4.1	The space of 3-dimensional Riemannian manifolds	31
4.2	Why do all the invariants of a gravitational wave vanish? . . .	35
4.2.1	Preliminaries	36
4.2.2	Gravitational waves	39
4.2.3	Topological properties	41
4.3	Spacetimes which cannot be distinguished by invariants	43
5	Surface layers and relativistic surface tensions	45
5.1	Introduction to surface layers	45
5.2	Non-relativistic surface tensions	47
5.3	Mean curvature in curved spacetime	49
5.4	Equation of motion for the surface layer	51
5.5	Notation, distributions and mean curvature	52

6	The massive scalar field in a Friedmann universe	57
6.1	Some closed-form approximations	59
6.2	The qualitative behaviour	62
6.2.1	Existence of a maximum	63
6.2.2	The space of solutions	63
6.2.3	From one extremum to the next	64
6.2.4	The periodic solutions	66
6.2.5	The aperiodic perpetually oscillating solutions	67
6.3	Problems with the probability measure	70
6.3.1	The spatially flat Friedmann model	70
6.3.2	The Bianchi-type I model	71
6.3.3	The closed Friedmann model	72
6.4	Discussion of the inflationary phase	73
7	The superspace of Riemannian metrics	75
7.1	The superspace	75
7.2	Coordinates in superspace	76
7.3	Metric in superspace	77
7.4	Signature of the superspace metric	79
7.5	Supercurvature and superdeterminant	81
7.6	Gravity and quantum cosmology	83
7.7	The Wheeler-DeWitt equation	85
8	Scalar fields and $f(R)$ for cosmology	87
8.1	Introduction to scalar fields	87
8.2	The Higgs field	89
8.3	The non-linear gravitational Lagrangian	90
8.3.1	Calculation of the ground states	92

8.3.2	Definition of the masses	93
8.4	The cosmological model	94
8.4.1	The field equation	94
8.4.2	The masses	95
8.4.3	The Friedmann model	96
8.5	The generalized Bicknell theorem	100
8.6	On Ellis' programme within homogeneous world models	104
8.6.1	Ellis' programme	104
8.6.2	The massive scalar field in a Bianchi-type I model . . .	106
8.6.3	The generalized equivalence	110
8.6.4	The fourth-order gravity model	111
9	Models with De Sitter and power-law inflation	113
9.1	Differential-geometrical properties	114
9.2	Scale-invariant field equations	115
9.3	Cosmological Friedmann models	116
9.3.1	The closed and open models	117
9.3.2	The spatially flat model	119
9.4	Discussion of power-law inflation	120
9.5	The cosmic no hair theorem	121
9.5.1	Introduction to no hair theorems	122
9.5.2	Definition of an asymptotic de Sitter spacetime	130
9.5.3	Lagrangian $F(R, \square R, \square^2 R, \dots, \square^k R)$	134
9.5.4	No hair theorems for higher-order gravity	135
9.5.5	Diagonalizability of Bianchi models	138
9.5.6	Discussion of no hair theorems	141

10 Solutions of the Bach-Einstein equation	145
10.1 Introduction to the Bach equation	145
10.2 Notations and field equations	147
10.3 Linearizations	149
10.4 Non-standard analysis	151
10.5 Discussion – Einstein’s particle programme	154
10.6 Non-trivial solutions of the Bach equation exist	156
11 The Newtonian limit of non-linear gravity	159
11.1 The Newtonian limit of 4th-order gravity	159
11.2 Introduction to higher-order gravity	164
11.3 Lagrangian and field equation	165
11.4 Newtonian limit in higher-order gravity	166
11.5 Discussion of the weak-field limit	170
11.6 A homogeneous sphere	172
12 Cosmic strings and gravitational waves	175
12.1 Introduction to cosmic strings	175
12.2 The main formulae	176
12.3 The string in a Friedmann model	178
12.3.1 The open string	179
12.3.2 The closed string	181
12.4 The string in an anisotropic Kasner background	182
12.5 The string in a gravitational wave	183
13 Bibliography	187

1 Preface

We present mathematical details of several cosmological models, whereby the topological and the geometrical background will be emphasized.

This book arose from lectures I read as advanced courses in the following universities: 1990 at TU Berlin, 1991 in Münster, 1993 - 1996 and 2002, 2003 in Potsdam, 1999 and 2000 in Salerno, and 2001 at FU Berlin.

The reader is assumed to be acquainted with basic knowledge on general relativity, e.g. by knowing Stephani's book [250]. As a rule, the deduction is made so explicit, that one should be able to follow the details without further background knowledge. The intention is to present a piece of mathematical physics related to cosmology to fill the gap between standard textbooks like MTW¹, LL², or Weinberg³, on the one hand and to the current research literature on the other. I hope that this will become useful for both students and researchers who are interested in cosmology, mathematical physics or differential geometry.

To the reference list: If more than one source is given in one item, then a dot is put between them. If, however, two versions of the same source are given, then a semicolon is put between them. The gr-qc number in the reference list refers to the number in the preprint archive

<http://xxx.lanl.gov/find/>

and the numbers in italics given at the end of an item show the pages where this item is mentioned. This way, the Bibliography simultaneously serves as

¹C. Misner, K. Thorne, J. Wheeler *Gravitation*, Freeman, San Francisco 1973

²L. Landau, E. Lifschitz *Field Theory*, Nauka, Moscow 1973

³S. Weinberg *Gravitation and Cosmology*, J. Wiley, New York 1972

author index.

Parts of this book⁴ are composed from revised versions of papers of mine published earlier in research journals. They are reprinted with the kind permissions of the copyright owners: Akademie-Verlag Berlin for *Astron. Nachr.* and Plenum Press New York for *Gen. Rel. Grav.* The details about the originals can also be seen in

<http://www.physik.fu-berlin.de/~hjschmi>

I thank the following colleagues: Luca Amendola, Michael Bachmann, H. Baumgärtel, W. Benz, Claudia Bernutat, Ulrich Bleyer, A. Borde, H.-H. v. Borzeszkowski, Salvatore Capozziello, G. Dautcourt, Vladimir Dzhunushaliev, Mauro Francaviglia, T. Friedrich, V. Frolov, H. Goenner, Stefan Gottlöber, V. Gurovich, Graham Hall, F. Hehl, Olaf Heinrich, Alan Held, L. Herrera, Rainer John, U. Kasper, Jerzy Kijowski, Hagen Kleinert, B. Klotzek, Sabine Kluske, O. Kowalski, Andrzej Krasinski, W. Kühnel, W. Kundt, K. Lake, K. Leichtweiss, D.-E. Liebscher, Malcolm MacCallum, W. Mai, R. Mansouri, V. Melnikov, P. Michor, Salvatore Mignemi, Volker Müller, S. Nikcevic, I. Novikov, F. Paiva, Axel Pelster, V. Perlick, K. Peters, Ines Quandt, L. Querella, M. Rainer, J. Reichert, Stefan Reuter, W. Rinow, M. Sanchez, R. Schimming, M. Schulz, E. Schmutzer, U. Semmelmann, I. Shapiro, A. Starobinsky, P. Teyssandier, H.-J. Treder, H. Tuschik, L. Vanhecke and A. Zhuk for valuable comments, and my wife Renate for patience and support.

Hans-Jürgen Schmidt, Potsdam, July 2004

⁴The details are mentioned at the corresponding places.

2 Cosmological models with changes of the Bianchi type

We investigate such cosmological models which instead of the usual spatial homogeneity property only fulfil the condition that in a certain synchronized system of reference all spacelike sections $t = \text{const.}$ are homogeneous manifolds.

This allows time-dependent changes of the Bianchi type. Discussing differential-geometrical theorems it is shown which of them are permitted. Besides the trivial case of changing into type I there exist some possible changes between other types. However, physical reasons like energy inequalities partially exclude them.

2.1 Introduction to generalized Bianchi models

Recently, besides the known Bianchi models, there are investigated certain classes of inhomogeneous cosmological models. This is done to get a better representation of the really existent inhomogeneities, cf. e.g. Bergmann [23], Carmeli [51], Collins [55], Spero [240], Szekeres [251] and Wainwright [265].

We consider, similar as in Collins [55], such inhomogeneous models V_4 which in a certain synchronized system of reference possess homogeneous sections $t = \text{const.}$, called $V_3(t)$. This is analogous to the generalization of the concept of spherical symmetry in Krasinski [136], cf. also [137].

In this chapter, which is based on [200], we especially investigate which time-dependent changes of the Bianchi type are possible. Thereby we impose, besides the twice continuous differentiability, a physically reasonable

condition: the energy inequality,

$$T_{00} \geq |T_{\alpha\beta}|, \quad (2.1)$$

holds in each Lorentz frame.

We begin with some globally topological properties of spacetime: Under the physical condition (2.1) the topology of the sections $V_3(t)$ is, according to Lee [142], independent of t . Hence, the Kantowski-Sachs models, with underlying topology $S^2 \times \mathbb{R}$ or $S^2 \times S^1$ and the models of Bianchi type IX with underlying topology S^3 or continuous images of it as $SO(3)$ may not change, because all other types are represented by the $\mathbb{R}^3$ -topology, factorized with reference to a discrete subgroup of the group of motions. But the remaining types can all be represented in $\mathbb{R}^3$ -topology itself; therefore we do not get any further global restrictions. All homogeneous models, the above mentioned Kantowski-Sachs model being excluded, possess simply transitive groups of motion. If the isometry group G has dimension ≥ 4 , then this statement means: G possesses a 3-dimensional simply-transitive subgroup acting transitively on the $V_3(t)$. Hence we do not specialize if we deal only with locally simply-transitive groups of motions. We consequently do not consider here trivial changes of the Bianchi type, e. g. from type III to type VIII by means of an intermediate on which a group of motions possessing transitive subgroups of both types acts.

Next, we consider the easily tractable case of a change to type I: For each type M there one can find a manifold V_4 such that for each $t \leq 0$ the section $V_3(t)$ is flat and for each $t > 0$ it belongs to type M. Indeed, one has simply to use for every $V_3(t)$ such a representative of type M that their curvature vanishes as $t \rightarrow 0_+$. Choosing exponentially decreasing curvature one can

obtain an arbitrarily high differentiable class for the metric, e. g.

$$ds^2 = -dt^2 + dr^2 + h^2 \cdot (d\psi^2 + \sin^2 \psi d\phi^2), \quad (2.2)$$

where

$$h = \begin{cases} r & \text{for } t \leq 0 \\ \exp(t^{-2}) \cdot \sinh^2(r \cdot \exp(-t^{-2})) & \text{else.} \end{cases} \quad (2.3)$$

This is a C^∞ -metric whose slices $V_3(t)$ belong to type I for $t \leq 0$ and to type V for $t > 0$. In $t = 0$, of course, it cannot be an analytical one. The limiting slice belongs necessarily to type I by continuity reasons. Applying this fact twice it becomes obvious that by the help of a flat intermediate of finite extension or only by a single flat slice, all Bianchi types can be matched together. However, if one does not want to use such a flat intermediate the transitions of one Bianchi type to another become a non-trivial problem. It is shown from the purely differential-geometrical as well as from the physical points of view, sections 2.3 and 2.4 respectively, which types can be matched together immediately without a flat intermediate. To this end we collect the following preliminaries.

2.2 Spaces possessing homogeneous slices

As one knows, in cosmology the homogeneity principle is expressed by the fact that to a spacetime V_4 there exists a group of motions acting transitively on the spacelike hypersurfaces $V_3(t)$ of a slicing of V_4 . Then the metric is given by

$$ds^2 = -dt^2 + g_{ab}(t)\omega^a\omega^b \quad (2.4)$$

where the $g_{ab}(t)$ are positively definite and ω^a are the basic 1-forms corresponding to a certain Bianchi type; for details see e.g. [134]. If the ω^a are

related to a holonomic basis x^i , using type-dependent functions $A_i^a(x^j)$, one can write:

$$\omega^a = A_i^a dx^i. \quad (2.5)$$

For the spaces considered here we have however: there is a synchronized system of reference such that the slices $t = \text{const.}$ are homogeneous spaces $V_3(t)$. In this system of reference the metric is given by

$$ds^2 = -dt^2 + g_{ij} dx^i dx^j \quad (2.6)$$

where $g_{ij}(x^i, t)$ are twice continuously differentiable and homogeneous for constant t . Hence it is a generalization of eqs. (2.4) and (2.5). This is a genuine generalization because there is only posed the condition $g_{0\alpha} = -\delta_{0\alpha}$ on the composition of the homogeneous slices $V_3(t)$ to a V_4 . This means in each slice only the first fundamental form g_{ij} is homogeneous; but in the contrary to homogeneous models, the second fundamental form Γ_{0ab} need not have this property. Therefore, also the curvature scalar ${}^{(4)}R$, and with it the distribution of matter, need not be constant within a $V_3(t)$.

Now let t be fixed. Then one can find coordinates $x_t^i(x^j, t)$ in $V_3(t)$ such that according to eqs. (2.4) and (2.5) the inner metric gets the form

$$g_{ab}(t) \omega_t^a \omega_t^b \quad \text{where} \quad \omega_t^a = A_i^a(x_t^i) dx_t^i. \quad (2.7)$$

Transforming this into the original coordinates x^i one obtains for the metric of the full V_4

$$g_{0\alpha} = -\delta_{0\alpha}, \quad g_{ij} = g_{ab}(t) A_k^a(x_t^i) A_l^b(x_t^i) x_{t,i}^k x_{t,j}^l. \quad (2.8)$$

If the Bianchi type changes with time one has to take such an ansatz⁵ like eqs. (2.2), (2.3) for each interval of constant type separately; between them one has to secure a C^2 -joining. An example of such an inhomogeneous model is, cf. Ellis 1967 [70]

$$ds^2 = -dt^2 + t^{-2/3} [t + C(x)]^2 dx^2 + t^{4/3} (dy^2 + dz^2). \quad (2.9)$$

The slices $t = \text{const.}$ are flat, but only for constant C it belongs to Bianchi type I, cf. also [99] for such a model. Later, see eq. (3.19), we will present a more general model.

2.3 Continuous changes of the Bianchi type

Using the usual homogeneity property the same group of motions acts on each slice $V_3(t)$. Hence the Bianchi type is independent of time by definition. But this fails to be the case for the spaces considered here. However, we can deduce the following: completing $\partial/\partial t$ to an anholonomic basis which is connected with Killing vectors in each $V_3(t)$ one obtains for the structure constants associated to the commutators of the basis:

$$C_{\alpha\beta}^0 = 0; \quad C_{jk}^i(t) \quad \text{depends continuously on time.} \quad (2.10)$$

Note: These structure constants are calculated as follows: Without loss of generality let $x_t^i(0, 0, 0, t) = 0$. At this point $\partial/\partial x_t^i$ and $\partial/\partial x^i$ are taken as initial values for Killing vectors within $V_3(t)$. The structure constants obtained by these Killing vectors are denoted by $\bar{C}_{jk}^i(t)$ and $C_{jk}^i(t)$ respectively, where $\bar{C}_{jk}^i(t)$ are the canonical ones. Between them it holds at $(0, 0, 0, t)$:

$$x_{t,j}^i C_{kl}^j = \bar{C}_{mn}^i x_{t,k}^m x_{t,l}^n. \quad (2.11)$$

⁵The word “ansatz” (setting) stems from the German noun “der Ansatz”; that word stems from the German verb “setzen” (to set).

Using only the usual canonical structure constants then of course no type is changeable continuously. To answer the question which Bianchi types may change continuously we consider all sets of structure constants C_{jk}^i being antisymmetric in jk and fulfilling the Jacobi identity. Let C_S be such a set belonging to type S. Then we have according to eq. (2.10): Type R is changeable continuously into type S, symbolically expressed by the validity of $R \rightarrow S$, if and only if to each C_S there exists a sequence $C_R^{(n)}$ such that one has

$$\lim_{n \rightarrow \infty} C_R^{(n)} = C_S. \quad (2.12)$$

The limit has to be understood componentwise.

It holds that $R \rightarrow S$ if and only if there are a C_S and a sequence $C_R^{(n)}$ such that eq. (2.12) is fulfilled. The equivalence of both statements is shown by means of simultaneous rotations of the basis. In practice one takes as C_S the canonical structure constants and investigates for which types R there can be found corresponding sequences $C_R^{(n)}$: the components of C_S are subjected to a perturbation not exceeding ε and their Bianchi types are calculated. Finally one looks which types appear for all $\varepsilon > 0$. Thereby one profits, e.g., from the statement that the dimension of the image space of the Lie algebra, which equals 0 for type I, ..., and equals 3 for types VIII and IX, cannot increase during such changes; see [189] and [210] for more details about these Lie algebras.

One obtains the following diagram. The validity of $R \rightarrow S$ and $S \rightarrow T$ implies the validity of $R \rightarrow T$, hence the diagram must be continued transitively. The statement $VI_\infty \rightarrow IV$ expresses the facts that there exists a sequence

$$C_{VI_h}^{(n)} \rightarrow C_{IV} \quad (2.13)$$

and that in each such a sequence the parameter h must necessarily tend to infinity.

Further $\text{VI}_h \rightarrow \text{II}$ holds for every h . Both statements hold analogously for type VII_h . In MacCallum [146] a similar diagram is shown, but there it is only investigated which changes appear if some of the canonical structure constants are vanishing. So e.g., the different transitions from type VI_h to types II and IV are not contained in it.




Fig. 1, taken from [200]

To complete the answer to the question posed above it must be added: To each transition shown in the diagram, there one can indeed find a spacetime V_4 in which it is realized. Let $f(t)$ be a C^∞ -function with $f(t) = 0$ for $t \leq 0$ and $f(t) > 0$ else. Then, e.g.,

$$ds^2 = -dt^2 + dx^2 + e^{2x} dy^2 + e^{2x} (dz + x f(t) dy)^2 \quad (2.14)$$

is a C^∞ -metric whose slices $V_3(t)$ belong to type V and IV for $t \leq 0$ and $t > 0$ respectively. Presumably it is typical that in a neighbourhood of $t = 0$ the curvature becomes singular at spatial infinity; this is at least the case for metric (2.14). And, for the transition $R \rightarrow S$ the limiting slice belongs necessarily to type S.

2.4 Physical conditions

To obtain physically reasonable spacetimes one has at least to secure the validity of an energy inequality like (2.1), $T_{00} \geq |T_{\alpha\beta}|$ in each Lorentz frame. Without this requirement the transition $II \rightarrow I$ is possible. We prove that the requirement $T_{00} \geq 0$ alone is sufficient to forbid this transition. To this end let V_4 be a manifold which in a certain synchronized system of reference possesses a flat slice $V_3(0)$, and for all $t > 0$ has Bianchi type II-slices $V_3(t)$. Using the notations of eq. (2.11) we have: $\bar{C}_{23}^1 = -\bar{C}_{32}^1 = 1$ are the only non-vanishing canonical structure constants of Bianchi type II. With the exception $A_2^1 = -x^3$ we have $A_i^a = \delta_i^a$.

By the help of eq. (2.11) one can calculate the structure constants $C_{jk}^i(t)$. The flat-slice condition is equivalent to

$$\lim_{t \rightarrow 0} C_{jk}^i(t) = 0. \quad (2.15)$$

First we consider the special case $x_t^i = a^i(t) \cdot x^i$ (no sum), i. e. extensions of the coordinate axes. It holds $x_{t,j}^i = \delta_j^i a^i$ (no sum). The only non-vanishing $C_{jk}^i(t)$ are $C_{23}^1 = -C_{32}^1 = a(t)$, where $a = a^2 \cdot a^3 / a^1$. Then eq. (2.15) reads $\lim_{t \rightarrow 0} a(t) = 0$. The coefficients $g_{ab}(t)$ have to be chosen such that the g_{ij} , according to eq. (2.8), remain positive definite and twice continuously differentiable. Let ${}^{(3)}R$ be the scalar curvature within the slices. ${}^{(3)}R > 0$ appears only in Bianchi type IX and in the Kantowski-Sachs models. But changes from these types are just the cases already excluded by global considerations. Inserting g_{ij} into the Einstein equation by means of the Gauss-Codazzi theorem one obtains

$$\kappa T_{00} = R_{00} - \frac{1}{2} g_{00} {}^{(4)}R = \frac{1}{2} {}^{(3)}R + \frac{1}{4} g^{-1} \cdot H, \quad g = \det g_{ij}, \quad (2.16)$$

${}^{(3)}R$ being the scalar curvature within the slices, hence

$$\begin{aligned} {}^{(3)}R \leq 0 \quad \text{and} \quad H &= g_{11} [g_{22,0} g_{33,0} - (g_{23,0})^2] \\ &- 2 \cdot g_{12} [g_{33,0} g_{12,0} - g_{13,0} g_{23,0}] + \text{cyclic perm.} \end{aligned} \quad (2.17)$$

In our case H is a quadratic polynomial in x^3 whose quadratic coefficient reads

$$-g_{11} [g_{11} g_{33} - (g_{13})^2] \cdot \left[\frac{\partial a(t)}{\partial t} \right]^2. \quad (2.18)$$

Hence, for sufficiently large values x^3 and values t with $\partial a / \partial t \neq 0$ we have $H < 0$, and therefore $T_{00} < 0$.

Secondly we hint at another special case, namely rotations of the coordinate axes against each other, e.g.

$$x_t^1 = x^1 \cos \omega + x^2 \sin \omega, \quad x_t^2 = x^2 \cos \omega - x^1 \sin \omega, \quad x_t^3 = x^3, \quad \omega = \omega(t).$$

There one obtains in analogy to eq. (2.18) a negative T_{00} . Concerning the general case we have: loosely speaking, each diffeomorphism is a composite of such extensions and rotations. Hence, for each $\text{II} \rightarrow \text{I}$ -transition one would obtain points with negative T_{00} .

Hence, the energy condition is a genuine restriction to the possible transitions of the Bianchi types. Concerning the other transitions we remark: equations (2.16), (2.17) keep valid, and the remaining work is to examine the signs of the corresponding expressions H . Presumably one always obtains points with negative H , i.e., also under this weakened homogeneity presumptions the Bianchi types of the spacelike hypersurfaces at different distances from the singularity must coincide.

3 Inhomogeneous cosmological models with flat slices

A family of cosmological models is considered which in a certain synchronized system of reference possess flat slices $t = \text{const.}$ They are generated from the Einstein-De Sitter universe by a suitable transformation. Under physically reasonable presumptions these transformed models fulfil certain energy conditions. In Wainwright [265] a class of inhomogeneous cosmological models is considered which have the following property: there exists a synchronized system of reference of such a kind that the slices $t = \text{const.}$ are homogeneous manifolds.

Here we consider a special family of such models which possess flat slices. Following [201], we use the transformation formalism developed in chapter 2. In Krasinski [137], e.g. in Fig. 2.1., it is shown how this approach fits into other classes of cosmological solutions of the Einstein field equation. Additionally, we require that these transformations leave two coordinates unchanged; this implies the existence of a 2-dimensional Abelian group of motions. A similar requirement is posed in [265], too.

Starting from a Friedmann universe, we investigate whether the energy inequalities are fulfilled in the transformed model, too. In general this fails to be the case, but starting from the Einstein-De Sitter universe [68], cf. also Tolman [256], and requiring perfect fluid for the transformed model, the energy inequalities in the initial model imply their validity in the transformed model. A new review on exact solutions can be found in [26], and a new proof of Birkhoff's theorem is given in [226].

3.1 Models with flat slices

The transformation formalism of chapter 2 restricted to Bianchi type I reads as follows: using the same notations eq. (2.5), we have $A_i^a = \delta_i^a$, $\omega^a = dx^a$ and

$$ds^2 = -dt^2 + g_{ab}(t)dx^a dx^b \quad (3.1)$$

as the initial hypersurface-homogeneous model instead of eq. (2.4). Now let us consider the time-dependent transformation $x_t^a(x^i, t)$, where for each t it has to be a diffeomorphism of R^3 . Then one obtains in place of eq. (2.8)

$$g_{0\alpha} = -\delta_{0\alpha}, \quad g_{ij} = g_{ab}(t)x_{t,i}^a x_{t,j}^b \quad (3.2)$$

as the transformed model. It is no restriction to insert $g_{ab}(t) = \delta_{ab}$ in eq. (3.1), i.e. to start from Minkowski spacetime. In the following we consider only transformations which leave two coordinates unchanged, i.e., now writing t, x, y, z instead of $x^0, \dots, x^3$ resp., we restrict to transformations which read as follows

$$x_t(x, t), \quad y_t = y, \quad z_t = z. \quad (3.3)$$

These transformations we shall call x -transformations.

Using x -transformations, the Killing vectors $\partial/\partial y$ and $\partial/\partial z$ of the initial model remain Killing vectors. They form a 2-dimensional Abelian group of motions. All others, including the rotation $z(\partial/\partial y) - y(\partial/\partial z)$, may fail to remain Killing vectors. On the contrary to the general case, the x -transformed models depend genuinely on the initial ones. In the following, the spatially flat Friedmann universe with power-law behaviour of the cosmic scale factor shall be used as initial model, i.e.

$$g_{ab}(t) = \delta_{ab}K^2(t) \quad \text{with} \quad K(t) = t^\tau. \quad (3.4)$$

Together with eq. (3.1) one obtains from the Einstein field equation

$$\kappa\mu = \kappa T_{00} = 3\tau^2/t^2, \quad p = T_2^2 = \alpha\mu \quad \text{with} \quad \alpha = \frac{2}{3\tau} - 1; \quad (3.5)$$

where μ is the energy density and p is the pressure.

Now, inserting eq. (3.4) in eq. (3.1) and transforming to eq. (3.2) with restriction eq. (3.3) one obtains for the metric of the x -transformed model

$$g_{11} = t^{2\tau} \cdot (x_{t,1})^2 \equiv t^{2\tau} \cdot h(x, t), \quad g_{22} = g_{33} = t^{2\tau}, \quad g_{\alpha\beta} = \eta_{\alpha\beta} \text{ else.} \quad (3.6)$$

Here, $\eta_{\alpha\beta} = \text{diag}(-1, 1, 1, 1)$ is the metric of the flat Minkowski spacetime. This metric (3.6) belongs to the so-called Szekeres class, cf. [251]. Defining $a(x, t)$ by

$$g_{11} = e^{2a(x,t)}, \quad (3.7)$$

a coordinate transformation $\tilde{x}(x)$ yields $a(x, 1) = 0$. If $v = a_0 t$, then we have

$$a(x, t) = \int_1^t v(x, \tilde{t}) \cdot \tilde{t}^{-1} d\tilde{t}, \quad (3.8)$$

$v(x, t)$ being an arbitrary twice continuously differentiable function which may be singular at $t = 0$ and $t = \infty$. The initial model is included by setting $x_t = x$, hence $a = \tau \ln t$, $v = \tau$.

For $\tau \neq 0$ different functions v correspond to the same model only if they are connected by a translation into x -direction. Inserting eq. (3.6) with eq. (3.7) and eq. (3.8) into the Einstein field equation one obtains the energy-momentum tensor

$$\begin{aligned} \kappa T_{00} &= (\tau^2 + 2v\tau)/t^2, \\ \kappa T_1^1 &= (2\tau - 3\tau^2)/t^2, \\ \kappa T_2^2 = \kappa T_3^3 &= -v_0 t^{-1} + (\tau - \tau^2 - v\tau - v^2 + v)/t^2, \end{aligned}$$

$$\begin{aligned} \kappa T_{\alpha\beta} &= 0 \quad \text{else,} \quad \text{and} \\ \kappa T &= \kappa T_{\alpha}^{\alpha} = -2v_{,0}t^{-1} - 2(3\tau^2 - 2\tau + 2v\tau + v^2 - v)/t^2. \end{aligned} \quad (3.9)$$

The question, in which cases equation (3.6) gives a usual hypersurface-homogeneous model, can be answered as follows: metric (3.6) is a Friedmann universe, if and only if $h_{,0} = 0$. For this case it is isometric to the initial model. Metric (3.6) is a hypersurface-homogeneous model, if and only if functions A and B exist for which holds $h(x, t) = A(x) \cdot B(t)$. Because of $h > 0$ this is equivalent to $h_{,01} \cdot h = h_{,0} \cdot h_{,1}$. In this case it is a Bianchi type I model.

3.2 Energy inequalities and perfect fluid models

In this section it shall be discussed, in which manner the geometrically defined models described by eqs. (3.6), (3.7), (3.8) and (3.9) fulfil some energy conditions. Here we impose the following conditions: each observer measures non-negative energy density, time- or lightlike energy flow and spacelike tensions which are not greater than the energy density. In our coordinate system these conditions are expressed by the following inequalities

$$T_{00} \geq |T_1^1|, \quad (3.10)$$

$$T_{00} \geq |T_2^2|, \quad (3.11)$$

$$\text{and} \quad T \leq 0. \quad (3.12)$$

For the initial model this means $\tau = 0$ or $\tau \geq 1/2$, i.e. Minkowski spacetime or $-1 < \alpha \leq 1/3$. For the limiting case $\alpha = -1$ one gets the de Sitter spacetime in the original model (3.5) which does not have the assumed power-law behaviour for the scale factor. Using the energy-momentum tensor eq.

(3.9), eqs. (3.11) and (3.12) read

$$v_{,0} t \geq -v^2 - (3\tau - 1)v + \tau - 2\tau^2, \quad (3.13)$$

$$v_{,0} t \leq -v^2 + (\tau + 1)v + \tau \quad \text{and} \quad (3.14)$$

$$v_{,0} t \geq -v^2 - (2\tau - 1)v + 2\tau - 3\tau^2. \quad (3.15)$$

Now, if $\tau < 0$, i.e. $\alpha < -1$, then eq. (3.10) reads $v \leq \tau - 1$; together with eq. (3.14) one obtains $v_{,0} t \leq \tau - 2$. This implies the existence of a $\tilde{t} > 0$ with $v(\tilde{t}) \geq -1$ in contradiction to condition (3.10). Therefore, an initial model with $\tau < 0$, which itself contradicts the energy inequalities, cannot produce transformed models which always fulfil them.

If $\tau = 0$, then eqs. (3.13) and (3.14) imply $v_{,0} t = v - v^2$. This equation has the solutions $v = 0$ and $v = t(t + C)^{-1}$ with arbitrary $C(x) \geq 0$. This yields $a = 0$ and $a = \ln(t + C) - \ln(1 + C)$ with eq. (3.7), and $g_{11} = 1$ and $g_{11} = (t + C)^2 \cdot (1 + C)^{-2}$ resp. with eq. (3.8). Then eq. (3.6) shows that this is the Minkowski spacetime itself. Therefore, the Minkowski spacetime does not produce any new models.

Finally, if $\tau > 0$, hence $\alpha > -1$, eq. (3.10) then reads $v \geq \max(\tau - 1, 1 - 2\tau)$. A lengthy calculation shows which transformed models fulfil the energy inequalities. For each $\tau > 0$ models exist which do and models which do not fulfil them.

The situation described above changes if one requires that the transformed model consists of perfect fluid with an equation of state. The velocity vector must be $(1, 0, 0, 0)$ and

$$\kappa(T_1^1 - T_2^2) \equiv \frac{h_{,0}}{(1 + \alpha)t \cdot h} + \frac{1}{2\sqrt{h}} \left(\frac{h_{,0}}{\sqrt{h}} \right)_{,0} = 0 \quad (3.16)$$

must be fulfilled. Eq. (3.16) expresses the condition that the pressure has to be isotropic. $T_2^2 = T_3^3$ is fulfilled anyhow due to our assumption eq. (3.3).

If $f = h_0 h^{-1/2}$, then eq. (3.16) reads

$$\frac{f}{(1+\alpha)t} + \frac{f_{,0}}{2} = 0,$$

hence $f_{,0}f^{-1}$ does not depend on x , therefore $f_{,01} \cdot f = f_{,0} \cdot f_{,1}$, and we can use the ansatz $f = a(t) \cdot b(x)$. Inserting this into eq. (3.16), one obtains

$$h = \left[b(x) \cdot t^{(\alpha-1)(\alpha+1)} + c(x) \right]^2, \quad \text{where} \quad c(x) = \sqrt{h(x, 0)} \quad (3.17)$$

with arbitrary non-negative functions b and c fulfilling $b(x) + c(x) > 0$ for all x . For energy density and pressure we then obtain

$$\begin{aligned} \kappa\mu &= \frac{4}{3(1+\alpha)^2 t^2} - \frac{4(1-\alpha) \cdot b}{3(1+\alpha)^2 t(bt + ct^{2/(1+\alpha)})}, \\ \kappa p &= \frac{4\alpha}{3(1+\alpha)^2 t^2}. \end{aligned} \quad (3.18)$$

An equation of state means that p uniquely depends on μ . This takes place if and only if $\alpha = 0$ or $b/c = \text{const.}$ The latter is equivalent to the hypersurface-homogeneity of the model and is of lower interest here. For $\alpha = 0$ the initial model is the dust-filled Einstein-De Sitter model [68].

With eq. (3.18) we obtain $p = 0$ and $\mu \geq 0$ and may formulate: If the Einstein-De Sitter universe is x -transformed into a perfect fluid model, then this model also contains dust and fulfils the energy conditions.

These models have the following form: inserting eq. (3.17) with $\alpha = 0$ into eq. (3.6) we get a dust-filled model

$$ds^2 = -dt^2 + t^{4/3} \{ [b(x)/t + c(x)]^2 dx^2 + dy^2 + dz^2 \} \quad (3.19)$$

with arbitrary b, c as before. A subcase of eq. (3.19) is contained in Szekeres [251] as case (iii), but the parameter ε used there may now be non-constant. For $b = 1$, eq. (3.19) reduces to eq. (2.9). Of course, locally one can achieve a constant b by a transformation $\tilde{x} = \tilde{x}(x)$, but, e.g. changes in the sign of b then are not covered. As an illustration we give two examples of this model eq. (3.19):

1. If $b = 1, c > 0$, then

$$\kappa\mu = \frac{4c}{3t(1+ct)}, \quad (3.20)$$

hence the density contrast at two different values x_1, x_2 reads

$$\frac{\mu_1}{\mu_2} = \frac{c_2}{c_1} \cdot \frac{1+c_1t}{1+c_2t} \quad (3.21)$$

and tends to 1 as $t \rightarrow \infty$. This shows that one needs additional presumptions if one wants to prove an amplification of initial density fluctuations.

2. If $b = 1, c = 0$, then eq. (3.19) is the axially symmetric Kasner vacuum solution. If now c differs from zero in the neighbourhoods of two values x_1, x_2 , then the model is built up from two thin dust slices and Kasner-like vacuum outside them. The invariant distance of the slices is

$$At^{-1/3} + Bt^{2/3} \quad (3.22)$$

with certain positive constants A and B . In [225] it is argued that the invariant distance along spacelike geodesics need not coincide with what we measure as spatial distance. The distinction comes from the fact, that a geodesic within a prescribed spacelike hypersurface need not coincide with a spacelike geodesic in spacetime. This eq. (3.22) looks like gravitational repulsion, because the distance has a minimum at a positive value. But the

$t^{-1/3}$ -term is due to the participation of the slices in cosmological expansion and the remaining $t^{2/3}$ -term is due to an attractive gravitational force in parabolic motion. A more detailed discussion of this behaviour eq. (3.22) is given in Griffiths [99]: he argues, “however inhomogeneous the mass distribution, matter on each plane of symmetry has no net attraction to matter on other planes.”

The transformation of a hypersurface-homogeneous cosmological model considered here preserves all inner properties, expressed by the first fundamental form, of the slices $t = \text{const.}$, and also preserves the property that t is a synchronized time, but may change all other outer properties, which are expressed by the second fundamental form. The investigations of section 3.2 show that energy conditions are preserved under very special presumptions only.

One may consider these transformations as a guide in the search for new exact solutions of Einstein’s field equation. The new models are close to the initial hypersurface-homogeneous ones, if the transformation is close to the identical one. Thus, one can perturb a Bianchi model with exact solutions without use of any approximations.

3.3 A simple singularity theorem

What do we exactly know about solutions when no exact solution, in the sense of “solution in closed form”, is available? In which sense these solutions have a singularity? Here we make some remarks to these questions, see [228].

In [221], the following simple type of singularity theorems was discussed: The coordinates t, x, y, z shall cover all the reals, and $a(t)$ shall be an

arbitrary strictly positive monotonously increasing smooth function defined for all real values t , where “smooth” denotes “ C^∞ -differentiable”. Then it holds:

The Riemannian space defined by

$$ds^2 = dt^2 + a^2(t)(dx^2 + dy^2 + dz^2) \quad (3.23)$$

is geodesically complete. This fact is well-known and easy to prove; however, on the other hand, for the same class of functions $a(t)$ it holds:

The spacetime defined by

$$ds^2 = dt^2 - a^2(t)(dx^2 + dy^2 + dz^2) \quad (3.24)$$

is lightlike geodesically complete if and only if

$$\int_{-\infty}^0 a(t) dt = \infty \quad (3.25)$$

is fulfilled. The proof is straightforwardly done by considering lightlike geodesics in the $x - t$ -plane. So, one directly concludes that the statements do not depend on the number n of spatial dimensions, only $n > 0$ is used, and the formulation for $n = 3$ was chosen as the most interesting case. Moreover, this statement remains valid if we replace “lightlike geodesically complete” by “lightlike and timelike geodesically complete”.

Further recent results on singularity theorems are reviewed in Senovilla [233]. This review and most of the research concentrated mainly on the 4-dimensional Einstein theory. Probably, the majority of arguments can be taken over to the higher-dimensional Einstein equation without change, but this has not yet worked out up to now. And singularity theorems for $F(R)$ -gravity are known up to now only for very special cases only.

The most frequently used $F(R)$ -Lagrangian is

$$L = \left(\frac{R}{2} - \frac{l^2}{12} R^2 \right) \sqrt{-g} \quad \text{where } l > 0, \quad (3.26)$$

here we discuss the non-tachyonic case only. From this Lagrangian one gets a fourth-order field equation; only very few closed-form solutions, “exact solutions”, are known. However, for the class of spatially flat Friedmann models, the set of solutions is qualitatively completely described, but not in closed form, in [163]. We call them “rigorous solutions”.

One of them, often called “Starobinsky inflation”, can be approximated with

$$a(t) = \exp\left(-\frac{t^2}{12l^2}\right) \quad (3.27)$$

This approximation is valid in the region $t \ll -l$. However, this solution does not fulfil the above integral condition. Therefore, Starobinsky inflation does not represent a lightlike geodesically complete cosmological model as has been frequently stated in the literature.

To prevent a further misinterpretation let me reformulate this result as follows: In spite of the fact that the Starobinsky model is regular, in the sense that $a(t) > 0$ for arbitrary values of synchronized time t , every past-directed lightlike geodesic terminates in a curvature singularity, i.e., $|R| \rightarrow \infty$, at a finite value of its affine parameter. Therefore, the model is not only geodesically incomplete in the coordinates chosen, moreover, it also fails to be a subspace of a complete one.

In contrast to this one can say: the spatially flat Friedmann model with $a(t) = \exp(Ht)$, H being a positive constant, is the inflationary de Sitter spacetime. According to the above integral condition, it is also incomplete.

However, contrary to the Starobinsky model, it is a subspace of a complete spacetime, namely of the de Sitter spacetime represented as a closed Friedmann model.

The application of conformal transformations represents one of the most powerful methods to transform different theories into each other. In many cases this related theories to each other which have been originally considered to be independent ones. The most often discussed question in this context is “which of these metrics is the physical one”, but one can, of course, use such conformal relations also as a simple mathematical tool to find exact solutions without the necessity of answering this question. However, concerning a conformal transformation of singularity theorems one must be cautious, because typically, near a singularity in one of the theories, the conformal factor diverges, and then the conformally transformed metric need not be singular there.

Finally, let me again mention the distinct properties of the Euclidean and the Lorentzian signature: Smooth connected complete Riemannian spaces are geodesically connected. However, smooth connected complete spacetimes, e.g. the de Sitter space–time, need not be geodesically connected, see for instance Sanchez [196]. The topological origin of this distinction is the same as the distinction between the two above statements: for the positive definite signature case one uses the compactness of the rotation group, whereas in the indefinite case the noncompactness of the Lorentz group has to be observed [227].

4 Properties of curvature invariants

In section 4.1, which is an extended version of [223] we answer the following question: Let λ, μ, ν be arbitrary real numbers. Does there exist a 3-dimensional homogeneous Riemannian manifold whose eigenvalues of the Ricci tensor are just λ, μ and ν ?

In section 4.2 we discuss, following [224], why all the curvature invariants of a gravitational wave vanish.

In section 4.3 we present the example [222] that it may be possible that non-isometric spacetimes with non-vanishing curvature scalar cannot be distinguished by curvature invariants.

4.1 The space of 3-dimensional homogeneous Riemannian manifolds

The curvature of a 3-manifold is completely determined by the Ricci tensor, and the Ricci tensor of a homogeneous manifold has constant eigenvalues. However, it is essential to observe, that, nevertheless, the constancy of all three eigenvalues of the Ricci tensor of a 3-manifold does not imply the manifold to be locally homogeneous, see [132] for examples.

Assume λ, μ, ν to be arbitrary real numbers. First question: Does there exist a 3-dimensional homogeneous Riemannian manifold whose eigenvalues of the Ricci tensor are just λ, μ and ν ? Second question: If it exists, is it uniquely determined, at least locally up to isometries? The answer to these questions does not alter if we replace “homogeneous” by “locally homogeneous”. In principle, they could have been answered by the year 1905

already. But in fact, it was not given until 1995: the first of these problems have been solved independently by Kowalski and Nikcevic [133] on the one hand, and by Rainer and Schmidt [189], on the other hand; the second one is solved several times, e.g. in [189]. Curiously enough, the expected answer “Yes” turns out to be wrong.

Of course, both answers [133] and [189] are equivalent. But the formulations are so different, that it is useful to compare them here. In [133] one defines

$$\rho_1 = \max\{\lambda, \mu, \nu\}, \quad \rho_3 = \min\{\lambda, \mu, \nu\} \quad (4.1)$$

and

$$\rho_2 = \lambda + \mu + \nu - \rho_1 - \rho_3 \quad (4.2)$$

and formulates the conditions as inequalities between the ρ_i .

In [189] the formulation uses curvature invariants. First step: one observes that the answer does not alter, if one multiplies λ, μ, ν with the same positive constant. Note: this fails to be the case if we take a negative constant. Second step: we take curvature invariants which do not alter by this multiplication, i.e. we look for suitable homothetic invariants. Third step: we formulate the answer by use of the homothetic invariants.

First step: this can be achieved by a homothetic transformation of the metric, i.e., by a conformal transformation with constant conformal factor. Then one excludes the trivial case $\lambda = \mu = \nu = 0$. Note: It is due to the positive definiteness of the metric that $\lambda = \mu = \nu = 0$ implies the flatness of the space.

Second step:

$$R = \lambda + \mu + \nu, \quad N = \lambda^2 + \mu^2 + \nu^2 \quad (4.3)$$

are two curvature invariants with $N > 0$, and

$$\hat{R} = R/\sqrt{N} \quad (4.4)$$

is a homothetic invariant. For defining a second homothetic invariant we introduce the trace-free part of the Ricci tensor

$$S_{ij} = R_{ij} - \frac{R}{3}g_{ij} \quad (4.5)$$

and its invariant

$$S = S_i^j S_j^k S_k^i. \quad (4.6)$$

The desired second homothetic invariant reads

$$\hat{S} = S/\sqrt{N^3}. \quad (4.7)$$

Of course, one can also directly define $\hat{R}$ and $\hat{S}$ as functions of λ , μ and ν , then S is the cubic polynomial,

$$S = \left((2\lambda - \mu - \nu)^3 + (2\mu - \nu - \lambda)^3 + (2\nu - \lambda - \mu)^3 \right) / 27, \quad (4.8)$$

see [189] for details. Remark: the inequalities $S > 0$ and $\rho_2 < (\rho_1 + \rho_3)/2$ are equivalent.

Third step: The answer is “yes” if and only if one of the following conditions (4.9) till (4.14) are fulfilled:

$$N = 0, \quad \text{flat space} \quad (4.9)$$

$$\hat{R}^2 = 3 \quad \text{and} \quad \hat{S} = 0, \quad \text{non-flat spaces of constant curvature} \quad (4.10)$$

$$\hat{R}^2 = 2 \quad \text{and} \quad 9\hat{R}\hat{S} = -1, \quad \text{real line times constant curvature space} \quad (4.11)$$

$$\hat{R}^2 = 1 \quad \text{and} \quad 9\hat{R}\hat{S} = 2 \quad (4.12)$$

$$18\hat{S} > \hat{R}(9 - 5\hat{R}^2), \quad \text{equivalent to} \quad \lambda\mu\nu > 0 \quad (4.13)$$

$$-\sqrt{3} < \hat{R} < -\sqrt{2} \quad \text{and} \quad 0 < \hat{S} \leq \left(3 - \hat{R}^2\right)^2 \frac{6 - 5\hat{R}^2}{18\hat{R}^3}. \quad (4.14)$$

Remarks: 1. In [133] the last of the four inequalities (4.14) is more elegantly written as

$$\rho_2 \geq \frac{\rho_1^2 + \rho_3^2}{\rho_1 + \rho_3}. \quad (4.15)$$

2. There exists a one-parameter set of Bianchi-type VI manifolds, with parameter a where $0 < a < 1$, having $\rho_1 = -a - a^2$, $\rho_2 = -1 - a^2$, $\rho_3 = -1 - a$; they represent the case where in (4.14) the fourth inequality is fulfilled with “=”.

Now we turn to the second question: Let the 3 numbers λ, μ, ν be given such that a homogeneous 3-space exists having these 3 numbers as its eigenvalues of the Ricci tensor; is this 3-space locally uniquely determined? The first part of the answer is trivial:

If $\lambda = \mu = \nu$, then “yes”, it is the space of constant curvature. The second part of the answer is only slightly more involved: If λ, μ and ν represent 3 different numbers, then the 3 eigendirections of the Ricci tensor are uniquely determined, and then we can also prove: “yes”. The third part is really surprising: If

$$\hat{R} = 1 \quad \text{and} \quad \hat{S} = \frac{2}{9} \quad (4.16)$$

then the answer is “no”, otherwise “yes”.

Clearly, eq. (4.16) represents a subcase of (4.12). What is the peculiarity with the values (4.16) of the homothetic invariants? They belong to Bianchi type IX, have one single and one double eigenvalue, and with these values, a one-parameter set of examples with 3-dimensional isometry group, and another example with a 4-dimensional isometry group exist.

4.2 Why do all the curvature invariants of a gravitational wave vanish?

We prove the theorem valid for Pseudo-Riemannian manifolds V_n : “Let $x \in V_n$ be a fixed point of a homothetic motion which is not an isometry then all curvature invariants vanish at x .” and get the Corollary: “All curvature invariants of the plane wave metric

$$ds^2 = 2 du dv + a^2(u) dw^2 + b^2(u) dz^2 \quad (4.17)$$

identically vanish.”

Analysing the proof we see: The fact that for definite signature flatness can be characterized by the vanishing of a curvature invariant, essentially rests on the compactness of the rotation group $SO(n)$. For Lorentz signature, however, one has the non-compact Lorentz group $SO(3, 1)$ instead of it.

A further and independent proof of the corollary uses the fact, that the Geroch limit does not lead to a Hausdorff topology, so a sequence of gravitational waves can converge to the flat spacetime, even if each element of the sequence is the same pp-wave.

The energy of the gravitational field, especially of gravitational waves, within General Relativity was subject of controversies from the very beginning, see Einstein [66]. Global considerations - e.g. by considering the far-field of asymptotically flat spacetimes - soon led to satisfactory answers. Local considerations became fruitful if a system of reference is prescribed e.g. by choosing a timelike vector field. If, however, no system of reference is preferred then it is not a priori clear whether one can constructively distinguish

flat spacetime from a gravitational wave. This is connected with the generally known fact, that for a pp-wave, see e.g. Stephani [250] especially section 15.3. and [65] all curvature invariants vanish, cf. Hawking and Ellis [107] and Jordan et al. [123], but on the other hand: in the absence of matter or reference systems - only curvature invariants are locally constructively measurable. See also the sentence “R. Penrose has pointed out, in plane-wave solutions the scalar polynomials are all zero but the Riemann tensor does not vanish.” taken from page 260 of [107]. At page 97 of [123] it is mentioned that for a pp-wave all curvature invariants constructed from

$$R_{ijkl;i_1\dots i_r} \tag{4.18}$$

by products and traces do vanish.

It is the aim of this section to explain the topological origin of this strange property.

4.2.1 Preliminaries

Let V_n be a C^∞ -Pseudo-Riemannian manifold of arbitrary signature with dimension $n > 1$. The metric and the Riemann tensor have components g_{ij} and R_{ijlm} resp. The covariant derivative with respect to the coordinate x^m is denoted by “; m ” and is performed with the Christoffel affinity Γ_{lm}^i . We define

I is called a generalized curvature invariant of order k if it is a scalar with dependence

$$I = I(g_{ij}, R_{ijlm}, \dots, R_{ijlm;i_1\dots i_k}). \tag{4.19}$$

By specialization we get the usual Definition: I is called a curvature invariant of order k if it is a generalized curvature invariant of order k which depends continuously on all its arguments. The domain of dependence is requested to contain the flat space, and

$$I(g_{ij}, 0, \dots, 0) \equiv 0. \quad (4.20)$$

Examples: Let

$$I_0 = \text{sign}\left(\sum_{i,j,l,m=1}^n |R_{ijlm}|\right), \quad (4.21)$$

where the sign function is defined by $\text{sign}(0) = 0$ and $\text{sign}(x) = 1$ for $x > 0$. Then I_0 is a generalized curvature invariant of order 0, but it fails to be a curvature invariant. It holds: V_n is flat if and only if $I_0 \equiv 0$. Let further

$$I_1 = R_{ijlm} R^{ijlm} \quad (4.22)$$

which is a curvature invariant of order 0. If the metric has definite signature or if $n = 2$ then it holds: V_n is flat if and only if $I_1 \equiv 0$. Proof: For definite signature $I_0 = \text{sign}(I_1)$; for $n = 2$, $I_1 \equiv 0$ implies $R \equiv 0$, hence flatness. q.e.d.

For all other cases, however, the vanishing of I_1 does not imply flatness. Moreover, there does not exist another curvature invariant serving for this purpose, it holds:

For dimension $n \geq 3$, arbitrary order k and indefinite metric it holds: To each curvature invariant I of order k there exists a non-flat V_n with $I \equiv 0$.

Proof: Let $n = 3$. We use

$$ds^2 = 2 du dv \pm a^2(u) dw^2 \quad (4.23)$$

with a positive non-linear function $a(u)$. The “ $\pm$ ” covers the two possible indefinite signatures for $n = 3$. The Ricci tensor is $R_{ij} = R^m{}_{imj}$ and has with $u = x^1$

$$R_{11} = -\frac{1}{a} \cdot \frac{d^2 a}{du^2} \quad (4.24)$$

and therefore, eq. (4.23) represents a non-flat metric. Now let $n > 3$. We use the Cartesian product of eq. (4.23) with a flat space of dimension $n - 3$ and arbitrary signature. So we have for each $n \geq 3$ and each indefinite signature an example of a non-flat V_n . It remains to show that for all these examples, all curvature invariants of order k vanish. It suffices to prove that at the origin of the coordinate system, because at all other points it can be shown by translations of all coordinates accompanied by a redefinition of $a(u)$ to $a(u - u_0)$. Let I be a curvature invariant of order k . Independent of the dimension, i.e., how many flat spaces are multiplied to metric eq. (4.23), one gets for the case considered here that

$$I = I(a^{(0)}(u), a^{(1)}(u), \dots, a^{(k+2)}(u)) \quad (4.25)$$

where $a^{(0)}(u) = a(u)$, $a^{(m+1)}(u) = \frac{d}{du}a^{(m)}(u)$, and

$$I(a^{(0)}(u), 0, \dots, 0) = 0. \quad (4.26)$$

This is valid because each

$$R_{ijlm;i_1 \dots i_p} \quad (4.27)$$

continuously depends on $a^{(0)}(u)$, $a^{(1)}(u)$, $\dots$, $a^{(p+2)}(u)$ and on nothing else; and for $a = \text{const.}$, metric (4.23) represents a flat space.

Now we apply a coordinate transformation: Let $\varepsilon > 0$ be fixed, we replace u by $u \cdot \varepsilon$ and v by v/ε . This represents a Lorentz boost in the $u - v$ -plane.

Metric eq. (4.23) remains form-invariant by this rotation, only $a(u)$ has to be replaced by $a(u \cdot \varepsilon)$. At $u = 0$ we have

$$I = I(a^{(0)}(0), a^{(1)}(0), \dots, a^{(k+2)}(0)) \quad (4.28)$$

which must be equal to

$$I_\varepsilon = I(a^{(0)}(0), \varepsilon \cdot a^{(1)}(0), \dots, \varepsilon^{k+2} \cdot a^{(k+2)}(0)) \quad (4.29)$$

because I is a scalar. By continuity and by the fact that flat space belongs to the domain of dependence of I , we have

$$\lim_{\varepsilon \rightarrow 0} I_\varepsilon = 0. \quad (4.30)$$

All values I_ε with $\varepsilon > 0$ coincide, and so $I = 0$. q.e.d.

4.2.2 Gravitational waves

New results on gravitational waves can be found in [72], and [84]. A pp-wave is a plane-fronted gravitational wave with parallel rays, see [77]. It is a non-flat solution of Einstein's vacuum equation $R_{ij} = 0$ possessing a non-vanishing covariantly constant vector; this vector is then automatically a null vector. The simplest type of pp-waves can be represented similar as metric eq. (4.23)

$$ds^2 = 2 du dv + a^2(u) dw^2 + b^2(u) dz^2, \quad (4.31)$$

where

$$b \cdot \frac{d^2 a}{du^2} + a \cdot \frac{d^2 b}{du^2} = 0. \quad (4.32)$$

This metric represents flat spacetime if and only if both a and b are linear functions. Using the arguments of subsection 4.2.1 one sees that all curvature invariants this metric identically vanish. Here we present a second proof of

that statement which has the advantage to put the problem into a more general framework and to increase the class of spacetimes covered, e.g. to the waves

$$ds^2 = dx^2 + dy^2 + 2du dv + H(x, y, u)du^2 \quad (4.33)$$

Hall [102] considers fixed points of homothetic motions, which cannot exist in compact spacetimes, and shows that any plane wave, not only vacuum plane waves, admits, for each x , a homothetic vector field which vanishes at x . It holds

Let $x \in V_n$ be a fixed point of a homothetic motion which is not an isometry then all curvature invariants vanish at x .

Proof: The existence of a homothetic motion which is not an isometry means that V_n is selfsimilar. Let the underlying differentiable manifold be equipped with two metrics g_{ij} and

$$\tilde{g}_{ij} = e^{2C} g_{ij} \quad (4.34)$$

where C is a non-vanishing constant. The corresponding Riemannian manifolds are denoted by V_n and $\tilde{V}_n$ resp. By assumption, there exists an isometry from V_n to $\tilde{V}_n$ leaving x fixed. Let I be a curvature invariant. I can be represented as continuous function which vanishes if all the arguments do of finitely many of the elementary invariants. The elementary invariants are such products of factors g^{ij} with factors of type

$$R_{ijlm;i_1 \dots i_p} \quad (4.35)$$

which lead to a scalar, i.e., all indices are traced out. Let J be such an elementary invariant. By construction we have $J(x) = e^{qC} J(x)$ with a non-vanishing natural q , which depends on the type of J . Therefore, $J(x) = 0$. q.e.d.

From this it follows: All curvature invariants of metric (4.17) identically vanish. This statement refers not only to the 14 independent elementary invariants of order 0, see Harvey [104] and Lake 1993 [140] for a list of them, but for arbitrary order.

Proof: We have to show that for each point x , there exists a homothetic motion with fixed point x which is not an isometry. But this is trivially done by suitable linear coordinate transformations of v , w , and z . q.e.d.

4.2.3 Topological properties

Sometimes it is discussed that the properties of spacetime which can be locally and constructively, i.e., by rods and clocks, measured are not only the curvature invariants but primarily the projections of the curvature tensor and its covariant derivatives to an orthonormal tetrad, called 4-bein.⁶ The continuity presumption expresses the fact that a small deformation of spacetime should also lead to a correspondingly small change of the result of the measurement. To prevent a preferred system of reference one can construct curvature invariants like

$$I_2 = \inf \sum_{i,j,l,m} |R_{ijlm}| \quad (4.36)$$

where the infimum, here the same as the minimum, is taken over all orthonormal tetrads. From the first glance one could believe that $I_2 \equiv 0$ if and only if the space is flat. But for indefinite signature this would contradict the result of subsection 4.2.1. What is the reason? For definite signature the infimum is to be taken about the rotation group $SO(4)$, or $O(4)$ if one allows

⁶The German word “das Bein” denotes both “bone” and “leg”, here it is used in the second sense.

orientation-reversing systems; this group is compact. One knows: A positive continuous function over a compactum possesses a positive infimum. So, if one of the R_{ijlm} differs from zero, then $I_2 > 0$ at that point. For Lorentz signature, however, the infimum is to be taken about the non-compact Lorentz group $SO(3, 1)$ and so $R_{ijlm} \neq 0$ does not imply $I_2 \neq 0$.

Another topological argument, which underlies our subsection 4.2.1, is connected with the Geroch limit of spacetimes [86], we use the version of [210]. Theorem 3.1 of reference [210] reads:

1: For local Riemannian manifolds with definite signature, Geroch's limit defines a Hausdorff topology.

2: For indefinite signature this topology is not even T_1 .

Explanation: A topology is Hausdorff if each generalized Moore - Smith sequence possesses at most one limit, and it is T_1 if each constant sequence possesses at most one limit. The main example is a sequence, where each element of the sequence is the same pp-wave, and the sequence possesses two limits: flat spacetime and that pp-wave. Here the reason is: Only for definite signature, geodetic ε -balls form a neighbourhood basis for the topology.

Final remarks: The change from Euclidean to Lorentzian signature of a Pseudo-Riemannian space is much more than a purely algebraic duality - an impression which is sometimes given by writing an imaginary time coordinate: One loses all the nice properties which follow from the compactness of the rotation group. Lake 1993 [140] pointed out that also for the Robinson-Trautman vacuum solutions of Petrov type III all 14 curvature invariants vanish.

4.3 On spacetimes which cannot be distinguished by curvature invariants

For a positive C^∞ -function $a(u)$ let

$$ds^2 = \frac{1}{z^2} [2 du dv - a^2(u) dy^2 - dz^2] . \quad (4.37)$$

In the region $z > 0$, ds^2 represents the anti-de Sitter spacetime if and only if $a(u)$ is linear in u . Now, let $d^2a/du^2 < 0$ and

$$\phi := \frac{1}{\sqrt{\kappa}} \int \left(-\frac{1}{a} \frac{d^2a}{du^2} \right)^{1/2} du . \quad (4.38)$$

Then

$$\square \phi = \phi_{,i} \phi^{,i} = 0 \quad (4.39)$$

and

$$R_{ij} - \frac{R}{2} g_{ij} = \Lambda g_{ij} + \kappa T_{ij} \quad (4.40)$$

with $\Lambda = -3$ and $T_{ij} = \phi_{,i} \phi_{,j}$. So (ds^2, ϕ) represents a solution of Einstein's equation with negative cosmological constant Λ and a minimally coupled massless scalar field ϕ .

Let I be a curvature invariant of order k , i.e., I is a scalar

$$I = I(g_{ij}, R_{ijlm}, \dots, R_{ijlm;i_1 \dots i_k}) \quad (4.41)$$

depending continuously on all its arguments. Then for the metric ds^2 , I does not depend on the function $a(u)$.

This seems to be the first example that non-isometric spacetimes with non-vanishing curvature scalar cannot be distinguished by curvature invariants. The proof essentially uses the non-compactness of the Lorentz group

$SO(3, 1)$, here of the boosts $u \rightarrow u \lambda$, $v \rightarrow v/\lambda$. One can see this also in the representation theory in comparison with the representations of the compact rotation group $SO(4)$.

Let $v \in R^4$ be a vector and $g \in SO(4)$ such that $g(v) \uparrow\uparrow v$, then $g(v) = v$. In Minkowski spacetime M^4 , however, there exist vectors $v \in M^4$ and a $h \in SO(3, 1)$ with $h(v) \uparrow\uparrow v$ and $h(v) \neq v$. This is the reason why the null frame components of the curvature tensor called Cartan “scalars” are not always curvature scalars, see [224]. Further new results on curvature invariants are given in [28], [184] and [188]. Limits of spacetimes are considered in [176] and [230].

5 Surface layers and relativistic surface tensions

For a thin shell, the intrinsic 3-pressure will be shown to be analogous to $-A$, where A is the classical surface tension: First, interior and exterior Schwarzschild solutions will be matched together such that the surface layer generated at the common boundary has no gravitational mass; then its intrinsic 3-pressure represents a surface tension fulfilling Kelvin's relation between mean curvature and pressure difference in the Newtonian limit. Second, after a suitable definition of mean curvature, the general relativistic analogue to Kelvin's relation will be proven to be contained in the equation of motion of the surface layer.

5.1 Introduction to surface layers

In general relativity, an energy-momentum tensor concentrated on a timelike hypersurface is called a surface layer. Via Einstein's equations it is related to non-spurious jumps of the Christoffel affinities or equivalently to jumps of the second fundamental tensor. In [141], [178], [60], [117], [177] and [138] there have been given algorithms for their calculating, and in [141], [117], and [38], [63], [202], [260], [79] and [140] the spherically symmetric case was of a special interest. The discussion of disklike layers as models for accretion disks was initiated in [160] and [264], and [143] contains surface layers as sources of the Kerr geometry.

A surface layer has no component in the normal direction, otherwise the delta-like character would be destroyed. Hence, an ideal fluid with non-vanishing pressure cannot be concentrated on an arbitrarily thin region and

therefore in most cases, the layer is considered to be composed of dust. Additionally, an intrinsic 3-pressure is taken into account in [67], where for a spherically symmetric configuration each shell $r = \text{const.}$ is thought to be composed of identical particles moving on circular orbits without a preferred direction, and the tangential pressure, which is analogous to the intrinsic 3-pressure, is due to particle collisions, whereas a radial pressure does not appear. See also [202], [140], and [151], where the equation of motion for a spherically symmetric layer has been discussed.

Such an intrinsic 3-pressure as well as surface tensions are both of the physical dimension “force per unit length.” There the question arises whether an intrinsic 3-pressure of a surface layer may be related to a surface tension, and this chapter will deal with just this question. Then a general relativistic formulation of thermodynamics can be completed by equations for surface tensions to answer, e.g., the question how long a drop, say, a liquid comet, remains connected while falling towards a compact object. The only paper concerned with such questions seems to be [135]. There the influence of surface tensions on the propagation of gravitational waves has been calculated by perturbation methods, yielding a possibly measurable effect. For the background knowledge cf. [118] or [170] for relativistic thermodynamics and [174] for non-relativistic surface tensions.

This chapter, which is based on [203], proceeds as follows: Section 5.2 contains Kelvin’s relation for non-relativistic surface tensions and discusses the matching of the interior to the exterior Schwarzschild solution such that a surface tension appears, and compares with Kelvin’s relation in the Newtonian limit, calculations are found in the appendices, sections 5.5, 5.5 and 5.5. Sections 5.3 and 5.4 are devoted to the non-spherically symmetric case.

Section 5.3 contains a suitable definition of mean curvature and section 5.4 deduces Kelvin's relation from the equation of motion of the surface layer without any weak field assumptions.

5.2 Non-relativistic surface tensions

Imagine a drop of some liquid moving in vacuo. Its equilibrium configuration is a spherical one, and Kelvin's relation [255] between surface tension A , pressure difference ΔP , i.e. outer minus inner pressure, and mean curvature H , where $H = 1/R$ for a sphere of radius R , reads

$$\Delta P = -2HA. \quad (5.1)$$

A is a material-dependent constant. Equation (5.1) means, an energy $A \cdot \Delta F$ is needed to increase the surface area by ΔF . This supports our description of $-A$ as a kind of intrinsic pressure. But of course, it is a quite different physical process: Pressure, say, of an ideal gas, can be explained by collisions of freely moving particles, whereas the microphysical explanation of surface tensions requires the determination of the intermolecular potential, which looks like

$$\Phi(r) = -\mu r^{-6} + Ne^{-r/\rho} \quad (5.2)$$

with certain constants μ , N , and ρ ; see Ono [174]. In this context the surface has a thickness of about 10^{-7} cm but in most cases this thickness may be neglected.

In addition, for a non-spherically symmetric surface, the mean curvature H in equation eq. (5.1) may be obtained from the principal curvature radii R_1 , R_2 by means of the relation

$$H = \frac{1}{2R_1} + \frac{1}{2R_2}. \quad (5.3)$$

Now the drop shall be composed of an incompressible liquid. For a general relativistic description we have to take the interior Schwarzschild solution

$$ds^2 = - \left[\frac{3}{2}(1 - r_g/r_0)^{1/2} - \frac{1}{2}(1 - r_g r^2/r_0^3)^{1/2} \right]^2 dt^2 + \frac{dr^2}{1 - r_g r^2/r_0^3} + r^2 d\Omega^2 \quad \text{where} \quad d\Omega^2 = d\psi^2 + \sin^2 \psi d\varphi^2, \quad (5.4)$$

whose energy-momentum tensor represents ideal fluid with energy density $\mu = 3r_g/\kappa r_0^3$ and pressure

$$p(r) = \mu \cdot \frac{(1 - r_g r^2/r_0^3)^{1/2} - (1 - r_g/r_0)^{1/2}}{3(1 - r_g/r_0)^{1/2} - (1 - r_g r^2/r_0^3)^{1/2}}. \quad (5.5)$$

One has $p(r_0) = 0$, and therefore usually $0 \leq r \leq r_0$ is considered. But now we require only a non-vanishing pressure at the inner surface and take eq. (5.4) for values r with $0 \leq r \leq R$ and a fixed $R < r_0$ only.

The gravitational mass of the inner region equals

$$M = \mu \cdot 4\pi R^3/3 = r_g(R/r_0)^3/2 \quad (5.6)$$

where we assumed unit such that $G = c = 1$. The outer region shall be empty and therefore we have to insert the Schwarzschild solution for $r \geq R$. Neglecting the gravitational mass of the boundary $\Sigma \subset V_4$ which is defined by $r - R = 0$, just M of equation (5.6) has to be used as the mass parameter of the exterior Schwarzschild metric. Then only delta-like tensions appear at Σ , and $\Delta P = -p(R) < 0$. The vanishing of the gravitational mass is required to single out the properties of an intrinsic 3-pressure. In general, a surface layer is composed of both parts, not at least to ensure the validity of the energy condition $T_{00} \geq |T_{ik}|$ which holds for all known types of matter.

On Σ , the energy-momentum tensor is

$$T_i^k = \tau_i^k \cdot \delta_\Sigma \quad (5.7)$$

the non-vanishing components of which are

$$\tau_2^2 = \tau_3^3 = -p(R)R/2(1 - 2M/R)^{1/2} \quad (5.8)$$

cf. section 5.5.

The δ_Σ distribution is defined such that for all smooth scalar functions f the invariant integrals are related by

$$\int_\Sigma f d^{(3)}x = \int_{V_4} f \cdot \delta_\Sigma d^{(4)}x. \quad (5.9)$$

For a more detailed discussion of distribution-valued tensors in curved spacetime, cf. [252]. The fact that all components τ_0^α in eq. (5.7) vanish reflects the non-existence of a delta-like gravitational mass on Σ .

Now consider the Newtonian limit $M/R \ll 1$; then the mean curvature becomes again $H = 1/R$, and together with eq. (5.8) Kelvin's relation eq. (5.1) is just equivalent to

$$\tau_2^2 = \tau_3^3 = -A[1 + O(M/R)]. \quad (5.10)$$

Therefore: At least for static spherically symmetric configurations and weak fields a delta-like negative tangential pressure coincides with the classical surface tension.

In the next two sections we investigate to what extent these presumptions are necessary.

5.3 Mean curvature in curved spacetime

To obtain a general relativistic analogue to Kelvin's relation eq. (5.1) we have to define the mean curvature H of the timelike hypersurface Σ contained in a spacetime V_4 such that for weak fields just the usual mean curvature arises.

To get the configuration we have in mind, we make the ansatz

$$\tau_{\alpha\beta} = -A(g_{\alpha\beta} + u_\alpha u_\beta) \quad u_\alpha u^\alpha = -1 \quad (5.11)$$

with $A > 0$. Thereby again the gravitational mass of Σ will be neglected. Now the mean curvature shall be defined. But there is a problem: In general, there does not exist a surface $S \subset \Sigma$ which can serve as “boundary at a fixed moment” for which we are to determine the mean curvature. To circumvent this problem we start considering the special case

$$u_{\alpha\parallel\beta} = u_{\beta\parallel\alpha}. \quad (5.12)$$

Then there exists a scalar t on Σ such that $u_\alpha = t_{\parallel\alpha}$, and the surface $S \subset \Sigma$ defined by $t = 0$ may be called “boundary at a fixed moment.” Now S has to be embedded into a “space at a fixed moment”: We take intervals of geodesics starting from points of S in the normal direction n_i and the opposite one.

The union V_3 of these geodetic segments will be called “space at a fixed moment,” and $S \subset V_3$ is simply a two-surface in a three-dimensional positive definite Riemannian manifold, for which mean curvature has a definite sense: Let v^α , w^α be the principal curvature directions inside S and R_1 , R_2 the corresponding principal curvature radii, then equation (5.3) applies to obtain H .

Of course, $v^\alpha w_\alpha = 0$ holds, and $v^\alpha v_\alpha = w^\alpha w_\alpha = 1$ shall be attained. Then, inserting the second fundamental tensor cf. section 5.5, this becomes equivalent to

$$H = \frac{1}{2}(v^\alpha v_\alpha + w^\alpha w_\alpha)k_{\alpha\beta} = \frac{1}{2}(g^{\alpha\beta} + u^\alpha u^\beta)k_{\alpha\beta}. \quad (5.13)$$

But this latter relation makes sense without any reference to condition eq. (5.12). Therefore, we define eq. (5.13) to be the general relativistic analogue

to the mean curvature of a surface. For the case $H^+ \neq H^-$ we take their arithmetic mean

$$H = (H^+ + H^-)/2, \quad (5.14)$$

and then we obtain from eqs. (5.11) and (5.13)

$$-2HA = \frac{1}{2} (k_{\alpha\beta}^+ + k_{\alpha\beta}^-) \tau^{\alpha\beta}. \quad (5.15)$$

This choice can be accepted noting that in the Newtonian limit

$$|H^+ - H^-| \ll |H^+ + H^-| \quad (5.16)$$

anyhow. To compare this with Kelvin's relation we have to relate the right-hand side of equation (5.15) to the pressure difference ΔP at Σ . To this end we investigate the equation of motion for the surface layer.

5.4 Equation of motion for the surface layer

The equation of motion, $T_{i;k}^k = 0$, contains products of δ distributions and θ -step functions, where $\theta(x) = 1$ for $x \geq 0$, and $\theta(x) = 0$ else, at points where Γ_{jk}^i has a jump discontinuity. These products require special care; cf. [53] for a discussion of his point. But defining $\theta \cdot \delta = \frac{1}{2}\delta$ we obtain, cf. section 5.5

$$\Delta P \equiv \Delta n_i n^k T_k^i = \frac{1}{2} (k_{\alpha\beta}^+ + k_{\alpha\beta}^-) \tau^{\alpha\beta} \quad \text{and} \quad (5.17)$$

$$\Delta T_\alpha^1 \equiv \Delta n_i e_\alpha^k T_k^i = -\tau_{\alpha\parallel\beta}^\beta. \quad (5.18)$$

From equation (5.18) we see the following: The equation $\tau_{\alpha\parallel\beta}^\beta = 0$ holds only under the additional presumption that the regular, i.e., not delta-like, part of T_α^1 has no jump on Σ .

This condition is fulfilled e.g., presuming Σ to be such a boundary that the regular energy flow does not cross it and the four-velocity is parallel to Σ in both V_+ and V_- . This we will presume in the following. Then ΔP is indeed the difference of the pressures on both sides, and together with equations (5.15) and (5.17) we obtain exactly Kelvin's relation eq. (5.1). That means, it is the definition of mean curvature used here that enables us to generalize Kelvin's formula to general relativity. Furthermore, $O(M/R)$ of equation (5.10) vanishes.

Finally we want to discuss the equation $\tau_{\alpha\parallel\beta}^\beta = 0$. Transvection with u^α and $\delta_\gamma^\alpha + u^\alpha u_\gamma$ yields

$$u_{\parallel\alpha}^\alpha = 0 \quad \text{and} \quad u^\alpha u_{\gamma\parallel\alpha} + (\ln A)_{\parallel\alpha} (\delta_\gamma^\alpha + u^\alpha u_\gamma) = 0 \quad (5.19)$$

respectively. But A is a constant here, and therefore u^α is an expansion-free geodesic vector field in Σ . But observe that the u^α lines are geodesics in V_4 under additional presumptions only.

Here, we have only considered a phenomenological theory of surface tensions, and, of course, a more detailed theory has to include intermolecular forces. But on that phenomenological level equations (5.11) and (5.19) together with Kelvin's relation eq. (5.1), which has been shown to follow from the equation of motion, and $A = \text{const}$ as a solely temperature-dependent equation of state complete the usual general relativistic Cauchy problem for a thermodynamical system by including surface tensions.

5.5 Notation, distributions and mean curvature

To make the equations better readable, some conventions and formulas shall be given. Let ξ^α , $\alpha = 0, 2, 3$, be coordinates in Σ and x^i , $i = 0, 1, 2, 3$,

those for V_4 . The embedding $\Sigma \subset V_4$ is performed by functions $x^i(\xi^\alpha)$ whose derivatives

$$e_\alpha^i = \partial x^i / \partial \xi^\alpha \equiv x_{,\alpha}^i \quad (5.20)$$

form a triad field in Σ . Σ divides, at least locally, V_4 into two connected components, V_+ and V_- , and the normal n_i , defined by

$$n_i n^i = 1, \quad n_i e_\alpha^i = 0 \quad (5.21)$$

is chosen into the V_+ direction, which can be thought being the outer region. Possibly V_+ and V_- are endowed with different coordinates x_+^i , x_-^i and metrics g_{ik+} and g_{ik-} , respectively. For this case all subsequent formulas had to be indexed with $+/-$, and only the inner metric of Σ , its first fundamental tensor

$$g_{\alpha\beta} = e_\alpha^i e_\beta^k g_{ik} \quad (5.22)$$

has to be the same in both cases. As usual, we require g_{ik} to be C^2 -differentiable except for jumps of $g_{ij,k}$ at Σ . Covariant derivatives within V_4 and Σ will be denoted by ∇ and $\parallel$ respectively.

The second fundamental tensor $k_{\alpha\beta}^\pm$ on both sides of Σ is defined by

$$k_{\alpha\beta}^\pm = \left(e_\alpha^i e_\beta^k n_{i;k} \right)^\pm = \left(e_{\alpha;k}^i e_\beta^k n_i \right)^\pm \quad (5.23)$$

and the difference, $\Delta k_{\alpha\beta} = k_{\alpha\beta}^+ - k_{\alpha\beta}^-$, $\Delta k = g^{\alpha\beta} \Delta k_{\alpha\beta}$, enters the energy-momentum tensor via equation (5.7) and the relation

$$\tau^{ik} \equiv e_\alpha^i e_\beta^k \tau^{\alpha\beta}, \quad \text{where} \quad \kappa \tau_{\alpha\beta} = g_{\alpha\beta} \Delta k - \Delta k_{\alpha\beta} \quad (5.24)$$

cf. e.g., [117]. From equation (5.21) and the Lanczos equation (5.24) one obtains $n^i \tau_{ik} = 0$, i.e., indeed the absence of a delta-like energy flow in the normal direction.

Now take a special coordinate system: $x^\alpha = \xi^\alpha$, and the x^1 lines are geodesics starting from Σ into n_i direction with natural parameter x^1 . Then the line element of V_4 reads

$$ds^2 = -\left(dx^1\right)^2 + g_{\alpha\beta}dx^\alpha dx^\beta \quad (5.25)$$

and the only jumps of Γ_{jk}^i are

$$\Gamma_{\alpha\beta}^{\pm 1} = -k_{\alpha\beta}^{\pm 1} = -\frac{1}{2}g_{\alpha\beta,1}^\pm.$$

The most natural definition of $\theta \cdot \delta$ is $\frac{1}{2}\delta$ being equivalent to the choice

$$\Gamma_{jk}^i = \frac{1}{2}\left(\Gamma_{jk}^{+i} + \Gamma_{jk}^{-i}\right) \quad \text{on} \quad \Sigma. \quad (5.26)$$

But cf. Dautcourt 1963 [60] for another choice of Γ_{jk}^i with the consequence that $T_{i;k}^k \neq 0$ at Σ .

Now the δ part of the equation $T_{1;k}^k = 0$ reads

$$\Delta T_1^1 \equiv T_{+1}^1 - T_{-1}^1 = \frac{1}{2}\left(k_{\alpha\beta}^+ + k_{\alpha\beta}^-\right)\tau^{\alpha\beta}. \quad (5.27)$$

Analogously one obtains for the other components

$$\Delta T_\alpha^1 = -\tau_{\alpha\parallel\beta}^\beta. \quad (5.28)$$

Reintroducing the original coordinate system, the left-hand sides of equations (5.27) and (5.28) have to be replaced by $\Delta P_1 = \Delta n_i n^k T_k^i$ and $\Delta P_\alpha = \Delta n_i e_\alpha^k T_k^i$, respectively; cf. [151]. Thereby, ΔP_i is the difference of the energy flows on both sides of Σ .

To deduce equation eq. (5.8) we take proper time ξ^0 and angular coordinates $\psi = \xi^2$ and $\varphi = \xi^3$. Then, inside Σ ,

$$ds^2 = -\left(d\xi^0\right)^2 + R^2 d\Omega^2. \quad (5.29)$$

Using equation (5.23) and the exterior Schwarzschild solution one obtains $k_{\alpha\beta}^+$ the non-vanishing components of which are

$$k_{00}^+ = -M/R^2(l - 2M/R)^{1/2}, \quad k_{22}^+ = R(1 - 2M/R)^{1/2} \quad (5.30)$$

and

$$k_{33}^+ = k_{22}^+ \sin^2 \psi \quad (5.31)$$

because of spherical symmetry. To avoid long calculations with the metric (5.4) one can proceed as follows. By construction, $\tau_{00} = 0$, and together with equation (5.24) and the spherical symmetry $k_{22}^- = k_{22}^+$, $k_{33}^- = k_{33}^+$ follows. From equation (5.27), equation (5.8) follows then immediately without the necessity of determining the actual value of k_{00}^- .

And to be independent of the discussions connected with equation (5.26), we deduce equation (5.27) another way. First, independent of surface layers, for an arbitrary timelike hypersurface and a coordinate system such that eq. (5.25) holds, we have

$$\kappa T_{11} = \frac{1}{2} \left({}^{(3)}R + k^2 - k_{\alpha\beta} k^{\alpha\beta} \right) \quad (5.32)$$

where ${}^{(3)}R$ is the curvature scalar within that surface. Now turn to a surface layer with $k_{\alpha\beta}^+ \neq k_{\alpha\beta}^-$. Then equation (5.32) splits into a “+” and a “−” equation, having in common solely ${}^{(3)}R$. Inserting all this into equation (5.24), one obtains

$$\begin{aligned} \frac{1}{2} (k_{\alpha\beta}^+ + k_{\alpha\beta}^-) \tau^{\alpha\beta} &= \frac{1}{2\kappa} (k_{\alpha\beta}^+ + k_{\alpha\beta}^-) (g^{\alpha\beta} \Delta k - \Delta k^{\alpha\beta}) \\ &= \frac{1}{2\kappa} \left[(k^+)^2 - (k^-)^2 - k_{\alpha\beta}^+ k^{+\alpha\beta} + k_{\alpha\beta}^- k^{-\alpha\beta} \right] = T_{11}^+ - T_{11}^- \end{aligned} \quad (5.33)$$

i.e., just equation (5.27).

6 The massive scalar field in a closed Friedmann universe

For the minimally coupled scalar field in Einstein's theory of gravitation we look for the space of solutions within the class of closed Friedmann universe models. We prove $D \geq 1$, where D is the dimension of the set of solutions which can be integrated up to $t \rightarrow \infty$. $D > 0$ was conjectured by Page in 1984 [175]. We discuss concepts like “the probability of the appearance of a sufficiently long inflationary phase” and argue that it is primarily a probability measure μ in the space V of solutions and not in the space of initial conditions, which has to be applied. The measure μ is naturally defined for Bianchi-type I cosmological models because V is a compact cube. The problems with the closed Friedmann model, which led to controversial claims in the literature, will be shown to originate from the fact that V has a complicated non-compact non-Hausdorff Geroch topology: no natural definition of μ can be given. We conclude: the present state of our universe can be explained by models of the type discussed, but thereby the anthropic principle cannot be fully circumvented. We consider a closed Friedmann cosmological model,

$$ds^2 = g_{ij}dx^i dx^j = dt^2 - a^2(t)[dr^2 + \sin^2 r(d\psi^2 + \sin^2 \psi d\chi^2)], \quad (6.1)$$

with the cosmic scale factor $a(t)$. We apply Einstein's General Relativity Theory and take a minimally coupled scalar field ϕ without self-interaction as source, i.e., the Lagrangian is

$$\mathcal{L} = \frac{R}{16\pi G} + \frac{1}{2}g^{ij}\nabla_i\phi\nabla_j\phi - \frac{1}{2}m^2\phi^2. \quad (6.2)$$

In this chapter we assume units such that $\hbar = c = 1$; R is the scalar curvature, G Newton's constant, and m the mass of the scalar field. The variational

derivative $\delta\mathcal{L}/\delta\phi = 0$ yields the field equation for the scalar field

$$(m^2 + \square)\phi = 0, \quad \square \equiv g^{ij}\nabla_i\nabla_j \quad (6.3)$$

where $\square$ denotes the covariant D'Alembertian. The variation of the Lagrangian with respect to the metric

$$\delta\mathcal{L}\sqrt{-\det g_{kl}}/\delta g^{ij} = 0 \quad (6.4)$$

yields Einstein's field equation

$$R_{ij} - \frac{R}{2}g_{ij} = 8\pi GT_{ij}, \quad T_{ij} \equiv \nabla_i\phi\nabla_j\phi - \frac{1}{2}\left(\nabla^k\phi\nabla_k\phi - m^2\phi^2\right)g_{ij}, \quad (6.5)$$

with Ricci tensor R_{ij} . It is the aim of this chapter, which is based on [215], to present some rigorous results about the space V of solutions of eqs. (6.3), (6.5) with metric (6.1). This is done in sections 6.1, 6.2 and 6.3. We discuss them in the context of cosmology in section 6.4: the probability of the appearance of a sufficiently long inflationary phase and the anthropic principle.

Schrödinger [234] already dealt with the massive scalar field in the closed Friedmann universe in 1938, but there the back-reaction of the scalar field on the evolution of the cosmic scale factor had been neglected. 35 years later the model enjoyed a renewed interest, especially as a semi-classical description of quantum effects, cf. e.g. Fulling and Parker [82], [83], Starobinsky [243], Barrow and Matzner [14], Gottlöber [92] and Hawking and Luttrell [108]. One intriguing property - the possibility of a bounce, i.e., a positive local minimum of the cosmic scale factor - made it interesting in connection with a possible avoidance of a big bang singularity $a(t) \rightarrow 0$. The existence of periodic solutions $(\phi(t), a(t))$ of eqs. (6.1), (6.3) and (6.5) became clear in 1984 by independent work of Hawking [106] and Gottlöber and Schmidt [96].

The existence of an inflationary phase, which is defined by

$$|dh/dt| \ll h^2, \quad \text{where} \quad h = d(\ln a)/dt \quad (6.6)$$

is the Hubble parameter, in the cosmic evolution is discussed in many papers to that model, cf e.g. Belinsky et al. [20], [21], Gottlöber and Müller [93] and Page 1987 [175]. Soon it became clear that the satisfactory results obtained for the spatially flat model – the existence of a naturally defined measure in the space of solutions and with this measure the very large probability to have sufficient inflation – cannot be generalized to the closed model easily. We shall turn to that point in section 6.4.

6.1 Some closed-form approximations

We consider the system (6.3), (6.5) with metric (6.1). Sometimes, one takes it as an additional assumption that ϕ depends on the coordinate t only, but it holds, cf. Turner [258]: the spatial homogeneity of metric (6.1) already implies this property. Proof: For $i \neq j$ we have $R_{ij} = g_{ij} = 0$ and, therefore we get $\nabla_i \phi \nabla_j \phi = 0$. This means, locally ϕ depends on one coordinate x^i only. Supposed $i \neq 0$, then $g^{ij} T_{ij} - 2T_{00} = m^2 \phi^2$. The l.h.s. depends on t only. For $m \neq 0$ this is already a contradiction. For $m = 0$ we have $R_{ij} = \nabla_i \phi \nabla_j \phi$, which is a contradiction to the spatial isotropy of R_{ij} , q.e.d.

We always require $a(t) > 0$, for otherwise the metric (6.1) is degenerated, leading to a “big bang”. Inserting metric (6.1), eqs. (6.3), (6.5) reduce to

$$d^2 \phi / dt^2 + 3h d\phi / dt + m^2 \phi = 0 \quad (6.7)$$

and

$$3(h^2 + a^{-2}) = 4\pi G [m^2 \phi^2 + (d\phi/dt)^2]. \quad (6.8)$$

Eq. (6.8) is the 00-component of eq. (6.5), the other components are a consequence of it. To get the analogous equations for the spatially flat Friedmann model, the l.h.s. of eq. (6.8) has to be replaced by $3h^2$.

For the massless case $m = 0$ representing stiff matter, i.e., pressure equals energy density, eqs. (6.7), (6.8) can be integrated in closed form. First, for the spatially flat model, we get solution eq. (3.4) with $\tau = 1/3$, i.e. $a(t) = t^{1/3}$; this means $\alpha = 1$ in eq. (3.5). Second, for the closed model, let $\psi = d\phi/dt$, then eq. (6.7) implies $\psi a^3 = \text{const}$. Inserting this into eq. (6.8), we get

$$da/dt = \pm \sqrt{L^4/a^4 - 1} \quad (6.9)$$

with a constant $L > 0$ and $0 < a \leq L$. We get $a(t)$ via the inverted function up to a t translation from the following equation

$$\pm t = \frac{1}{2}L \arcsin(a/L) - \frac{a}{2}\sqrt{1 - a^2/L^2}. \quad (6.10)$$

The minimally coupled massless scalar field in Einstein's theory is conformally equivalent to the theory of gravity following from the Lagrangian $L = R^2$, cf. Bicknell [29], or [245] and [16]. Details of this Bicknell theorem are presented in chapter 8, especially section 8.5.

For $a \ll L$ we have from eq. (6.10) $a \sim t^{1/3}$ and $\phi = \phi_0 + \phi_1 \ln t$. The limit $L \rightarrow \infty$ is, only locally of course, the limit from the closed to the spatially flat Friedmann model. Thus we recover, as it must be the case, $\tau = 1/3$ given above. Eq. (6.10) shows that the function $a(t)$ has a maximum at $a = L$. $a(t) > 0$ is fulfilled for a t -interval of length $\Delta t = \pi L/2$ only, and there is no bounce. In which range can one except the massless case to be a good approximation for the massive case? To this end we perform the following

substitutions:

$$\tilde{t} = t/\varepsilon, \quad \tilde{a} = a/\varepsilon, \quad \tilde{m} = m\varepsilon, \quad \tilde{\phi} = \phi. \quad (6.11)$$

They do not change the differential equations. Therefore, to get a solution $a(t)$ with a maximum $a_{\max} = \varepsilon \ll 1$ for m fixed, we can transform to a solution with $\tilde{a}_{\max} = 1$, $\tilde{m} = \varepsilon m \ll m$ and apply solutions $\tilde{a}(\tilde{t})$ with $\tilde{m} = 0$ as a good approximation. Then $a(t) > 0$ is fulfilled for a t -interval of length $\Delta t \approx \pi a_{\max}/2$ only. This is in quite good agreement with the estimate in eq. (22) of Page 1984 [175]. But in the contrary to the massless case it holds: to each $\varepsilon > 0$ there exist bouncing solutions which possess a local maximum $a_{\max} = \varepsilon$.

Let henceforth be $m > 0$. Then we use $\sqrt{4\pi G/3} \cdot \phi$ instead of ϕ as scalar field and take units such $m = 1$. With a dot denoting d/dt we finally get from eqs. (6.7), (6.8)

$$\ddot{\phi} + 3h\dot{\phi} + \phi = 0, \quad h^2 + a^{-2} = \phi^2 + \dot{\phi}^2. \quad (6.12)$$

$\phi \rightarrow -\phi$ is a Z_2 -gauge transformation, where Z_2 denotes the two-point group. Derivating eq. (6.12) we can express ϕ and $\dot{\phi}$ as follows

$$\phi = \pm \sqrt{2 + 2\dot{a}^2 + a\ddot{a}} / \sqrt{3a^2}, \quad \dot{\phi} = \pm \sqrt{1 + \dot{a}^2 - a\ddot{a}} / \sqrt{3a^2}. \quad (6.13)$$

Inserting eq. (6.13) into eq. (6.12) we get

$$a^2 \frac{d^3 a}{dt^3} = 4\dot{a}(1 + \dot{a}^2) - 3a\ddot{a} \pm 2a\sqrt{2 + 2\dot{a}^2 + a\ddot{a}} \sqrt{1 + \dot{a}^2 - a\ddot{a}}. \quad (6.14)$$

On the r.h.s. we have “+” if $\dot{\phi} > 0$ and “−” otherwise. At all points t , where one of the roots becomes zero, “+” and “−” have to be interchanged.

To get the temporal behaviour for very large values $a \gg 1$ but small values $|h|$, we make the ansatz

$$a(t) = 1/\varepsilon + \varepsilon A(t), \quad \varepsilon > 0, \quad \varepsilon \approx 0. \quad (6.15)$$

In lowest order of ε we get from eq. (6.14)

$$\frac{d^3 A}{dt^3} = \pm 2 \sqrt{2 + \ddot{A}} \sqrt{1 - \ddot{A}}. \quad (6.16)$$

An additive constant to $A(t)$ can be absorbed by a redefinition of ε , eq. (6.15), so we require $A(0) = 0$. After a suitable translation of t , each solution of eq. (6.16) can be represented as

$$A(t) = \alpha t - t^2/4 + \frac{3}{8} \sin(2t), \quad |\alpha| \leq \pi/4. \quad (6.17)$$

$\alpha = \pi/4$ and $\alpha = -\pi/4$ represent the same solution, so we have a S^1 space of solutions. Here, S^n denotes the n -dimensional sphere. $a(t) > 0$ is fulfilled for $A(t) > -1/\varepsilon^2$, i.e. $|t| < 2/\varepsilon$ only. In dependence of the value α , $A(t)$ has one or two maxima and, accordingly, null or one minimum. The corresponding intervals for α meet at two points, $\alpha \approx 0$ and $|\alpha| \approx \pi/4$, where one maximum and one horizontal turning point

$$\dot{a} = \ddot{a} = 0, \quad \frac{d^3 a}{dt^3} \neq 0 \quad (6.18)$$

exist.

6.2 The qualitative behaviour

Eqs. (6.12) represent a regular system: at $t = 0$ we prescribe a_0 , $\text{sgn } \dot{a}_0$, ϕ_0 , $\dot{\phi}_0$ fulfilling

$$a_0^2 (\phi_0^2 + \dot{\phi}_0^2) \leq 1 \quad (6.19)$$

and then all higher derivatives can be obtained by differentiating:

$$|\dot{a}| = \sqrt{a^2(\phi^2 + \dot{\phi}^2) - 1}, \ddot{\phi} = -3\dot{a}\dot{\phi}/a - \phi, \quad (6.20)$$

the r.h.s. being smooth functions. It follows

$$d/dt (h^2 + a^{-2}) = -6h(d\phi/dt)^2. \quad (6.21)$$

6.2.1 Existence of a maximum

The existence of a local maximum for each solution $a(t)$ already follows from the “closed universe recollapse conjecture”, but we shall prove it for our model as follows: if we start integrating with $h \geq 0$; otherwise, $t \rightarrow -t$ serves to reach that; then $h^2 + a^{-2}$ is a monotonously decreasing function as long as $h \geq 0$ holds, because $h\dot{\phi} = 0$ holds at isolated points t only, cf. eq. (6.21). We want to show that after a finite time, h changes its sign giving rise to a local maximum of $a(t)$. If this is not the case after a short time, then we have after a long time $h \ll 1$, $a \gg 1$, and eqs. (6.15), (6.17) become a good approximation to the exact solution. The approximate solution (6.15), (6.17) has already shown to possess a local quadratic maximum and this property is a stable one within C^2 -perturbations. So the exact solution has a maximum, too, q.e.d.

6.2.2 The space of solutions

We denote the space of solutions for eqs. (6.1), (6.12) by V and endow it with Geroch’s topology from 1969, see [86] and subsection 6.3.3 for further details. Applying the result of subsection 6.2.1, V can be constructed as follows: the set of solutions will not be diminished if we start integrating

with $\dot{a}_0 = 0$. We prescribe

$$f = a_0^{-1} \quad \text{and} \quad g = (1 - a_0 \ddot{a}_0) \cdot \text{sgn}(\phi_0 \dot{\phi}_0), \quad (6.22)$$

then all other values are fixed. f and g are restricted by $f > 0$; $|g| \leq 3$, where $g = 3$ and $g = -3$ describe the same solution. Therefore

$$V = (\mathcal{R} \times S^1)/Q, \quad (6.23)$$

where $\mathcal{R}$ denotes the real line, considered as topological space, and Q is an equivalence relation defined as follows: Some solutions $a(t)$ have more than one, but at most countably many, extremal points, but each extremum defines one point in $\mathcal{R} \times S^1$; these are just the points being Q -equivalent.

Considering eqs. (6.12), (6.22) in more details, we get the following: for $|g| < 1$ we have a minimum and for $|g| > 1$ a maximum for $a(t)$.

$$\frac{d^3 a_0}{dt^3} > 0 \quad (6.24)$$

holds for $0 < g < 3$. $a(t)$ is an even function for $g = 0$ and $g = \pm 3$ only. For $g = 0$ it is a symmetric minimum of $a(t)$ and ϕ is even, too. For $g = \pm 3$ it is a symmetric maximum of $a(t)$ and ϕ is odd. For $|g| = 1$ one has a horizontal turning point for $a(t)$. Time reversal $t \rightarrow -t$ leads to $g \rightarrow -g$.

6.2.3 From one extremum to the next

To prove our result, $D \geq 1$, where D is the dimension of the set of solutions which can be integrated up to $t \rightarrow \infty$, we need a better knowledge of the equivalence relation Q . To this end we define a map

$$p : \mathcal{R} \times S^1 \rightarrow \mathcal{R} \times S^1 \quad (6.25)$$

as follows: For $x = (f, g)$ we start integrating at $t = 0$. Let

$$t_1 = \min\{t | t > 0, \dot{a}(t) = 0\}. \quad (6.26)$$

For $t_1 < \infty$ we define

$$p(x) = \bar{x} = (\bar{f}, \bar{g}) = (f(t_1), g(t_1)) \quad (6.27)$$

and otherwise p is not defined. In words: p maps the initial conditions from one extremal point of $a(t)$ to the next one. It holds: p is injective and xQy if and only if there exists an integer m such that $p^m(x) = y$. Let $V_m \subset \mathcal{R} \times S^1$ be that subspace for which p^m is defined. Then the inclusion

$$\{(f, g) | -1 < g \leq 1\} \subset V_1 \quad (6.28)$$

means: a minimum is always followed by a maximum; $|g| < 1$ implies $|\bar{g}| \geq 1$. $V_2 \neq \emptyset$ follows from the end of section 6.1. For $x = (f, g)$ we define $-x = (f, -g)$. With this notation it holds, see the end of subsection 6.2.2,

$$p^m(V_m) = V_{-m} = -V_m, \quad (6.29)$$

and for $x \in V$ we have

$$p^m(-p^m(x)) = -x \quad \text{and} \quad p^{-m}(V_m \cap V_{-m}) = V_{2m}. \quad (6.30)$$

A horizontal turning point can be continuously deformed to a pair of extrema; such points give rise to discontinuities of the function p , but for a suitably defined non-constant integer power m the function p^m is a continuous one. If $x \in V_1 \setminus \text{int}(V_1)$ where int denotes the topological interior, then $a(t)$ possesses a horizontal turning point. To elucidate the contents of these sentences we give an example:

For very small values f and $\bar{f}$ we may apply eqs. (6.15), (6.17) to calculate the function p^m . In the approximation used, no more than 3 extremal points appear, so we have to consider

$$p^m, \quad m \in \{-2, -1, 0, 1, 2\} \quad (6.31)$$

only. f and $\bar{f}$ approximately coincide, so we concentrate on the function $\bar{g}(g)$. The necessary power m is sketched at the curve. To come from m to $-m$, the curve has to be reflected at the line $\bar{g} = g$. Time reversal can be achieved, if it is reflected at $\bar{g} = -g$. V_1 is the interval $-1 < g \leq 2.8853$. The jump discontinuity of p at $g = 1$ and the boundary value $g = 2.8853 \dots$ are both connected with horizontal turning points $g, \bar{g} = \pm 1$. The shape of the function $\bar{g}(g)$ for $m = 1$ can be obtained by calculating the extrema of $A(t)$ eq. (6.17) in dependence of α and then applying eq. (6.22). For $m \neq 1$ one applies eq. (6.30).

6.2.4 The periodic solutions

The periodic solutions $a(t)$ are characterized by the fixed points of some p^m , $m \geq 2$, whereas p itself has no fixed points. The notation p^m means, that the map p , eqs. (6.25), (6.27), will be applied m times. The existence of the fixed points can be proved as follows: we start integrating at $t = 0$ with a symmetric minimum, $g = 0$, of $a(t)$ and count the number m of zeros of $\phi(t)$ in the time interval $0 < t < t_2$, where $a(t_2)$ is the first local maximum of $a(t)$. The number m depends on $f = 1/a_0$ and has jump discontinuities only at points where $\phi(t_2) = 0$. For initial values $f = f_k$, where the number m jumps from k to $k + 1$, the function $a(t)$ is symmetric about $t = t_2$. But a function, which is symmetric about two different points, is a periodic one. We call periodic solutions obtained by this procedure, periodic solutions of

the first type, they represent fixed points of the map p^2 . Page 1984 [175] gives the numerically obtained result:

$$f_1 = 1/a_0 \quad \text{with} \quad a_0 = 0.76207 \dots \quad (6.32)$$

That f_k exists also for very large values k can be seen as follows: take solution eqs. (6.15), (6.17) with a very small value ε and insert it into eq. (6.13); we roughly get $m = 1/\varepsilon\pi$.

Fixed points of p^4 , which are not fixed points of p^2 , are called periodic solutions of the second type. They can be constructed as follows: we start with $g = 0$ but $f \neq f_k$ and count the number n of zeros of $\dot{\phi}(t)$ in the interval $0 < t < t_3$, where $a(t_3) > 0$ is next local minimum of $a(t)$. For $f \approx f_k$, t_3 is defined by continuity reasons. For $f = f^l$, the number n jumps from l to $l + 1$, and we have a periodic solution. Numerical evaluations yield

$$f^1 = 1/a_0, \quad a_0 = 0.74720 \dots \quad (6.33)$$

The existence of f^l for very large values l is again ensured by solution (6.15), (6.17).

6.2.5 The aperiodic perpetually oscillating solutions

Now we look at the solutions in the neighbourhood of the periodic ones. Let $x_0 = (f_k, 0) \in V_2$ be one of the fixed points of p^2 representing a periodic solution of the first type; $a_0(t)$, the corresponding solution has no horizontal turning points. Therefore, the function p is smooth at x_0 . Let us denote the circle with boundary of radius ε around x_0 by $K(\varepsilon)$. By continuity reasons there exists an $\varepsilon > 0$ such that $K(\varepsilon) \subset \text{int}(V_2 \cap V_{-2})$ and the first two extrema of the functions $a(t)$ corresponding to points of $K(\varepsilon)$ are either maxima or

minima. Let $R_0 = K(\varepsilon)$ and for $n \geq 1$

$$R_n := R_0 \cap p^2(R_{n-1}) ; \quad R_{-n} := R_0 \cap p^{-2}(R_{1-n}) . \quad (6.34)$$

By assumption, $p^{\pm 2}$ is defined in R_0 , so R_m is a well-defined compact set with $x_0 \in \text{int } R_m$ for each integer m . It holds

$$R_{n+1} \subset R_n , \quad R_{-n} = -R_n \quad (6.35)$$

and

$$R_n = \{x | p^{2m}(x) \in R_0 \text{ for } m = 0, \dots, n\} . \quad (6.36)$$

Then the set

$$R_\infty := \bigcap_{n=0}^{\infty} R_n \quad (6.37)$$

is a non-empty compact set with $x_0 \in R_\infty$. In words: R_n represents the set of all these solutions which have n different maxima. R_∞ then corresponds to the set of solutions possessing infinitely many maxima. For each $x \in R_\infty$, the corresponding solution $a(t)$ can be integrated up to $t \rightarrow \infty$. The last statement follows from the fact that the time from one extremum to the next is bounded from below by a positive number within the compact set R_0 , i.e., an infinite number of extrema can be covered only by an infinite amount of time. Analogous statements hold for $R_{-\infty} = -R_\infty$ and $t \rightarrow -\infty$.

Let us fix an integer $m \geq 1$. We start integrating at $x_\delta := (f_k, \delta) \in R_0$, $0 < \delta \leq \varepsilon$.

$$\delta(m) := \max\{\delta | \delta \leq \varepsilon, p^{2k}(x_\delta) \in R_0 \text{ for } k = 0, \dots, m\} \quad (6.38)$$

exists because of compactness, i.e., $x_{\delta(m)} \in R_m$ and there is an integer $k(m) \leq m$ such that

$$y(m) := p^{2k(m)}(x_{\delta(m)}) \in \delta R_0 , \quad (6.39)$$

δR_0 being the boundary of R_0 .

$$j(m) := \begin{cases} 1, & \text{if } k(m) < m/2 \\ 0 & \text{otherwise.} \end{cases} \quad (6.40)$$

The sequence

$$(y(m), j(m)) \subset S^1 \times Z_2 \quad (6.41)$$

possesses a converging subsequence with

$$y(\infty) := \lim_{i \rightarrow \infty} y(m_i), \quad j(\infty) := \lim_{i \rightarrow \infty} j(m_i). \quad (6.42)$$

Now let us start integrating from $y(\infty)$ forward in time for $j(\infty) = 1$ and backwards in time for $j(\infty) = 0$. We consider only the case $j(\infty) = 1$, the other case will be solved by $t \rightarrow -t$. For each I with $m_i > 2m$ we have $y(m_i) \in R_m$, therefore, $y(\infty) \in R_m$ for all m , i.e., $y(\infty) \in R_\infty$. $y(\infty) \in \delta R_0$ and for all $n \geq 1$, $p^{2n}(y(\infty)) \in R_0$.

By continuously diminishing ε we get a one-parameter set of solutions $a_\varepsilon(t)$, which can be integrated up to $t \rightarrow \infty$. To this end remember that one solution $a(t)$ is represented by at most countably many points of R_0 . Supposed $a_\varepsilon(t)$ is a periodic function. By construction this solution has a symmetric minimum and there exist only countably many such solutions. Let the set M of solutions be defined as follows: $M = \{a(t) | a(0) \text{ is a minimum parametrized by a point of } R_0, a(t_1) \text{ is the next minimum, and } 0 \leq t \leq t_1\}$; then we get $M = [a_{\min}, a_{\max}]$ with $a_{\min} > 0$, $a_{\max} < \infty$, and for each ε and each $t \geq 0$ it holds $a_{\min} \leq a_\varepsilon(t) \leq a_{\max}$. So we have proven: in each neighbourhood of the periodic solutions of the first type there exists a set of Hausdorff dimension $D \geq 1$ of uniformly bounded aperiodic perpetually oscillating solutions which can be integrated up to $t \rightarrow \infty$.

6.3 Problems with die probability measure

To give concepts like “the probability p of the appearance of a sufficiently long inflationary phase” a concrete meaning, we have to define a probability measure μ in the space V of solutions. Let us suppose we can find a hypersurface H in the space G of initial conditions such that each solution is characterized by exactly one point of H . Then V and H are homeomorphic and we need not to make a distinction between them. Let us further suppose that $H \subset G$ is defined by a suitably chosen physical quantity ψ to take the Planckian value. Then we are justified to call H the quantum boundary. By construction, H divides G into two connected components; $\psi \leq \psi_{\text{Pl}}$ defines the classical region. All classical trajectories start their evolution at H and remain in the classical region forever.

Let us remember the situation for the spatially flat Friedmann model Belinsky et al. [20] subsection 6.3.1, and for the Bianchi type I model, Lukash and Schmidt [145] subsection 6.3.2, before we discuss the closed Friedmann model in subsection 6.3.3.

6.3.1 The spatially flat Friedmann model

For this case, eq. (6.21) reduces to $h\dot{h} = -3h\dot{\phi}^2$, i.e., each solution crosses the surface $h = h_{\text{Pl}}$ exactly once, the only exception is the flat Minkowski spacetime $h \equiv 0$. Reason: For $\dot{\phi} \neq 0$ we get $\dot{h} < 0$. The corresponding physical quantity

$$\psi = h^2 = \phi^2 + \dot{\phi}^2 \tag{6.43}$$

is the energy density. The space of non-flat spatially flat Friedmann models is topologically S^1 , and equipartition of initial conditions gives a natural

probability measure there. With this definition it turned out that for $m \ll m_{\text{Pl}}$ it holds

$$p \approx 1 - 8m/m_{\text{Pl}}, \quad (6.44)$$

cf. e.g. Müller and Schmidt [163], i.e., inflation becomes quite probable. If, on the other hand, equipartition is taken at some $h_0 \ll h_{\text{Pl}}m/m_{\text{Pl}}$ then inflation is quite improbable. The total space V of solutions has Geroch topology αS^1 , i.e., $V = S^1 \cup \{\alpha\}$, and the space itself is the only neighbourhood around the added point α which corresponds to $h \equiv 0$, because each solution is asymptotically flat for $t \rightarrow \infty$.

6.3.2 The Bianchi-type I model

With the metric

$$ds^2 = dt^2 - e^{2\alpha} [e^{2(s+\sqrt{3}r)} dx^2 + e^{2(s-\sqrt{3}r)} dy^2 + e^{-4s} dz^2], \quad h = \dot{\alpha} \quad (6.45)$$

the analogue to eq. (6.43) is

$$\psi = h^2 = \dot{\phi}^2 + \dot{\phi}^2 + \dot{r}^2 + \dot{s}^2, \quad (6.46)$$

and $h = h_{\text{Pl}}$ defines a sphere S^3 in eq. (6.46). Here all solutions cross this sphere exactly once, even the flat Minkowski spacetime: it is represented as (0 0 1)-Kasner solution α , so the space of solutions is $V = S^3/Q$, where Q is a 12-fold cover of S^3 composed of the Z_2 -gauge transformation $\phi \rightarrow -\phi$ and of the six permutations of the three spatial axes. Eq. (6.46) induces a natural probability measure on the space $\psi = \psi_{\text{Pl}}$ by equipartition, and the equivalence relation Q does not essentially influence this. As in subsection 6.3.1, $\alpha \in V$ has only one neighbourhood: V itself. Up to this exception, V is topologically a 3-dimensional cube with boundary. One diagonal line

through it represents the Kasner solution $\phi \equiv 0$. The boundary δV of V has topology S^2 and represents the axially symmetric solutions and one great circle of it the isotropic ones. As usual, the solutions with higher symmetry form the boundary of the space of solutions. p turns out to be the same as in subsection 6.3.1 and how ψ eq. (6.46) can be invariantly defined, is discussed in [145].

6.3.3 The closed Friedmann model

Now we come to the analogous questions concerning the closed Friedmann model. Before defining a measure, one should have a topology in a set. I feel it should be a variant of Geroch [86]. The Geroch topology, cf. [210], is defined as follows: let $x_i = (a_i(t), \phi_i(t))$ be a sequence of solutions and $x = (a(t), \phi(t))$ a further solution. Then $x_i \rightarrow x$ in Geroch's topology, if there exist suitable gauge and coordinate transformations after which $a_i(t) \rightarrow a(t)$ and $\phi_i(t) \rightarrow \phi(t)$ converge uniformly together with all their derivatives in the interval $t \in [-\varepsilon, \varepsilon]$ for some $\varepsilon > 0$. Because of the validity of the field equations, "with all their derivatives" may be substituted by "with their first derivatives".

With this definition one gets just the same space V as in subsection 6.2.2, eq. (6.23). The existence of aperiodic perpetually oscillating solutions, which go right across the region of astrophysical interest subsection 6.2.5 shows that for a subset of dimension $D \geq 1$ of $R \times S^1$, Q identifies countably many points. All these points lie in a compact neighbourhood of the corresponding periodic point $(f_k, 0)$ and possess therefore at least one accumulation point z . At these points z , V has a highly non-Euclidean topology. Further, V has a non-compact non-Hausdorff topology. So there is no chance to define a

probability measure in a natural way and no possibility to define a continuous hypersurface in the space of initial conditions which each trajectory crosses exactly once.

6.4 Discussion of the inflationary phase

Supposed, we had obtained a result of the type: “Each solution $a(t)$ has at least one but at most seven local maxima.” Then one could define — up to a factor $7 = O(1)$ — a probability measure. So it is just the existence of the perpetually bouncing aperiodic solutions which gives the problems. We conclude: it is not a lack of mathematical knowledge but an inherent property of the closed Friedmann model which hinders to generalize the convincing results obtained for the spatially flat model. So it is no wonder that different trials led to controversial results, cf. Belinsky et al. [20], [21] and Page 1987 [175]. One of these results reads “inflationary and non-inflationary solutions have both infinite measure”, hence, nothing is clear.

Let us now discuss the results of Starobinsky [243] and Barrow and Matzner [14] concerning the probability of a bounce. They have obtained a very low probability to get a bounce, but they used equipartition at some $h_0 \ll h_{\text{Pl}} m/m_{\text{Pl}}$. As seen in subsection 6.3.1 for the spatially flat Friedmann model concerning inflation, this low probability does not hinder to get a considerable large probability if equipartition is applied at $h = h_{\text{Pl}}$. We conclude, the probability of bouncing solutions is not a well-defined concept up to now. Well-defined is, on the other hand, some type of conditional probability. If we suppose that some fixed value a , say 10^{28} cm or so, and there some fixed value h , say 50 km/sec · Mpc or so, appear within the cosmic evolution, then the remaining degree of freedom is just the phase of

the scalar field, which is the compact set S^1 as configuration space. But this solves not all problems, because the perpetually bouncing solutions discussed in subsection 6.2.5 cross this range of astrophysical interest infinitely often. Calculating conditional probabilities instead of absolute probabilities, and, if this condition is related to our own human existence, then we have already applied the anthropic principle. Cf. similar opinions in Singh and Padmanabhan [238] concerning the so far proposed explanations of the smallness of the cosmological constant.

The massive scalar field in a closed Friedmann model with Einstein's theory of gravity cannot explain the long inflationary stage of cosmic evolution as an absolutely probable event and so some type of an anthropic principle has to be applied; see the well-balanced monograph by Barrow and Tipler [17] to this theme. We have discussed the solutions for the minimally coupled scalar field, but many results for the conformally coupled one are similar, see e.g. Turner and Widrow [259]; this fact in turn can be explained by the existence of a conformal transformation relating between them, see [213]. The problems in the case of defining a probability measure in the set of not necessarily spatially flat Friedmann models are also discussed in Madsen and Ellis [149]. They conclude that inflation need not to solve the flatness problem. The Gibbons-Hawking-Stewart approach [89] gives approximately the same probability measure as the equipartition of initial conditions used here. The Wheeler-de Witt equation for the massive scalar field in a closed Friedmann universe model is also discussed by Calzetta [46]. Possibly, this approach is the route out of the problems mentioned here.

7 The metric in the superspace of Riemannian metrics

The space of all Riemannian metrics is infinite-dimensional. Nevertheless a great deal of usual Riemannian geometry can be carried over. The superspace of all Riemannian metrics shall be endowed with a class of Riemannian metrics; their curvature and invariance properties are discussed. Just one of this class has the property to bring the Lagrangian of General Relativity into the form of a classical particle's motion. The signature of the superspace metric depends on the signature of the original metric in a non-trivial manner, we derive the corresponding formula. Our approach, which is based on [217], is a local one: the essence is a metric in the space of all symmetric rank-two tensors, and then the space becomes a warped product of the real line with an Einstein space.

7.1 The superspace

Let $n \geq 2$, n be the dimension of the basic Riemannian spaces. Let M be an n -dimensional differentiable manifold with an atlas x of coordinates x^i , $i = 1, \dots, n$. The signature s shall be fixed; s is the number of negative eigenvalues of the metric. Let V be the space of all Riemannian metrics $g_{ij}(x)$ in M with signature s , related to the coordinates x^i .

This implies that isometric metrics in M are different points in V in general. The V is called superspace, its points are the Riemannian metrics. The tangent space in V is the vector space

$$T = \{h_{ij}(x) | x \in M, h_{ij} = h_{ji}\}, \quad (7.1)$$

the space of all symmetric tensor fields of rank 2. All considerations are local ones, so we may have in mind one single fixed coordinate system in M .

7.2 Coordinates in superspace

Coordinates should possess one contravariant index, so we need a transformation of the type

$$y^A = \mu^{Aij} g_{ij}(x) \quad (7.2)$$

such that the y^A are the coordinates for V . To have a defined one-to-one correspondence between the index pairs (i, j) and the index A we require

$$A = 1, \dots, N = n(n+1)/2, \quad (7.3)$$

and $A = 1, \dots, N$ corresponds to the pairs

$$\begin{aligned} (1, 1), (2, 2), \dots, (n, n), (1, 2), (2, 3), \dots, (n-1, n), (1, 3), \\ \dots, (n-2, n), \dots, (1, n) \end{aligned} \quad (7.4)$$

consecutively. (i, j) and (j, i) correspond to the same A . We make the ansatz

$$\mu^{Aij} = \mu_{Aij} = \begin{cases} b & \text{for } i \neq j \\ c & \text{for } i = j \\ 0 & \text{if } (i, j) \text{ does not correspond to } A \end{cases} \quad (7.5)$$

with certain real numbers b and c to be fixed later and require the usual inversion relations

$$\mu^{Aij} \mu_{Bij} = \delta_B^A \quad \text{and} \quad \mu^{Aij} \mu_{Akl} = \delta_k^{(i} \delta_l^{j)}. \quad (7.6)$$

Bracketed indices are to be symmetrized, which is necessary because of symmetry of the metric g_{ij} . Inserting ansatz (7.5) into eq. (7.6) gives $c^2 = 1$,

$b^2 = 1/2$. Changing the sign of b or c only changes the sign of the coordinates, so we may put

$$c = 1, \quad b = 1/\sqrt{2}. \quad (7.7)$$

The object μ^{Aij} is analogous to the Pauli spin matrices relating two spinorial indices to one vector index.

7.3 Metric in superspace

The metric in the superspace shall be denoted by H_{AB} , it holds

$$H_{AB} = H_{BA} \quad (7.8)$$

and the transformed metric is

$$G^{ijkl} = H_{AB} \mu^{Aij} \mu^{Bkl}, \quad H_{AB} = \mu_{Aij} \mu_{Bkl} G^{ijkl}. \quad (7.9)$$

From eqs. (7.5) and (7.8) it follows that

$$G^{ijkl} = G^{jikl} = G^{klij}. \quad (7.10)$$

The inverse to H_{AB} is H^{AB} , and we define

$$G_{ijkl} = H^{AB} \mu_{Aij} \mu_{Bkl} \quad (7.11)$$

which has the same symmetries as eq. (7.10). We require G^{ijkl} to be a tensor and use only the metric $g_{ij}(x)$ to define it.

Then the ansatz

$$G^{ijkl} = z g^{i(k} g^{l)j} + \alpha g^{ij} g^{kl} \quad (7.12)$$

$$G_{ijkl} = v g_{i(k} g_{l)j} + \beta g_{ij} g_{kl} \quad (7.13)$$

where v , z , α and β are constants, is the most general one to fulfil the symmetries eq. (7.10). One should mention that also curvature-dependent constants could have been introduced.

The requirement that H_{AB} is the inverse to H^{AB} leads via eqs. (7.9) and (7.11) to

$$G_{ijkl} G^{klmp} = \delta_i^{(m} \delta_j^{p)}. \quad (7.14)$$

The requirement that G^{ijkl} is a tensor can be justified as follows: Let a curve $y^A(t)$, $0 < t < 1$ in V be given, then its length is

$$\sigma = \int_0^1 \left(H_{AB} \frac{dy^A}{dt} \frac{dy^B}{dt} \right)^{1/2} dt \quad (7.15)$$

i.e., with eqs. (7.2) and (7.9)

$$\sigma = \int_0^1 \left(G^{ijkl} \frac{dg_{ij}}{dt} \frac{dg_{kl}}{dt} \right)^{1/2} dt. \quad (7.16)$$

A coordinate transformation in $M : x^i \rightarrow \varepsilon x^i$ changes $g_{ij} \rightarrow \varepsilon^{-2} g_{ij}$.

We now require that σ shall not be changed by such a transformation. Then α and z are constant real numbers. Inserting eqs. (7.12) and (7.13) into eq. (7.14) gives $vz = 1$, hence $z \neq 0$. By a constant rescaling we get

$$v = z = 1 \quad (7.17)$$

and then eqs. (7.12), (7.13) and (7.14) yield the conditions

$$\alpha \neq \frac{-1}{n}, \quad \beta = \frac{-\alpha}{1 + \alpha n}. \quad (7.18)$$

So we have got a one-parameter set of metrics in V .

Eq. (7.18) fulfils the following duality relation: with

$$f(\alpha) = -\alpha/(1 + \alpha n), \quad (7.19)$$

the equation $f(f(\alpha)) = \alpha$ holds for all $\alpha \neq -1/n$.

Inserting eq. (7.17) into eqs. (7.12) and (7.13), we finally get for the superspace metric

$$\begin{aligned} G^{ijkl} &= g^{i(k} g^{l)j} + \alpha g^{ij} g^{kl} \quad \text{and} \\ G_{ijkl} &= g_{i(k} g_{l)j} + \beta g_{ij} g_{kl}. \end{aligned} \quad (7.20)$$

It holds: For $\alpha = -1/n$, the metric H_{AB} is not invertible.

Indirect proof: G^{ijkl} depends continuously on α , so it must be the case with the inverse. But

$$\lim_{\alpha \rightarrow -1/n}$$

applied to G_{ijkl} gives no finite result. Contradiction.

7.4 Signature of the superspace metric

Let S be the superspace signature, i.e., the number of negative eigenvalues of the superspace metric H_{AB} . S depends on α and s , but $\alpha = -1/n$ is excluded. For convenience we define

$$\Theta = \begin{cases} 0, & \alpha > -1/n \\ 1, & \alpha < -1/n \end{cases} \quad (7.21)$$

From continuity reasons it follows that S is a function of Θ and s : $S = S(\Theta, s)$. If we transform $g_{ij} \rightarrow -g_{ij}$ i.e., $s \rightarrow n - s$, then H_{AB} is not changed, i.e.,

$$S(\Theta, s) = S(\Theta, n - s). \quad (7.22)$$

We transform g_{ij} to diagonal form as follows

$$g_{11} = g_{22} = \dots = g_{ss} = -1, \quad g_{ij} = \delta_{ij} \quad \text{otherwise}. \quad (7.23)$$

Let us now give the signature for $\Theta = 0$. To calculate $S(0, s)$ we may put $\alpha = 0$ and get with eqs. (7.9), (7.12) and (7.17)

$$H_{AB} = \mu_{Aij} \mu_{Bkl} g^{ik} g^{jl} \quad (7.24)$$

which is a diagonal matrix. It holds $H_{11} = \dots = H_{nn} = 1$ and the other diagonal components are ± 1 . A full estimate gives in agreement with eq. (7.22)

$$S(0, s) = s(n - s). \quad (7.25)$$

Now, we look for the signature for $\Theta = 1$. To calculate $S(1, s)$ we may put $\alpha = -1$ and get

$$H_{AB} = \mu_{Aij} \mu_{Bkl} (g^{ik} g^{jl} - g^{ij} g^{kl}). \quad (7.26)$$

For $A \leq n < B$, $H_{AB} = 0$, i.e., the matrix H_{AB} is composed of two blocks. For $A, B \leq n$ we get

$$H_{AB} = \begin{cases} 0 & \text{for } A = B \\ 1 & \text{for } A \neq B \end{cases} \quad (7.27)$$

a matrix which has the $(n - 1)$ -fold eigenvalue 1 and the single eigenvalue $1 - n$. For $A, B > n$ we have the same result as for the case $\alpha = 0$, i.e., we get $S(1, s) = 1 + s(n - s)$.

Result: The signature of the superspace metric is

$$S = \Theta + s(n - s). \quad (7.28)$$

7.5 Supercurvature and superdeterminant

We use exactly the same formulae as for finite-dimensional Riemannian geometry to define Christoffel affinities Γ_{BC}^A and Riemann tensor R_{BCD}^A in superspace. We even omit the prefix “super” in the following. Using eq. (7.5) we write all equations with indices $i, j = 1, \dots, n$.

Then each pair of covariant indices i, j corresponds to one contravariant index A . The following formulae appear:

$$\frac{\partial g^{ij}}{\partial g_{km}} = -g^{i(k} g^{m)j} \quad (7.29)$$

$$\Gamma^{ijklmp} = -\frac{1}{2}g^{i(k} g^{l)(m} g^{p)j} - \alpha g^{ij} g^{k(m} g^{p)l} - \frac{1}{2}g^{j(k} g^{l)(m} g^{p)i} \quad (7.30)$$

and, surprisingly independent of α we get

$$\Gamma_{ij}^{klmp} = -\delta_{(i}^{(k} g^{l)(m} \delta_{j)}^{p)} . \quad (7.31)$$

Consequently, also Riemann- and Ricci tensor do not depend on α :

$$R_{rs}^{klmpij} = \frac{1}{2} \left(\delta_{(r}^{(k} g^{l)(m} g^{p)(i} \delta_{s)}^{j)} - \delta_{(r}^{(k} g^{l)(i} g^{j)(m} \delta_{s)}^{p)} \right) . \quad (7.32)$$

Summing over $r = m$ and $s = p$ we get

$$R^{klij} = \frac{1}{4} \left(g^{ij} g^{kl} - n g^{k(i} g^{j)l} \right) . \quad (7.33)$$

The Ricci tensor has one eigenvalue 0. Proof: It is not invertible because it is proportional to the metric for the degenerated case $\alpha = -1/n$, cf. eq. (7.20) in section 7.3.

The co-contravariant Ricci tensor reads

$$R_{kl}^{ij} = G_{klmp} R^{mpij} = \frac{1}{4} \left(g^{ij} g_{kl} - n \delta_k^{(i} \delta_l^{j)} \right) , \quad (7.34)$$

and the curvature scalar is

$$R = -\frac{1}{8}n(n-1)(n+2). \quad (7.35)$$

The eigenvector to the eigenvalue 0 of the Ricci tensor is g_{ij} . All other eigenvalues equal $-n/4$, and the corresponding eigenvectors can be parametrized by the symmetric traceless matrices, i.e. the multiplicity of the eigenvalue $-n/4$ is $(n-1)(n+2)/2$. This is another view to the well-known split of metric perturbations into conformal transformations, i.e., one bulk degree of freedom on the one hand, and the remaining volume-preserving degrees of freedom on the other.

We define the superdeterminant

$$H = \det H_{AB}. \quad (7.36)$$

H is a function of g , α and n which becomes zero for $\alpha = -1/n$, cf. section 7.3. We use eqs. (7.9) and (7.20) to look in more details for the explicit value of H . Though we assume $n \geq 2$, here the formal calculation for $n = 1$ makes sense; it leads to

$$H = H_{11} = G^{1111} = g^{11}g^{11} + \alpha g^{11}g^{11} = (1 + \alpha)g^{-2}. \quad (7.37)$$

Let us return to the general case $n \geq 2$. Multiplication of g_{ij} with ε gives $g \rightarrow \varepsilon^n g$, $H_{AB} \rightarrow \varepsilon^{-2} H_{AB}$ and $H \rightarrow \varepsilon^{-n(n+1)} H$.

So we get in an intermediate step

$$H = H_1 g^{-n-1} \quad (7.38)$$

where H_1 is the value of H for $g = 1$. H_1 depends on α and n only. To calculate H_1 we put $g_{ij} = \delta_{ij}$ and get via $H_{ij} = \delta_{ij} + \alpha$, $H_{Ai} = 0$ for $A > n$,

and $H_{AB} = \delta_{AB}$ for $A, B > n$ finally

$$H_1 = 1 + \alpha n. \quad (7.39)$$

This result is in agreement with the $n = 1$ -calculation eq. (7.37) and also with the fact that $\alpha = -1/n$ gives $H = 0$.

7.6 Gravity and quantum cosmology

Now, we come to the main application: The action for gravity shall be expressed by the metric of superspace. The purpose is to explain the mathematical background of quantum cosmology, see [9], [59], [73], [101], [126], [148], [158], [179], [182] and [198] for new papers on that topic.

We start from the metric

$$ds^2 = dt^2 - g_{ij} dx^i dx^j \quad (7.40)$$

$i, j = 1, \dots, n$ with positive definite g_{ij} and $x^0 = t$. We define the second fundamental form K_{ij} by

$$K_{ij} = \frac{1}{2} g_{ij,0}. \quad (7.41)$$

The Einstein action for metric (7.40) is

$$I = - \int \frac{1}{2} {}^*R \sqrt{g} d^{n+1}x \quad (7.42)$$

where $g = \det g_{ij}$ and *R is the $(n+1)$ -dimensional curvature scalar for eq. (7.40). Indices at K_{ij} will be shifted with g_{ij} , and $K = K_i^i$. With eq. (7.41) we get

$$(K\sqrt{g})_{,0} = (K_{,0} + K^2)\sqrt{g}. \quad (7.43)$$

This divergence can be added to the integrand of eq. (7.42) without changing the field equations. It serves to cancel the term $K_{,0}$ of I . So we get

$$I = \int \frac{1}{2} (K^{ij} K_{ij} - K^2 + R) \sqrt{g} d^{n+1}x \quad (7.44)$$

where R is the n -dimensional curvature scalar for g_{ij} .

Applying eq. (7.20), we now make the ansatz for the kinetic energy

$$W = \frac{1}{2} G^{ijmp} K_{ij} K_{mp} = \frac{1}{2} (K^{ij} K_{ij} + \alpha K^2) . \quad (7.45)$$

Comparing eq. (7.45) with eq. (7.44) we see that for $\alpha = -1$

$$I = \int \left(W + \frac{R}{2} \right) \sqrt{g} d^{n+1}x \quad (7.46)$$

holds.

Surprisingly, this value for α does not depend on n . Because of $n \geq 2$ this value α gives a regular superspace metric. For $n = 1$, eq. (7.42) is a divergence, and $\alpha = -1$ does not give an invertible superspace metric. This is another form of the result that Einstein gravity does not lead to a local field equation in 1+1-dimensional spacetime.

Using the μ^{Aij} and the notations $z^A = \mu^{Aij} g_{ij} / 2$ and $v^A = dz^A / dt$ we get from eqs. (7.45) and (7.46)

$$I = \int \frac{1}{2} (H_{AB} v^A v^B + R(z^A)) \sqrt{g} d^{n+1}x \quad (7.47)$$

i.e., the action has the classical form of kinetic plus potential energy. The signature of the metric H_{AB} is $S = 1$. This can be seen from eqs. (7.21) and (7.28).

7.7 The Wheeler-DeWitt equation

In eq. (7.47), Einstein gravity is given in a form to allow canonical quantization: The momentum v^A is replaced by $-i\partial/\partial z^A$ in units where $\hbar = 1$, and then the Wheeler-DeWitt equation for the world function $\psi(z^A)$ appears as Hamiltonian constraint in form of a wave equation:

$$\left(\square - R(z^A)\right)\psi = 0. \quad (7.48)$$

After early attempts in [7], the Wheeler - DeWitt equation has often been discussed, especially for cosmology, see e.g. [30], [89], [100] and [103]. Besides curvature, matter fields can be inserted as potential, too. It is remarkable that exactly for Lorentz and for Euclidean signatures in eq. (7.40), i.e., positive and negative definite g_{ij} respectively, the usual D'Alembert operator with $S = 1$ in eq. (7.48) appears. For other signatures in eq. (7.40), eq. (7.48) has at least two timelike axes.

8 Comparison of scalar fields and $f(R)$ for cosmology

Following [209], we generalize the well-known analogies between $m^2\phi^2$ and $R+R^2$ theories to include the self-interaction $\lambda\phi^4$ -term for the scalar field. It turns out to be the $R+R^3$ Lagrangian which gives an appropriate model for it. Considering a spatially flat Friedmann cosmological model, common and different properties of these models are discussed, e.g., by linearizing around a ground state the masses of the corresponding spin 0-parts coincide. Then we prove a general conformal equivalence theorem between a Lagrangian $L = L(R)$, $L'L'' \neq 0$, and a minimally coupled scalar field in a general potential in section 8.5. This theorem was independently deduced by several persons, and it is now known as Bicknell theorem [29]. In the final section 8.6, which is based on [212] we discuss Ellis' programme, on which length scale the Einstein field equation is valid, on microscopic or on cosmic distances?

8.1 Introduction to scalar fields

For the gravitational Lagrangian

$$L = (R/2 + \beta R^2)/8\pi G, \quad (8.1)$$

where β is some free but constant parameter, the value

$$R = R_{\text{crit}} = -1/4\beta \quad (8.2)$$

is the critical value of the curvature scalar, cf. Nariai [169] and Schmidt [206], [207]. It is defined by

$$\partial L/\partial R = 0. \quad (8.3)$$

In regions where

$$R/R_{\text{crit}} < 1 \quad (8.4)$$

holds, we can define

$$\psi = \ln(1 - R/R_{\text{crit}}) \quad (8.5)$$

and

$$\tilde{g}_{ij} = (1 - R/R_{\text{crit}}) g_{ij} . \quad (8.6)$$

In units where $8\pi G = 1$ we now obtain from the Lagrangian eq. (8.1) via the conformal transformation eq. (8.6) the transformed Lagrangian

$$\tilde{L} = \tilde{R}/2 - 3\tilde{g}^{ij}\psi_{;i}\psi_{;j}/4 - (1 - e^{-\psi})^2/16\beta \quad (8.7)$$

being equivalent to L , cf. Whitt [270]; see [207] for the version of this equivalence used here.

For $\beta < 0$, i.e., the absence of tachyons in L eq. (8.1), we have massive gravitons of mass

$$m_0 = (-12\beta)^{-1/2} \quad (8.8)$$

in L , cf. Stelle [248]. For the weak field limit, the potential in eq. (8.7) can be simplified to be $\psi^2/(16 \cdot \beta)$, i.e., we have got a minimally coupled scalar field whose mass is also m_0 . The superfluous factor $3/2$ in eq. (8.7) can be absorbed by a redefinition of ψ . Therefore, it is not astonishing, that all results concerning the weak field limit for both $R + R^2$ -gravity without tachyons and Einstein gravity with a minimally coupled massive scalar field exactly coincide. Of course, one cannot expect this coincidence to hold for the non-linear region, too, but it is interesting to observe which properties hold there also.

We give only one example here: we consider a cosmological model of the spatially flat Friedmann type, start integrating at the quantum boundary, which is obtained by

$$R_{ijkl}R^{ijkl}$$

on the one hand, and T_{00} on the other hand, to have Planckian values, with uniformly distributed initial conditions and look whether or not an inflationary phase of the expansion appears. In both cases we get the following result: The probability p to have sufficient inflation is about $p = 1 - \sqrt{\lambda}m_0/m_{\text{Pl}}$, i.e., $p = 99.992\%$ if we take $m_0 = 10^{-5}m_{\text{Pl}}$ from GUT and $\lambda = 64$, where e^λ is the linear multiplication factor of inflation. Cf. Belinsky et al. [19] for the scalar field and Schmidt [207] for $R + R^2$, respectively.

From Quantum field theory, however, instead of the massive scalar field, a Higgs field with self-interaction turns out to be a better candidate for describing effects of the early universe. One of the advances of the latter is its possibility to describe a spontaneous breakdown of symmetry. In the following, we try to look for a purely geometric model for this Higgs field which is analogous to the above mentioned type where $L = R + R^2$ modelled a massive scalar field.

8.2 The Higgs field

For the massive scalar Field ϕ we have the mater Lagrangian

$$L_m = - \left(\phi_{;i}\phi^{;i} - m^2\phi^2 \right) / 2, \quad (8.9)$$

and for the Higgs field to be discussed now,

$$L_\lambda = - \left(\phi_{;i}\phi^{;i} + \mu^2\phi^2 - \lambda\phi^4/12 \right) / 2. \quad (8.10)$$

The ground states are defined by $\phi = \text{const.}$ and $\partial L/\partial\phi = 0$. This means $\phi = 0$ for the scalar field, and the three ground states $\phi = \phi_0 = 0$, and

$$\phi = \phi_{\pm} = \pm\sqrt{6\mu^2/\lambda} \quad (8.11)$$

for the Higgs field.

The expression

$$\left(\partial^2 L/\partial\phi^2\right)^{1/2} \quad (8.12)$$

represents the effective mass at these points. This gives the value m for the scalar field eq. (8.9), so justifying the notation. Further, eq. (8.12) give mass $i\mu$ at $\phi = 0$ and $\sqrt{2}\mu$ at $\phi = \phi_{\pm}$ for the Higgs field eq. (8.10). The imaginary value of the mass at the ground state $\phi = 0$ shows the instability met there, and in the particle picture, this gives rise to a tachyon.

To have a vanishing Lagrangian at the ground state $\phi_{\pm}$ eq. (8.11) we add a constant

$$\Lambda = -3\mu^4/2\lambda \quad (8.13)$$

to the Lagrangian eq. (8.10). The final Lagrangian reads

$$L = R/2 + L_{\lambda} + \Lambda \quad (8.14)$$

with L_{λ} eq. (8.10) and Λ eq. (8.13).

8.3 The non-linear gravitational Lagrangian

Preliminarily we direct the attention to the following fact: on the one hand, for Lagrangians (8.9), (8.10) and (8.14) the transformation $\phi \rightarrow -\phi$ is a pure gauge transformation, it does not change any invariant or geometric objects. On the other hand,

$$R_{ijkl} \rightarrow -R_{ijkl} \quad (8.15)$$

or simpler

$$R \rightarrow -R \quad (8.16)$$

is a gauge transformation at the linearized level only: taking

$$g_{ik} = \eta_{ik} + \varepsilon h_{ik} , \quad (8.17)$$

where

$$\eta_{ik} = \text{diag}(1, -1, -1, -1) , \quad (8.18)$$

then $\varepsilon \rightarrow -\varepsilon$ implies curvature inversion eq. (8.15). To be strict at the linearized level in ε . On the other hand, the curvature inversion eq. (8.15), and even its simpler version eq. (8.16), fails to hold quadratic in ε . This corresponds to fact that the ε^2 -term in eq. (8.7), which corresponds to the ψ^3 -term in the development of $\tilde{L}$ in powers of ψ , is the first one to break the $\psi \rightarrow -\psi$ symmetry in eq. (8.7).

In fact, the potential is essentially

$$\left(1 - e^{(-x)}\right)^2 . \quad (8.19)$$

Calculating this to *Order* = 14, the mapleresult reads

$$\begin{aligned} & x^2 - x^3 + \frac{7}{12}x^4 - \frac{1}{4}x^5 + \frac{31}{360}x^6 - \frac{1}{40}x^7 + \\ & \frac{127}{20160}x^8 - \frac{17}{12096}x^9 + \frac{73}{259200}x^{10} - \frac{31}{604800}x^{11} + \\ & \frac{2047}{239500800}x^{12} - \frac{1}{760320}x^{13} + \text{O}(x^{14}) \end{aligned} \quad (8.20)$$

Now, let us introduce the general non-linear Lagrangian $L = L(R)$ which we at the moment only assume to be an analytical function of R . The ground states are defined by $R = \text{const.}$, i.e.,

$$L'R_{ik} - g_{ik}L/2 = 0 . \quad (8.21)$$

Here, $L' = \partial L / \partial R$.

8.3.1 Calculation of the ground states

From eq. (8.21) one immediately sees that $\partial L/\partial R = 0$ defines critical values of the curvature scalar. For these values $R = R_{\text{crit}}$ it holds: For $L(R_{\text{crit}}) \neq 0$ no such ground state exists, and for $L(R_{\text{crit}}) = 0$, we have only one equation $R = R_{\text{crit}}$ to be solved with 10 arbitrary functions g_{ik} . We call these ground states degenerated ones. For $L = R^2$, $R_{\text{crit}} = 0$, this has been discussed by Buchdahl [43]. Now, let us concentrate on the case $\partial L/\partial R \neq 0$. Then R_{ij} is proportional to g_{ij} with a constant proportionality factor, i.e., each ground state is an Einstein space

$$R_{ij} = R g_{ij}/4, \quad (8.22)$$

with a prescribed constant value R . Inserting eq. (8.22) into eq. (8.21) we get as condition for ground states

$$RL' = 2L. \quad (8.23)$$

As an example, let L be a third order polynomial

$$L = \Lambda + R/2 + \beta R^2 + \lambda R^3/12. \quad (8.24)$$

We consider only Lagrangians with a positive linear term as we wish to reestablish Einstein gravity in the $\Lambda \rightarrow 0$ weak field limit, and $\beta < 0$ to exclude tachyons there.

We now solve eq. (8.23) for the Lagrangian eq. (8.24). For $\lambda = 0$ we have, independently of β , the only ground state $R = -4\Lambda$. It is a degenerated one if and only if $\beta\Lambda = 1/16$. That implies that for usual $R + R^2$ gravity eq. (8.1), i.e. $\lambda = \Lambda = 0$, we get $R = 0$ as the only ground state; it is a non-degenerated one.

Now, let $\lambda \neq 0$ and $\Lambda = 0$. To get non-trivial ground states we need the additional assumption $\lambda > 0$. Then, besides $R = 0$, the ground states are

$$R = R_{\pm} = \pm\sqrt{6/\lambda} \quad (8.25)$$

being quite analogous to the ground states eq. (8.11) of the Higgs field eq. (8.10). The ground state $R = 0$ is not degenerated. Of course, this statement is independent of λ and holds true, as one knows, for $\lambda = 0$. To exclude tachyons, we require $\beta < 0$, then R_- is not degenerated and R_+ is degenerated if and only if $\beta = -\sqrt{6/\lambda}$. The case $\lambda\Lambda \neq 0$ will not be considered here.

8.3.2 Definition of the masses

For the usual $R + R^2$ theory eq. (8.1), the mass is calculated at the level of the linearized theory. Then the equivalence to the Einstein field equation with a scalar field applies, and we use eqs. (8.12) and (8.2) to calculate

$$m_0 = (R_{\text{crit}}/3)^{1/2} = (-12\beta)^{-1/2}. \quad (8.26)$$

Thus, we recover the value eq. (8.8). But how to define the graviton's masses for the Lagrangian eq. (8.24)? To give such a definition a profound meaning one should do the following: linearize the full vacuum field equation around the ground state, preferably de Sitter- or anti-de Sitter spacetime, respectively, decompose its solutions with respect to a suitably chosen orthonormal system, which is a kind of higher spherical harmonics, and look for the properties of its single modes. For L eq. (8.1) this procedure just gave m_0 .

A little less complicated way to look at this mass problem is to consider a spatially flat Friedmann cosmological model and to calculate the frequency

with which the scale factor oscillates around the ground state, from which the mass m_0 turned out to be the graviton's mass for L eq. (8.1), too.

Keep in mind, 1. that all things concerning a linearization around flat vacuous spacetime do not depend on the parameter λ neither for the Higgs field nor for the $L(R)$ model, and 2. that a field redefinition $R \rightarrow R^* + R_\pm$ is *not* possible like $\phi \rightarrow \phi^* + \phi_\pm$ because curvature remains absolutely present.

8.4 The cosmological model

Now we take as Lagrangian eq. (8.24) and as line element

$$ds^2 = dt^2 - a^2(t)(dx^2 + dy^2 + dz^2). \quad (8.27)$$

The dot denotes d/dt and $h = \dot{a}/a$. We have

$$R = -6\dot{h} - 12h^2, \quad (8.28)$$

and the field equation will be obtained as follows.

8.4.1 The field equation

For $L = L(R)$ the variation

$$\delta \left(L \sqrt{-g} \right) / \delta g^{ij} = 0 \quad (8.29)$$

gives with $L' = \partial L / \partial R$ the following fourth-order gravitational field equation

$$L' R_{ij} - g_{ij} L / 2 + g_{ij} \square L' - L'_{;ij} = 0, \quad (8.30)$$

cf. e.g., Novotny (1985) [172]; see also the ideas presented by Kerner [125]. It holds

$$L'_{;ij} = L'' R_{;ij} + L''' R_{;i} R_{;j}. \quad (8.31)$$

With eq. (8.31), the trace of eq. (8.30) reads

$$L'R - 2L + 3L''\square R + 3L''' R_{;k}R^{;k} = 0, \quad (8.32)$$

i.e., with L eq. (8.24)

$$-2\Lambda - R/2 + \lambda R^3/12 + 6\beta\square R + \frac{3\lambda}{2}(R\square R + R_{;k}R^{;k}) = 0. \quad (8.33)$$

Inserting eqs. (8.27), (8.28) and (8.31) into the 00-component of eq. (8.30) we get the equation

$$\begin{aligned} 0 = & h^2/2 - \Lambda/6 - 6\beta(2h\ddot{h} - \dot{h}^2 + 6h^2\dot{h}) \\ & + 3\lambda(\dot{h} + 2h^2)(6h\ddot{h} + 19h^2\dot{h} - 2\dot{h}^2 - 2h^4). \end{aligned} \quad (8.34)$$

The remaining components are a consequence of this one.

8.4.2 The masses

Linearizing the trace equation (8.32) around the flat spacetime, hence $\Lambda = 0$, gives independently of λ of course, $R = 12\beta\square R$, and the oscillations around the flat spacetime indeed correspond to a mass $m_0 = (-12\beta)^{-1/2}$. This once again confirms the evaluation eq. (8.8).

Now, let us linearize around the ground states eq. (8.25) by inserting $\Lambda = 0$ and $R = \pm\sqrt{6/\lambda} + Z$ into eq. (8.32). It gives

$$Z = \left(-6\beta \mp \sqrt{27\lambda/2}\right)\square Z, \quad (8.35)$$

and, correspondingly, comparing eq. (8.35) with the equation $(\square + m_\pm)Z = 0$, we get

$$m_\pm = \left(6\beta \pm \sqrt{27\lambda/2}\right)^{-1/2}. \quad (8.36)$$

For $\beta \ll -\sqrt{\lambda}$, $m_{\pm}$ is imaginary, and its absolute value differs by a factor $\sqrt{2}$ from m_0 . This is quite analogous to the $\lambda\phi^4$ -theory, cf. section 8.2. Therefore, we concentrate on discussing this range of parameters.

For the ground state for $\Lambda \neq 0$, $\lambda = 0$ we get with $R = -4\Lambda + Z$ just $Z = 12\beta\Box Z$, i.e., mass m_0 eq. (8.8) just as in the case $\lambda = \Lambda = 0$.

Let us generalize this estimate to $L = L(R)$; according to eq. (8.23), $R = R_0 = \text{const.}$ is a ground state if

$$L'(R_0) R_0 = 2 L(R_0) \quad (8.37)$$

holds. It is degenerated if $L'(R_0) = 0$. Now, we linearize around $R = R_0$: $R = R_0 + Z$. For $L''(R_0) = 0$, only $Z = 0$ solves the linearized equation, and $R = R_0$ is a singular solution. For $L''(R_0) \neq 0$ we get the mass

$$m = \left(R_0/3 - L'(R_0)/3L''(R_0) \right)^{1/2} \quad (8.38)$$

meaning the absence of tachyons for real values m . Eq. (8.38) is the analogue to eq. (8.12) for the general Lagrangian $L(R)$.

8.4.3 The Friedmann model

Here we only consider the spatially flat Friedmann model eq. (8.27). Therefore, we can discuss only de Sitter stages with $R < 0$, especially the ground state R_+ eq. (8.25) representing an anti-de Sitter spacetime does not enter our discussion, but R_- does.

Now, let $\Lambda = 0$. Solutions of eq. (8.34) with constant values h are $h = 0$ representing flat spacetime and in the case that $\lambda > 0$ also

$$h = \frac{1}{\sqrt[4]{24\lambda}} \quad (8.39)$$

representing the de Sitter spacetime. These are the non-degenerated ground states $R = 0$ and $R = R_- = -\sqrt{6}/\lambda$, respectively. Eq. (8.34) can be written as

$$0 = h^2(1 - 24\lambda h^4)/2 + h\ddot{h} \left(1/m_0^2 + 18\lambda(\dot{h} + 2h^2)\right) - 6\lambda\dot{h}^3 + \dot{h}^2(45\lambda h^2 - 1/2m_0^2) + 3h^2\dot{h}(1/m_0^2 + 36\lambda h^2). \quad (8.40)$$

First, let us consider the singular curve defined by the vanishing of the coefficient of $\ddot{h}$ in eq. (8.40) in the $h - \dot{h}$ -phase plane. It is, besides $h = 0$, the curve

$$\dot{h} = -2h^2 - 1/18\lambda m_0^2 \quad (8.41)$$

i.e., just the curve

$$R = 1/3\lambda m_0^2 = -4\beta/\lambda \quad (8.42)$$

which is defined by $L'' = 0$, cf. eq. (8.32). This value coincides with R_+ if $\beta = -\sqrt{3\lambda}/8$, this value we need not discuss here. Points of the curve eq. (8.41) fulfil eq. (8.40) for

$$h = \pm 1/18\lambda m_0^3 \sqrt{3} \sqrt{1 - 1/18\lambda m_0^4} \quad (8.43)$$

only, which is not real because of $\lambda \ll m_0^4$. Therefore, the space of solutions is composed of at least two connected components.

Second, for $h = 0$ we have $\dot{h} = 0$ or

$$\dot{h} = -1/12\lambda m^2. \quad (8.44)$$

From the field equation we get: $h = \dot{h} = 0$ implies $h\ddot{h} \geq 0$, i.e. h does not change its sign. We know such a behaviour already from the calculations in

[163], where the same model with $\lambda = 0$ was discussed. In a neighbourhood of eq. (8.44) we can make the ansatz

$$h = -t/12\lambda m_0^2 + \sum_{n=2}^{\infty} a_n t^n \quad (8.45)$$

which has solutions with arbitrary values a_2 . This means: one can change from expansion to subsequent recontraction, but only through the “eye of a needle” eq. (8.44). On the other hand, a local minimum of the scale factor never appears. Further, eq. (8.44) does not belong to the connected component of flat spacetime.

But we are especially interested in the latter one, and therefore, we restrict to the subset $\dot{h} > \dot{h}$ (eq. (8.41)) and need only to discuss expanding solutions $h \geq 0$. Inserting $\dot{h} = 0$,

$$\ddot{h} = h(24\lambda h^4 - l)/(2/m_0^2 + 72\lambda h^2) \quad (8.46)$$

turns out, i.e., $\ddot{h} > 0$ for $h > 1/\sqrt[4]{24\lambda}$ only. All other points in the $h-\dot{h}$ phase plane are regular ones, and one can write $d\dot{h}/dh \equiv \ddot{h}/\dot{h} = F(h, \dot{h})$ which can be calculated by eq. (8.40).

For a concrete discussion let $\lambda \approx 10^2 l_{\text{Pl}}^4$ and $m_0 = 10^{-5} m_{\text{Pl}}$. Then both conditions $\beta \ll -\sqrt{\lambda}$ and $|R_-| < l_{\text{Pl}}^{-2}$ are fulfilled. Now the qualitative behaviour of the solutions can be summarized: There exist two special solutions which approximate the ground state R_- for $t \rightarrow -\infty$. All other solutions have a past singularity $h \rightarrow \infty$. Two other special solutions approximate the ground state R_- for $t \rightarrow +\infty$. Further solutions have a future singularity $h \rightarrow \infty$, and all other solutions have a power-like behaviour for $t \rightarrow \infty$, $a(t)$ oscillates around the classical dust model $a(t) \sim t^{2/3}$. But if we restrict the

initial conditions to lie in a small neighbourhood of the unstable ground state R_- , only one of the following three cases appears:

1. Immediately one goes with increasing values h to a singularity.
2. As a special case: one goes back to the de Sitter stage R_- .
3. The only interesting one: One starts with a finite l_{Pl} -valued inflationary era, goes over to a GUT-valued second inflation and ends with a power-like Friedmann behaviour.

In the last case to be considered here, let $\lambda = 0$, $\Lambda > 0$ and $\beta < 0$. The analogue to eq. (8.40) then reads

$$0 = h^2/2 - \Lambda/6 + (2h\ddot{h} - \dot{h}^2 + 6h^2\dot{h})/2m_0^2. \quad (8.47)$$

Here, always $h \neq 0$ holds, we consider only expanding solutions $h > 0$. For $\dot{h} = 0$ we have

$$\ddot{h} = (\Lambda m_0^2/3 - m_0^2 h^2)/2h. \quad (8.48)$$

For $\ddot{h} = 0$ we have $\dot{h} > m_0^2/6$ and

$$h = (\Lambda/3 + \dot{h}^2/m_0^2)^{1/2} (1 + 6\dot{h}/m_0^2)^{-1/2}. \quad (8.49)$$

Using the methods of [163], where the case $\Lambda = 0$ has been discussed, we obtain the following result: All solutions approach the de Sitter phase $h^2 = \Lambda/3$ as $t \rightarrow \infty$. There exists one special solution approaching $\dot{h} = -m_0^2/6$ for $h \rightarrow \infty$, and all solutions have a past singularity $h \rightarrow 0$. For a sufficiently small value Λ we have again two different inflationary eras in most of all models.

8.5 The generalized Bicknell theorem

In this section we derive a general equivalence theorem between a non-linear Lagrangian $L(R)$ and a minimally coupled scalar field ϕ with a general potential with Einstein's theory. Instead of ϕ we take

$$\psi = \sqrt{2/3} \phi. \quad (8.50)$$

This is done to avoid square roots in the exponents. Then the Lagrangian for the scalar field reads

$$\tilde{L} = \tilde{R}/2 - 3\tilde{g}^{ij}\psi_{;i}\psi_{;j}/4 + V(\psi). \quad (8.51)$$

At ground states $\psi = \psi_0$, defined by $\partial V/\partial\psi = 0$ the effective mass is

$$m = \sqrt{2/3}\sqrt{\partial^2 V/\partial\psi^2}, \quad (8.52)$$

cf. eqs. (8.12) and (8.50). The variation $0 = \delta\tilde{L}/\delta\psi$ gives

$$0 = \partial V/\partial\psi + 3\tilde{g}^{ij}\tilde{\nabla}_i\tilde{\nabla}_j\psi/2 \quad (8.53)$$

and Einstein's equation is

$$\tilde{E}_{ij} = \kappa\tilde{T}_{ij} \quad (8.54)$$

with

$$\kappa\tilde{T}_{ij} = 3\psi_{;i}\psi_{;j}/2 + \tilde{g}_{ij}\left(V(\psi) - \frac{3}{4}\tilde{g}^{ab}\psi_{;a}\psi_{;b}\right). \quad (8.55)$$

Now, let

$$\tilde{g}_{ij} = e^\psi g_{ij}. \quad (8.56)$$

The conformal transformation eq. (8.56) shall be inserted into eqs. (8.53), (8.54) and eq. (8.55). One obtains from eq. (8.53) with

$$\begin{aligned} \psi^{;k} &:= g^{ik}\psi_{;i} \\ \square\psi + \psi^{;k}\psi_{;k} &= -2(e^\psi\partial V/\partial\psi)/3 \end{aligned} \quad (8.57)$$

and from eqs. (8.54), (8.55)

$$E_{ij} = \psi_{;ij} + \psi_{;i}\psi_{;j} + g_{ij} \left(e^\psi V(\psi) - \square\psi - \psi_{;a}\psi^{;a} \right). \quad (8.58)$$

Its trace reads

$$-R = 4e^\psi V(\psi) - 3\square\psi - 3\psi_{;a}\psi^{;a}. \quad (8.59)$$

Comparing with eq. (8.57) one obtains

$$R = R(\psi) = -2e^{-\psi} \partial \left(e^{2\psi} V(\psi) \right) / \partial \psi. \quad (8.60)$$

Now, let us presume $\partial R / \partial \psi \neq 0$, then eq. (8.60) can be inverted as

$$\psi = F(R). \quad (8.61)$$

In the last step, eq. (8.61) shall be inserted into eqs. (8.57), (8.58), (8.59).

Because of

$$F(R)_{;ij} = \partial F / \partial R \cdot R_{;ij} + \partial^2 F / \partial R^2 \cdot R_{;i} R_{;j} \quad (8.62)$$

and $\partial F / \partial R \neq 0$, eq. (8.58) is a fourth-order equation for the metric g_{ij} . We try to find a Lagrangian $L = L(R)$ such that the equation $\delta L \sqrt{-g} / \delta g^{ij} = 0$ becomes just eq. (8.58). For $L' = \partial L / \partial R \neq 0$, eq. (8.30) can be solved to be

$$E_{ij} = -g_{ij}R/2 + g_{ij}L/2L' - g_{ij}\square L'/L' - L'_{;ij}/L'. \quad (8.63)$$

We compare the coefficients of the $R_{;ij}$ terms in eqs. (8.58) and (8.63), this gives

$$\begin{aligned} \partial F / \partial R &= L'' / L', \quad \text{hence} \\ L(R) &= \mu \int_{R_0}^R e^{F(x)} dx + \Lambda_0 \end{aligned} \quad (8.64)$$

with suitable constants Λ_0 , μ , and R_0 , $\mu \neq 0$. We fix them as follows: We are interested in a neighbourhood of $R = R_0$ and require $L'(R_0) = 1/2$.

Otherwise L should be multiplied by a constant factor. Further, a constant translation of ψ can be used to obtain $F(R_0) = 0$, hence $\mu = 1/2$, $L(R_0) = \Lambda_0$, and

$$L'(R_0) = \partial F / \partial R(R_0) / 2 \neq 0. \quad (8.65)$$

With eq. (8.64) being fulfilled, the traceless parts of eqs. (8.58) and (8.64) identically coincide. Furthermore, we have

$$\square L' / L' = \square F + F^i{}_{;i} \quad (8.66)$$

and it suffices to test the validity of the relation

$$e^F V(F(R)) = -R/2 + L/2L'. \quad (8.67)$$

It holds

$$\begin{aligned} 2L' &= e^F, \quad \text{i.e.,} \\ e^{2F} V(F(R)) &= L - Re^F / 2. \end{aligned} \quad (8.68)$$

At $R = R_0$, this relation reads $V(0) = \Lambda_0 - R_0/2$. Applying $\partial/\partial R$ to eq. (8.68) gives just eq. (8.57), and, by the way, $V'(0) = R_0/2 - 2\Lambda_0$. In sum,

$$L(R) = V(0) + R_0/2 + \int_{R_0}^R e^{F(x)} dx / 2, \quad (8.69)$$

where $F(x)$ is defined via $F(R_0) = 0$,

$$\psi = F \left(-2e^{-\psi} \partial(e^{2\psi} V(\psi)) / \partial \psi \right). \quad (8.70)$$

Now, let us go the other direction: Let $L = L(R)$ be given such that at $R = R_0$, $L'L'' \neq 0$. By a constant change of L let $L'(R_0) = 1/2$. Define

$\Lambda_0 = L(R_0)$, $\psi = F(R) = \ln(2L'(R))$ and consider the inverted function $R = F^{-1}(\psi)$. Then

$$V(\psi) = (\Lambda_0 - R_0/2)e^{-2\psi} - e^{-2\psi} \int_0^\psi e^x F^{-1}(x) dx / 2 \quad (8.71)$$

is the potential ensuring the above mentioned conformal equivalence. This procedure is possible at all R -intervals where $L' L'' \neq 0$ holds. For analytical functions $L(R)$, this inequality can be violated for discrete values R only, or one has simply a linear function $L(R)$ being Einstein gravity with Λ -term.

Eq. (8.71) is given here in the form published in [209]. Later it turned out that this integral can be evaluated in closed form as follows:

$$V(\psi) = L(F^{-1}(\psi)) \cdot e^{-2\psi} - \frac{1}{2} F^{-1}(\psi) \cdot e^{-\psi}. \quad (8.72)$$

Examples: 1. Let $L = \Lambda + R^2$, $R_0 = 1/4$, then $4R = e^\psi$ and

$$V(\psi) = \Lambda e^{-2\psi} - 1/16. \quad (8.73)$$

For $\Lambda = 0$, this is proven in Bicknell [29] and Starobinsky and Schmidt [245].

2. Let $L = \Lambda + R/2 + \beta R^2 + \lambda R^3/12$, $R_0 = 0$, hence $\beta \neq 0$ is necessary.

We get

$$\begin{aligned} e^\psi - 1 &= 4\beta R + \lambda R^2/2 \quad \text{and} \\ V(\psi) &= \Lambda e^{-2\psi} + \\ 2\beta\lambda^{-1}e^{-2\psi} &\left(e^\psi - 1 - 16\beta^2(3\lambda)^{-1}((1 + \lambda(e^\psi - l)/8\beta^2)^{3/2} - 1) \right). \end{aligned} \quad (8.74)$$

The limit $\lambda \rightarrow 0$ in eq. (8.74) is possible and leads to

$$V(\psi) = \Lambda e^{-2\psi} - (1 - e^{-\psi})^2/16\beta, \quad (8.75)$$

so we get for $\Lambda = 0$ again the potential from eq. (8.7).

Now, let R_0 be a non-degenerated ground state, hence

$$L(R) = \Lambda_0 + (R - R_0)/2 + L''(R_0)(R - R_0)^2/2 + \dots \quad (8.76)$$

with $L''(R_0) \neq 0$ and $\Lambda_0 = R_0/4$, cf. subsection 8.3.1. Using eq. (8.71) we get $V'(0) = 0$ and

$$V''(0) = R_0/2 - 1/(4L''(R_0)). \quad (8.77)$$

Inserting this into eq. (8.52) we exactly reproduce eq. (8.38). This fact once again confirms the estimate eq. (8.38) and, moreover, shows it to be a true analogue to eq. (8.12). To understand this coincidence one should note that at ground states, the conformal factor becomes a constant = 1.

8.6 On Ellis' programme within homogeneous world models

For the non-tachyonic curvature squared action we show that the expanding Bianchi-type I models tend to the dust-filled Einstein-de Sitter model for t tending to infinity if the metric is averaged over the typical oscillation period. Applying a conformal equivalence between curvature squared action and a minimally coupled scalar field, which holds for all dimensions > 2 , the problem is solved by discussing a massive scalar field in an anisotropic cosmological model.

8.6.1 Ellis' programme

Ellis [70] has asked in 1984 on which length scale the Einstein field equation is valid. On laboratory or on cosmic distances? Here we extend this question to cover microscopic scales, too. This is a real problem as one knows:

1. An averaging procedure does not commute with a non-linear differential operator as the Einstein tensor is.
2. Einstein's theory is well tested at large distances $\gg 1$ cm.
3. The ultraviolet divergencies of Einstein gravity can be removed by adding curvature squared terms to the action, see Weinberg [267]; this is a microscopic phenomenon and forces to prefer a curvature squared action already on the level of classical field theory, as we are concerned here.

Now, we propose a synthesis of 1., 2., and 3. as follows: microscopically, we take

$$L_g = (R/2 - l^2 R^2)/8\pi G \quad (8.78)$$

the minus sign before the R^2 -term shows that we consider the non-tachyonic case only. By an averaging procedure we get Einstein gravity on large scales $\gg l \approx 10^{-28}$ cm. As we are dealing with homogeneous cosmological models, the average is taken over the time coordinate only: We will show: the curvature squared contribution represents effectively dust in the asymptotic region $t \rightarrow \infty$. For spatially flat Friedmann models this was already proven in Müller and Schmidt [163], here we generalize to models with less symmetry. In the present paper we consider only the vacuum case

$$\delta L_g \sqrt{-\det g_{ij}} \delta g^{kl} = 0, \quad (8.79)$$

and therefore, we interpret the effectively obtained dust as invisible gravitating matter necessary to get a spatially flat universe.

We conjecture that additional matter contributions, that means usual dust plus radiation, do not alter the result qualitatively. The general expectation for Lagrangian eq. (8.78) is the following: starting at die Planck

era

$$R_{ijkl}R^{ijkl} \approx 10^{131} \text{ cm}^{-4}, \quad (8.80)$$

the R^2 -term is dominant, the inflationary de Sitter phase is an attractor; its appearance becomes very probable, see [207] and [245]; the R -term in eq. (8.78) yields a parametric decay of the value of R and one turns over to the region $t \rightarrow \infty$ where R^2 gives just dust in the mean.

In Carfora and Marzuori [50] another approach to Ellis' programme was initiated: a smoothing out of the spatially closed 3-geometry in the direction of the spatial Ricci-tensor leads to different values of mean mass density before and after the smoothing out procedure.

8.6.2 The massive scalar field in a Bianchi-type I model

Consider a minimally coupled scalar field ϕ in a potential $V(\phi)$, where $8\pi G = 1$ is assumed,

$$L = R/2 - \frac{1}{2}\phi_{;i}\phi^{;i} + V(\phi). \quad (8.81)$$

We suppose $V(\phi)$ to be a C^3 -function and $V(0) = 0$ to be a local quadratic minimum of the potential V , i.e., $V'(0) = 0, V''(0) = m^2, m > 0$. To describe the asymptotic behaviour $\phi \rightarrow 0$ as $t \rightarrow \infty$ it suffices to use $V(\phi) = m^2\phi^2/2$ because the higher order terms do not affect it. This statement can be proved as follows: For each $\varepsilon > 0, \varepsilon < m$, there exists a $\phi_0 > 0$ such that for all ϕ with $|\phi| < \phi_0$ it holds

$$(m - \varepsilon)^2\phi^2 \leq 2V(\phi) \leq (m + \varepsilon)^2\phi^2. \quad (8.82)$$

Then all further development is enclosed by inequalities with $m \pm \varepsilon, \varepsilon \rightarrow 0$.

Here, we concentrate on a cosmological model of Bianchi-type I. In Misner parametrization it can be written as

$$ds^2 = dt^2 - e^{2\alpha} [e^{2(s+\sqrt{3}r)} dx^2 + e^{2(s-\sqrt{3}r)} dy^2 + e^{-4s} dz^2]. \quad (8.83)$$

The functions ϕ , α , r , s depend on t only. $\dot{\alpha} = d\alpha/dt = h$ is the Hubble parameter and $\eta = u/h$, $u = (\dot{r}^2 + \dot{s}^2)^{1/2}$ the anisotropy parameters. It holds $0 \leq \eta \leq 1$, $\eta = 0$ represents the isotropic model, and $\eta = 1$ gives $\phi \equiv 0$, therefore, we consider only the case $0 < \eta < 1$ in the following. $h = 0$ is possible for Minkowski spacetime only, and the time arrow is defined by $h > 0$, i.e., we restrict to expanding solutions.

In Lukash and Schmidt [145] it was shown that all such solutions tend to Minkowski spacetime for $t \rightarrow \infty$. More detailed: At $t = 0$, let $\alpha = r = s = 0$ by a coordinate transformation. Then for each prescribed quadruple $(\dot{r}, \dot{s}, \phi, \dot{\phi})$ at $t = 0$ the integration of the relevant system up to $t \rightarrow \infty$ is possible with $r, s, \phi, \dot{\phi}, h, \dot{h}$ tending to zero in that limit. The relevant equations are

$$(\phi^2 + \dot{\phi}^2 + u^2)^{1/2} = h \quad (8.84)$$

$$\ddot{\phi} + 3h\dot{\phi} + \phi = 0 \quad (8.85)$$

$$\dot{r} = C_r e^{-3\alpha}, \quad \dot{s} = C_s e^{-3\alpha}. \quad (8.86)$$

We put the mass $m = 1$; C_r, C_s are constants. Now, we consider this limit in more details. Up to now the following is known: For the isotropic models $u \equiv 0$, $e^\alpha = a$ one knows that the asymptotic behaviour is given by oscillations around $a \sim t^{2/3}$ i.e., we get the Einstein-de Sitter model in the mean, and the effective equation of state is that of dust, cf. e.g. Starobinsky [243]. In Gottlöber 1977 [92] this is generalized to a special class

of inhomogeneous, nearly isotropic models. After averaging over space and oscillation period one gets as effective equation of state

$$p \sim \rho/a^2 \quad (8.87)$$

i.e., also dust in the limit $t \rightarrow \infty$.

The question for the Bianchi-type I model is now: does one get the energy-momentum tensor of an ideal fluid in the mean, or will there appear a strongly anisotropic pressure? The first question to be answered is about the averaging procedure: From eq. (8.85) and the fact that $h \rightarrow 0$ as $t \rightarrow \infty$ one gets a fixed oscillation period: let $t = t_n$ be the n -th local extremum of $\phi(t)$, then

$$\lim_{n \rightarrow \infty} t_{n+1} - t_n = \pi. \quad (8.88)$$

Therefore, we average the metric about this period of time. Let us denote $r(t_n)$ by $r_n, \dots$. For a fixed finite time it is ambiguous how to perform this average, but in the limit $t \rightarrow \infty$ one can describe the system by adiabatic or parametric deviations from pure $\phi \sim \cos t$ -oscillations, and the averaging procedure consists of constructing monotonous smooth curves $\bar{\alpha}(t)$ fulfilling $\bar{\alpha}(t_n) = \alpha_n, \dots$, and the effective energy-momentum tensor is obtained from eq. (8.78) with $\alpha = \bar{\alpha}(t), \dots$ using the Einstein equation.

Derivating eq. (8.84) one gets with eq. (8.85)

$$0 \geq \dot{h} = 3\phi^2 - 3h^2 \geq -3h^2 \quad (8.89)$$

and therefore,

$$h_n \geq h_{n+1} > h_n - 3\pi h_n^2. \quad (8.90)$$

Let

$$h \geq \frac{1}{3\pi(n + n_0)} \quad (8.91)$$

be valid for one value $n = n_1$ then, by induction, this holds true also for all larger values n , i.e.,

$$h(t) \geq \frac{1}{3(t + t_0)} \quad (8.92)$$

for some t_0 , let $t_0 = 0$ subsequently. Derivating eq. (8.86) we get with eq. (8.84)

$$\dot{\eta} = -3\eta(1 - \eta^2)h \cdot \frac{\phi^2}{\phi^2 + \dot{\phi}^2}. \quad (8.93)$$

The last factor can be substituted by its mean value, which equals $1/2$. From eq. (8.92) one can see that for initial conditions $0 < \eta < 1$ as we met here, eq. (8.93) leads to $\eta \rightarrow 0$ as $t \rightarrow \infty$. Knowing this, we can perform a stronger estimate for h , because in the mean, $\dot{h} = -3h^2/2$, i.e., in the leading order we have $h(t) = 2/3t$ and, using eq. (8.93), we get

$$\dot{\eta} = -\eta/t, \quad \eta = \eta_0/t. \quad (8.94)$$

Remember: the inflationary era diminishes η exponentially with increasing time, but here we consider only the asymptotic region. In sum we get for metric (8.83)

$$e^\alpha = t^{2/3}, \quad r = -C_r/t, \quad s = -C_s/t. \quad (8.95)$$

Inserting eq. (8.95) into eq. (8.83) we get, via the Einstein equation, a diagonal energy-momentum tensor with $\rho = T_{00} = 4/3t^2$ and pressure of the order $(C_r^2 + C_s^2)/t^4$. The effective equation of state is $p \sim \rho/t^2 \sim \rho/a^3$, i.e., also dust in the limit $t \rightarrow \infty$ as eq. (8.87). From the details of the paper [145] one can see that for almost all models the end of the inflationary stage is just the beginning of the Einstein-de Sitter stage. A similar result occurs if not the metric, but ρ and p will be averaged, cf. Gottlöber [92].

Result: For the minimally coupled scalar field in a potential V which has a single quadratic minimum at $V = 0$ all expanding Bianchi-type I

cosmological solutions tend to the Einstein-de Sitter model for $t \rightarrow \infty$ if the metric is averaged over the oscillation period.

8.6.3 The generalized equivalence

Some types of a conformal equivalence theorem between fourth-order gravity and minimally coupled scalar fields are obtained in [209], [211]. This theorem was also independently obtained by Ferraris [76], see also Jakubiec and Kijowski [120], Goenner [90]. All of them are restricted to 4-dimensional spacetimes. On the other hand, both R^2 -terms and scalar fields have been discussed for higher-dimensional spacetimes, cf. e.g. Ishihara [116].

Therefore, it is worth mentioning that this conformal equivalence theorem can be formulated for arbitrary dimensions $n > 2$: Let

$$\tilde{\mathcal{L}} = \tilde{R}/2 - \frac{1}{2}\tilde{g}^{ij}\phi_{|i}\phi_{|j} + V(\phi) \quad (8.96)$$

and

$$\tilde{g}_{ij} = e^{\lambda\phi}g_{ij}, \quad \lambda = \frac{2}{(n-1)(n-2)} \quad (8.97)$$

be the conformally transformed metric. Then the solutions of the variation of eq. (8.96) are transformed by eq. (8.97) to the solutions of the variation of $L = L(R)$, where

$$R = -2e^{\lambda\phi} \left(\frac{nV}{n-2} + \mu \frac{dV}{d\phi} \right), \quad \mu = \sqrt{\frac{n-1}{n-2}} \quad (8.98)$$

is supposed to be locally, i.e. near $R = R_0$, invertible as

$$\begin{aligned} \phi &= F(R), \quad F(R_0) = 0, \quad F'(R_0) \neq 0 \\ \mathcal{L}(R) &= \frac{1}{2}R_0 + V(0) + \frac{1}{2} \int_{R_0}^R e^{F(x)/\mu} dx. \end{aligned} \quad (8.99)$$

The inverse direction is possible provided $L'(R) \cdot L''(R) \neq 0$, cf. [209], [211] for details with $n = 4$. Other cosmological models with higher dimensions are in [32], [119], and [190]. Two-dimensional models are discussed in [154] and [229].

8.6.4 The fourth-order gravity model

Now we come to the question posed in the introduction: Let $L(R)$ be a C^3 -function fulfilling $L(0) = 0$, $L'(0) \cdot L''(0) < 0$. Then we can write

$$\mathcal{L}(R) = \frac{R}{2} + \beta R^2 + O(R^3), \quad \beta < 0. \quad (8.100)$$

We consider the Bianchi-type I vacuum solutions which start in a neighbourhood of the Minkowski spacetime and ask for the behaviour as $t \rightarrow \infty$. Applying the equivalence theorem cited in subsection 8.6.3 we arrive at the models discussed in subsection 8.6.2, and this is applicable for $|R|$ being small enough. The conformal factor depends on t only, and therefore, the space of Bianchi-type I models will not be leaved, and we can formulate the following: In a neighbourhood of Minkowski spacetime, all Bianchi-type I models which represent a stationary point of the action eq. (8.100), can be integrated up to $t \rightarrow \infty$ or $-\infty$, let it be $+\infty$. One singular solution is the Kasner solution and all other solutions undergo isotropization and have an averaged equation of state $p = 0$ for $t \rightarrow \infty$.

Let us conclude: Anderson [6] discussed the possibility that curvature squared terms give an effective contribution to the energy density of a Friedmann model. This could explain the discrepancy between the observed mean mass density of about 1/10 the critical one and the predicted one, from inflationary cosmology, nearly the critical one. Here, we have shown that also for

a large initial anisotropy the oscillating curvature squared contributions give just dust in the mean and not an equally large anisotropic pressure as one could have expected. The next step would be to look for a generalization of this fact to inhomogeneous cosmological models. Further fourth-order gravity models can be seen in [11], [48], [49], [61], [74], [110], [111], [115], [124], [128], [139], [193], [214] and [246]. Another approach to averaging procedures, i.e., to Ellis' programme, is due to Zalaletdinov [275].

9 Friedmann models with de Sitter and power-law inflation

Following [216], we consider the spatially flat Friedmann model

$$ds^2 = dt^2 - a^2(t)(dx^2 + dy^2 + dz^2). \quad (9.1)$$

For $a \approx t^p$, especially, if $p \geq 1$, this is called power-law inflation. For the Lagrangian

$$L = R^m \quad \text{with} \quad p = -(m-1)(2m-1)/(m-2) \quad (9.2)$$

power-law inflation is an exact solution, as it is for Einstein gravity with a minimally coupled scalar field Φ in an exponential potential

$$V(\Phi) = \exp(\mu\Phi) \quad (9.3)$$

and also for the higher-dimensional Einstein equation with a special Kaluza-Klein ansatz. The synchronized coordinates are not adapted to allow a closed-form solution, so we write

$$ds^2 = a^2(Q^2(a) da^2 - dx^2 - dy^2 - dz^2). \quad (9.4)$$

The general solutions reads

$$Q(a) = (a^b + C)^{f/b} \quad (9.5)$$

with free integration constant C , where $C = 0$ gives exact power-law inflation, and m -dependent values b and f :

$$f = -2 + 1/p, \quad b = (4m - 5)/(m - 1). \quad (9.6)$$

Further, special solutions for the closed and open Friedmann model are found.

The de Sitter spacetime

$$ds^2 = dt^2 - e^{2Ht}(dx^2 + dy^2 + dz^2), \quad H \neq 0 \quad (9.7)$$

is the spacetime being mainly discussed to represent the inflationary phase of cosmic evolution. However, a spacetime defined by

$$ds^2 = dt^2 - |t|^{2p}(dx^2 + dy^2 + dz^2), \quad p \neq 0 \quad (9.8)$$

enjoys increasing interest for these discussions, too. Especially, eq. (9.8) with $p \geq 1$, $t > 0$ is called power-law inflation; and with $p < 0$, $t < 0$ it is called polar inflation.

We summarize some differential-geometrical properties of both de Sitter and power-law/polar inflation in section 9.1 and show, from which kind of scale-invariant field equations they arise in section 9.2; we give the complete set of solutions for the spatially flat and special solutions for the closed and open Friedmann models in closed form for field equations following from the Lagrangian $L = R^m$ in section 9.3, and discuss the results in section 9.4 under the point of view that power-law inflation is an attractor solution of the corresponding field equations.

9.1 Differential-geometrical properties

Eq. (9.8) defines a self-similar spacetime: if we multiply the metric ds^2 by an arbitrary positive constant a^2 , then the resulting $d\hat{s}^2 = a^2 ds^2$ is isometric to ds^2 . Proof: We perform a coordinate transformation $\hat{t} = at$, $\hat{x} = b(a, p)x$... On the other hand, the de Sitter spacetime eq. (9.7) is not self-similar, because it has a constant non-vanishing curvature scalar.

Power-law inflation is intrinsically time-oriented. Proof: The gradient of the curvature scalar defines a temporal orientation. On the other hand, the expanding ($H > 0$) and the contracting ($H < 0$) de Sitter spacetime can be transformed into each other by a coordinate transformation, because both of them can be transformed to the closed Friedmann universe with scale factor $\cosh(Ht)$, which is an even function of t . This property is connected with the fact that eq. (9.8) gives a global description, whereas eq. (9.7) gives only a proper subset of the full de Sitter spacetime.

For $p \rightarrow \infty$, eq. (9.8) tends to eq. (9.7). Such a statement has to be taken with care, even for the case with real functions. Even more carefully one has to deal with spacetimes. The most often used limit — the Geroch-limit [86] of spacetimes — has the property that a symmetry, here we take it to be self-similarity, of all the elements of the sequence must also be a symmetry of the limit.

From this it follows that the Geroch limit of spacetimes eq. (9.8) with $p \rightarrow \infty$ cannot be unique, moreover, it is just the one-parameter set eq. (9.7) parametrized by arbitrary values $H > 0$.

9.2 Scale-invariant field equations

A gravitational field equation is called scale-invariant, if to each solution ds^2 and to each positive constant c^2 the resulting homothetically equivalent metric $d\hat{s}^2 = c^2 ds^2$ is also a solution. One example of such field equations is that one following from eq. (9.2). Moreover, no Lagrangian $L = L(R)$ gives rise to a scale-invariant field equation which is not yet covered by eq. (9.2) already.

Secondly, for

$$L = R/16\pi G - \frac{1}{2}g^{ij}\Phi_{,i}\Phi_{,j} + V_0 \exp(\mu\Phi), \quad (9.9)$$

we assume $8\pi G = 1$ henceforth, the homothetic transformation has to be accompanied by a suitable translation of Φ to ensure scale-invariance. For $\mu \neq 0$, the value of V_0 can be normalized to 1, 0 or -1 .

A third example is the following: for the Kaluza-Klein ansatz eq. (9.10) the N -dimensional Einstein equation $R_{AB} = 0$ is scale-invariant. We take the ansatz as

$$dS^2 = ds^2(x^i) + W(x^i)d\tau^2(x^\alpha), \quad (9.10)$$

where $i, j = 0, \dots, 3$; $\alpha, \beta = 4, \dots, N-1$; $A, B = 0 \dots N-1$, and restrict to the warped product of the $(N-4)$ -dimensional internal space $d\tau^2$ with 4-dimensional spacetime ds^2 .

Let us consider the limits $m \rightarrow \infty$ and $m \rightarrow 0$ of eq. (9.2). One gets $L = \exp(R/R_0)$ and $L = \ln(R/R_0)$, respectively. Both of them give rise to a field equation which is not scale-invariant: a homothetic transformation changes also the reference value R_0 . For the second case this means a change of the cosmological constant. So we have a similar result as before: the limits exist, but they are not unique.

9.3 Cosmological Friedmann models

We consider the closed and open model in subsection 9.3.1. and the spatially flat model in subsection 9.3.2.

9.3.1 The closed and open models

Let us start with the closed model. We restrict ourselves to a region where one has expansion, so we may use the cosmic scale factor as timelike coordinate:

$$ds^2 = a^2 \left(Q^2(a) da^2 - d\sigma^2 \right), \quad Q(a) > 0, \quad (9.11)$$

where $d\sigma^2$, defined by

$$d\sigma^2 = dr^2 + \sin^2 r d\Omega^2, \quad d\Omega^2 = d\theta^2 + \sin^2 \theta d\psi^2 \quad (9.12)$$

is the positively curved 3-space of constant curvature. It holds: if the 00-component of the field equation, here: eq. (9.15) below, is fulfilled, then all other components are fulfilled, too. Such a statement holds true for all Friedmann models and “almost all sensible field theories”. With ansatz eqs. (9.11), (9.12) we get via

$$R_0^0 = 3(a^{-4}Q^{-2} + a^{-3}Q^{-3}dQ/da) \quad (9.13)$$

and

$$R = 6(a^{-3}Q^{-3}dQ/da - a^{-2}) \quad (9.14)$$

the result: the field equation following from the Lagrangian (9.2) is fulfilled for metric (9.11), (9.12), if and only if

$$mRR_0^0 - R^2/2 + 3m(m-1)Q^{-2}a^{-3}dR/da = 0 \quad (9.15)$$

holds. For $m = 1$, which represents Einstein’s theory, no solution exists. For all other values m , eqs. (9.13), (9.14) and (9.15) lead to a second order equation for $Q(a)$.

We look now for solutions with vanishing R_0^0 , i.e., with eq. (9.13) we get

$$Q = C/a, \quad C = \text{const.} > 0. \quad (9.16)$$

By the way, metric (9.11), (9.12) is self-similar if and only if eq. (9.16) holds. From eqs. (9.14), (9.16) we get

$$R = D/a^2, \quad D = -6(l + 1/C^2). \quad (9.17)$$

We insert eqs. (9.16), (9.17) into eq. (9.15) and get

$$C = (2m^2 - 2m - 1)^{1/2}, \quad (9.18)$$

which fulfils eq. (9.16) if

$$m > (1 + \sqrt{3})/2 \quad \text{or} \quad m < (1 - \sqrt{3})/2 \quad (9.19)$$

holds. Inserting eq. (9.16) and eq. (9.18) into eq. (9.11) and introducing synchronized coordinates we get as a result: if eq. (9.19) holds, then

$$ds^2 = dt^2 - t^2 d\sigma^2 / (2m^2 - 2m - 1) \quad (9.20)$$

is a solution of the fourth-order field equation following from Lagrangian (9.2). It is a self-similar solution, and no other self-similar solution describing a closed Friedmann model exists.

For the open Friedmann model all things are analogous, one gets for

$$(1 - \sqrt{3})/2 < m < (1 + \sqrt{3})/2 \quad (9.21)$$

and with $\sinh r$ instead of $\sin r$ in eq. (9.12) the only self-similar open solution, which is flat for $m \in \{0, 1\}$,

$$ds^2 = dt^2 - t^2 d\sigma^2 / (-2m^2 + 2m + 1). \quad (9.22)$$

9.3.2 The spatially flat model

The field equation for the spatially flat model can be deduced from that one of a closed model by a limiting procedure as follows: we insert the transformation $r \rightarrow \varepsilon r$, $a \rightarrow a/\varepsilon$, $Q \rightarrow Q\varepsilon^2$ and apply the limit $\varepsilon \rightarrow 0$ afterwards. One gets via

$$\lim_{\varepsilon \rightarrow 0} \sin(\varepsilon r) = r \quad (9.23)$$

the metric

$$ds^2 = a^2 \left(Q^2(a) da^2 - dx^2 - dy^2 - dz^2 \right) \quad (9.24)$$

with unchanged eqs. (9.13) and (9.15), whereas eq. (9.14) yields

$$R = 6a^{-3}Q^{-3}dQ/da. \quad (9.25)$$

The trivial solutions are the flat Minkowski spacetime and the model with constant value of Q , i.e., $R = 0$, a valid solution for $m > 1$ only, which has the geometry of Friedmann's radiation model.

Now, we consider only regions with non-vanishing curvature scalar. For the next step we apply the fact that the spatially flat model has one symmetry more than the closed one: the spatial part of the metric is self-similar. In the coordinates eq. (9.24) this means that each solution $Q(a)$ may be multiplied by an arbitrary constant. To cancel this arbitrariness, we define a new function

$$P(a) = d(\ln Q)/da. \quad (9.26)$$

We insert eqs. (9.13), (9.25) into eq. (9.15), and then eq. (9.26) into the resulting second order equation for Q . We get the first order equation for P

$$0 = m(m-1)dP/da + (m-1)(1-2m)P^2 + m(4-3m)P/a. \quad (9.27)$$

As it must be the case, for $m = 0, 1$, only $P = 0$ is a solution. For $m = 1/2$,

$$P \sim a^5 \quad \text{and} \quad Q = \exp(c a^6), \quad (9.28)$$

where c denotes an integration constant. For the other values m we define $z = aP$ as new dependent and $t = \ln a$ as new independent variable. Eq. (9.27) then becomes

$$0 = dz/dt + gz^2 - bz, \quad g = l/m - 2, \quad b = (4m - 5)/(m - 1). \quad (9.29)$$

For $m = 5/4$ we get

$$z = -5/(6t - c), \quad \text{i.e.,} \quad P = -5/(6a \ln(a/c)), \quad Q = (\ln(a/c))^{-5/6}. \quad (9.30)$$

For the other values we get

$$z = f/(\exp(-bt + c) - 1), \quad f = -b/g, \quad P = f/(e^c a^{1-b} - a) \\ Q = (\pm a^b + c)^{1/g}. \quad (9.31)$$

For $m \rightarrow 1/2$ we get $b \rightarrow 6$ and $1/g \rightarrow \infty$; for $m \rightarrow 5/4$ we get $b \rightarrow 0$ and $1/g \rightarrow -5/6$, so the two special cases could also have been obtained by a limiting procedure from eq. (9.31).

Metric (9.24) with eq. (9.31) can be explicitly written in synchronized coordinates for special examples only, see e.g. Burd and Barrow [45].

9.4 Discussion of power-law inflation

We have considered scale-invariant field equations. The three examples mentioned in section 9.2 can be transformed into each other by a conformal transformation of the four-dimensional spacetime metric. The parameters of eqs.

(9.2) and (9.9) are related by $\sqrt{3}\mu = \sqrt{2}(2-m)/(m-1)$, cf. [214], a similar relation exists to the internal dimension in eq. (9.10), one has

$$m = 1 + 1/\left\{1 + \sqrt{3}[1 + 2/(N-4)]^{\pm 1/2}\right\}, \quad (9.32)$$

for details see [31]. The necessary conformal factor is a suitable power of the curvature scalar eq. (9.2). A further conformal transformation in addition with the field redefinition $\theta = \tanh \Phi$ leads to the conformally coupled scalar field θ in the potential

$$(1 + \theta)^{2+\mu} (1 - \theta)^{2-\mu}, \quad (9.33)$$

the conformal factor being $\cosh^4 \Phi$, cf. [213]. So, equations stemming from quite different physical foundations are seen to be equivalent. We have looked at them from the point of view of self-similar solutions and of limiting processes changing the type of symmetry.

The general solution for the spatially flat Friedmann models in fourth-order gravity eq. (9.2), eqs. (9.24), (9.31) can be written for small c in synchronized coordinates as follows

$$ds^2 = dt^2 - a^2(t)(dx^2 + dy^2 + dz^2), \quad a(t) = t^p(1 + \varepsilon t^{-bp} + O(\varepsilon^2)). \quad (9.34)$$

$\varepsilon = 0$ gives exact power-law inflation with $p = (m-1)(2m-1)/(2-m)$ and $b = (4m-5)/(m-1)$. We see: in the range $1 < m < 2$, power-law inflation is an attractor solution within the set of spatially flat Friedmann models for $m \geq 5/4$ only, because otherwise b would become negative. The generalization to polar inflation with $m > 2$ and $p < 0$ is similar.

9.5 The cosmic no hair theorem

Following [129], we now discuss the cosmic no hair theorem, which tells under which circumstances the de Sitter spacetime represents an attractor solution

within the set of other nearby solutions. This property ensures the inflationary model to be a typical solution. The notion “cosmic no hair theorem” is chosen because of its analogous properties to the “no hair theorem” for black holes, for which one can cf. [194].

After a general introduction we restrict our consideration to spatially flat Friedmann models. In this section, we choose gravitational Lagrangians

$$R \square^k R \sqrt{-g} \tag{9.35}$$

and linear combinations of them. They are motivated from trials how to overcome the non-renormalizability [248] of Einstein’s theory of gravity. Results are: For arbitrary k , i.e., for arbitrarily large order $2k + 4$ of the field equation, one can always find examples where the attractor property takes place. Such examples necessarily need a non-vanishing R^2 -term. The main formulas do not depend on the dimension, so one gets similar results also for 1+1-dimensional gravity and for Kaluza-Klein cosmology.

9.5.1 Introduction to no hair theorems

Over the years, the notion “no hair conjecture” drifted to “no hair theorem” without possessing a generally accepted formulation or even a complete proof. Several trials have been made to formulate and prove it at least for certain special cases. They all have the overall structure: “For a geometrically defined class of spacetimes and physically motivated properties of the energy-momentum tensor, all the solutions of the gravitational field equation asymptotically converge to a space of constant curvature.”

The paper Weyl [269] is cited in [13] with the phrase: “The behaviour of every world satisfying certain natural homogeneity conditions in the large

follows the de Sitter solution asymptotically.” to be the first published version of the no hair conjecture. Barrow and Götz [13] apply the formulation: “All ever-expanding universes with $\Lambda > 0$ approach the de Sitter spacetime locally.” The first proof of the stability of the de Sitter solution, in this case within the steady-state theory, is due to Hoyle and Narlikar 1963 [112].

The probability of inflation is large if the no hair theorem is valid, cf. Alshuler [1]. Peter, Polarski, Starobinsky [181] and Kofman, Linde, Starobinsky [130] compared the double-inflationary models with cosmological observations. Brauer, Rendall, Reula [40] discussed the no hair conjecture within Newtonian cosmological models. Hübner and Ehlers [113] and Burd [44] considered inflation in an open Friedmann universe and have noted that inflationary models need not to be spatially flat.

Gibbons and Hawking 1977 [88] have found two of the earliest strict results on the no hair conjecture for Einstein’s theory, cf. also Hawking, Moss [109] and Demianski [62]. Barrow 1986 [12] gave examples that the no hair conjecture fails if the energy condition is relaxed and points out, that this is necessary to solve the graceful exit problem. He uses the formulation of the no hair conjecture “in the presence of an effective cosmological constant, stemming e.g. from viscosity, the de Sitter spacetime is a stable asymptotic solution”.

Usually, energy inequalities are presumed for formulating the no hair conjecture. Nakao et al. [168] found some cases where it remains valid also for negative Abbott-Deser mass. The latter goes over to the well-known ADM-mass [7] for $\Lambda \rightarrow 0$. In Murphy [165], viscosity terms as source are considered to get a singularity-free cosmological model. In the eighties, these

non-singular models with viscosity where reinterpreted as inflationary ones, cf. Oleak [173].

In the three papers [186], Prigogine et al. developed a phenomenological model of particle and entropy creation. It allows particle creation from space-time curvature, but the inverse procedure, i.e. particle decay into spacetime curvature is forbidden. This breaks the $t \rightarrow -t$ -invariance of the model. Within that model, the expanding de Sitter spacetime is an attractor solution independently of the initial fluctuations; this means, only the expanding de Sitter solution is thermodynamically possible.

Vilenkin 1992 [262] and Borde [33], [34] discussed future-eternal inflating universe models; they must have a singularity if the condition D: “There is at least one point p such that for some point q to the future of p the volume of the difference of the pasts of p and q is finite” is fulfilled. Mondaini and Vilar [157] have considered recollapse and the no hair conjecture in closed higher-dimensional Friedmann models.

In Shiromizu et al. [237], Shibata et al. [235] and Chiba, Maeda [52] the following argument is given: If the matter distribution is too clumpy, then a large number of small black holes appears. Then one should look for an inflationary scenario where these black holes are harmless. They cannot clump together to one giant black hole because of the exponential expansion of the universe; this explains the existing upper bound of the mass of black holes in the quasi-de Sitter model: above

$$M_{\text{crit}} = \frac{1}{3\sqrt{\Lambda}} \quad (9.36)$$

there do not exist horizons; this restriction is called cosmic hoop conjecture. The paper [52] from 1994 is entitled “Cosmic hoop conjecture?”, and that

conjecture is formulated as: when an apparent horizon forms in a gravitational collapse, the matter must be sufficiently compactified such that the circumference C satisfies the condition

$$C < 4\pi M < 4\pi M_{\text{crit}} , \quad (9.37)$$

where M is the Abbott–Deser mass of the collapsed body.

Xu, Li and Liu [272] proved the instability of the anti-de Sitter spacetime against gravitational waves and dust matter perturbations in 1994. Coley and Tavakol [54] discussed the robustness of the cosmic no hair conjecture under using the concept of the structural stability. Sirousse-Zia [239] considered the Bianchi type IX model in Einstein’s theory with a positive Λ -term and got an asymptotic isotropization of the mixmaster model. She uses methods of Belinsky et al. [22]. Müller [162] used $L = R^2$ and discussed the power-asymptotes of Bianchi models. Barrow and Sirousse-Zia [16] discussed the mixmaster R^2 -model and the question, under which conditions the Bianchi type IX model becomes asymptotic de Sitter. In Yokoyama, Maeda [273] and Cotsakis et al. [57] the no hair conjecture for Bianchi type IX models and Einstein’s theory with a cosmological term are discussed; they get R^2 –inflation in anisotropic universe models and typically, an initial anisotropy helps to enhance inflation. They got some recollapsing solutions besides those converging to the de Sitter solution. In Spindel [241] also general Bianchi type I models in general dimensions are considered.

Breizman et al. 1970 [41] considered

$$L = R + \alpha R^{4/3} \quad (9.38)$$

to get a singularity-free model, the solutions are of a quasi de Sitter type. The papers Barrow [12] from 1987, Pollock [185], Müller et al. [164] and

Mijic et al. [155] consider the no hair conjecture for R^2 models, they use the formulation “asymptotical de Sitter, at least on patch”. The restriction “on patch” is not strictly defined but refers to a kind of local validity of the statement, e.g., in a region being covered by one single synchronized system of reference in which the spatial curvature is non-positive and the energy conditions are fulfilled. The Starobinsky model is one of those which does not need an additional inflaton field to get the desired quasi de Sitter stage.

One should observe a notational change: Originally,

$$L = R + aR^2 \ln R \quad (9.39)$$

was called Starobinsky model, whereas

$$L = R + aR^2 \quad (9.40)$$

got the name “improved Starobinsky model” - but now the latter carries simply the name “Starobinsky model”, see e.g. Starobinsky 1983 [244], Hwang [114], Gottlöber et al. [95] and Amendola et al. [3] and [4]. For the inflationary phase, both versions are quite similar. A further result of the papers [12] and [185] is that by the addition of a cosmological term, the Starobinsky model leads naturally to double inflation. Let us comment this result: It is correct, but one should add that this is got at the price of getting a “graceful exit problem”, by this phrase there is meant the problem of how to finish the inflationary phase dynamically - in the Starobinsky model this problem is automatically solved by the fact that the quasi de Sitter phase is a transient attractor only.

The papers Maeda [150], Maeda et al. [152] and Barrow, Saich [15] discuss the no hair conjecture within R^2 -models and found inflation as a transient

attractor in fourth-order gravity. The papers Feldman [75], Rogers, Isaacson [192] and [220] discuss the stability of inflation in R^2 -gravity. The papers Bičák, Podolsky [27], Cotsakis, Flessas [58], Borzeszkowski, Treder [36] and Contreras et al. [56] discuss generalized cosmic no hair theorems for quasi exponential expansion.

In Starobinsky 1983 [244] the no hair theorem for Einstein's theory with a positive Λ -term is tackled by using a power series expansion as ansatz to describe a general spacetime. This includes a definition of an asymptotic de Sitter spacetime even for inhomogeneous models. However, the convergence of the sequence is not rigorously proven. In [245], the ansatz [244] was generalized to consider also the cosmic no hair theorem for $L = R^2$. Shiromizu et al. [237] discussed an inflationary inhomogeneous scenario and mentioned the open problem how to define asymptotic de Sitter spacetimes.

Jensen, Stein-Schabes [122] consider the no hair theorem for a special class of inhomogeneous models and give partial proofs. Morris [161] considers inhomogeneous models for $R + R^2$ -cosmology. In Calzetta, Sakellariadou [47] inflation in inhomogeneous but spherically symmetric cosmological models is obtained only if the Cauchy data are homogeneous over several horizon lengths. The analogous problem is considered in Nakao et al. [166], [167]; in Shinkai, Maeda [236] also the inclusion of colliding plane gravitational waves is considered, they give a numerical support of the no hair conjecture by concentrating on the dynamics of gravitational waves. Berkin [24] gets as further result, that for $L = f(R)$, a diagonal Bianchi metric is always possible. Barrow and Sirousse-Zia [16] and Spindel [241] also worked on the diagonalization problem. They apply the diagonalization condition of MacCallum et al. [146], [147].

Kottler 1918 [131] found the spherically symmetric vacuum solution for Einstein's theory with Λ -term

$$ds^2 = A(r)dt^2 - \frac{dr^2}{A(r)} - r^2(d\theta^2 + \sin^2\theta d\phi^2) \quad (9.41)$$

with

$$A(r) = 1 - \frac{2m}{r} - \frac{\Lambda}{3}r^2. \quad (9.42)$$

At the horizon $A(r) = 0$ the Killing vector changes its sign and one gets by interchanging the coordinates t and r the corresponding Kantowski-Sachs model. The cosmic no hair conjecture within Kantowski-Sachs models and $\Lambda > 0$ is discussed by Moniz [159]. To get a theory renormalizable to one-loop order one needs at least curvature squared terms in the Lagrangian, and they lead to a fourth-order theory, Stelle [248]. But there exist several further motivations to consider fourth-order theories; to get an overview for early papers on fourth-order gravity see e.g. [199].

Let us now discuss sixth-order equations. Stelle [248] 1977 considers mainly fourth-order R^2 -models; in the introduction he mentioned that in the next order, terms like

$$R^3 + R_{ij;k}R^{ij;k} \quad (9.43)$$

become admissible, but the pure R^3 -term is not admissible. The paper Buchdahl 1951 [42] deals with Lagrangians of arbitrarily high order. Its results are applied in [218], Quandt et al. [187], and Battaglia Mayer et al. [18] to general Lagrangians $F(R, \square)$. In the paper Vilkovisky [263], the Sacharov-approach was generalized. The original idea of A. Sakharov [195] from 1967, see the commented reprint from 2000, was to define higher order curvature corrections to the Einstein action to get a kind of elasticity of the vacuum. Then the usual breakdown of measurements at the Planck length, such a

short de Broglie wave length corresponds to such a large mass which makes the measuring apparatus to a black hole, is softened.

Vilkovisky [263] discusses the effective gravitational action in the form $Rf(\square)R$, where

$$f(\square) = \int \frac{1}{\square - x} \rho(x) dx. \quad (9.44)$$

Martin and Mazzitelli [153] discuss the non-local Lagrangian $R\frac{1}{\square}R$ as conformal anomaly in two dimensions. Treder 1991 [257] used higher-order Lagrangians, especially R^2 -terms, and he mentioned that for

$$R + R_{,i}R_{,k}g^{ik} \quad (9.45)$$

a sixth-order field equation appears. Remark: This Lagrangian leads to the same field equation as $R - R\square R$. Lu and Wise [144] consider the gravitational Lagrangian as a sequence $S = S_0 + S_1 + S_2 + \dots$ ordered with respect to physical dimension. So, $S_0 = R$ and S_1 sums up the R^2 -terms. Kirsten et al. [127] consider the effective Lagrangian for self-interacting scalar fields; in the renormalized action, the term

$$\frac{\square R}{c + R} \quad (9.46)$$

appears. Wands [266] classifies Lagrangians of the type $F(R, \Phi)\square R$ and mentions that not all of them can be conformally transformed to Einstein's theory. Amendola [2] considers the Lagrangian $\Phi^2\square R$, Gottlöber et al. [97] the Lagrangian $R\square R$, Gottlöber et al. [94] and Schmidt [219] discuss double inflation from Φ and R^2 -terms. Besides $R\square R$, Berkin [24] and Berkin, Maeda [25] consider the de Sitter spacetime as attractor solution for field equations where the variational derivative of the term $C_{ijkl}C^{ijkl}$ is included.

This section is organized as follows: subsection 9.5.2 compares several possible definitions of an asymptotic de Sitter spacetime. Subsection 9.5.3 deals with the Lagrangian and corresponding field equations for higher-order gravity, and 9.5.4 gives the no hair theorem for it. In subsection 9.5.5, we determine under which circumstances the Bianchi models in higher-order gravity can be written in diagonal form without loss of generality, and in subsection 9.5.6 we discuss and summarize the results.

9.5.2 Definition of an asymptotic de Sitter spacetime

In this subsection we want to compare some possible definitions of an asymptotic de Sitter spacetime. To this end let us consider the metric

$$ds^2 = dt^2 - e^{2\alpha(t)} \sum_{i=1}^n d(x^i)^2 \quad (9.47)$$

which is the spatially flat Friedmann model in n spatial dimensions. We consider all values $n \geq 1$, but then concentrate on the usual case $n = 3$. The Hubble parameter is

$$H = \dot{\alpha} \equiv \frac{d\alpha}{dt}. \quad (9.48)$$

We get

$$R_{00} = -n \left(\frac{dH}{dt} + H^2 \right), \quad R = -2n\dot{H} - n(n+1)H^2. \quad (9.49)$$

Then it holds: The following 4 conditions for metric (9.47) are equivalent. 1: The spacetime is flat. 2: It holds $R = R_{00} = 0$. 3: The curvature invariant $R_{ij}R^{ij}$ vanishes. 4: Either $\alpha = \text{const.}$ or

$$n = 1 \quad \text{and} \quad \alpha = \ln |t - t_0| + \text{const.} \quad (9.50)$$

For the last case with $n = 1$ one has to observe that

$$ds^2 = dt^2 - (t - t_0)^2 dx^2 \quad (9.51)$$

represents flat spacetime in polar coordinates. For the proof we use the identity

$$R_{ij}R^{ij} = (R_{00})^2 + \frac{1}{n}(R - R_{00})^2. \quad (9.52)$$

An analogous characterization is valid for the de Sitter spacetime. The following 4 conditions for metric (9.47) are equivalent. 1: It is a non-flat spacetime of constant curvature. 2: $R_{00} = R/(n+1) = \text{const.} \neq 0$. 3: $(n+1)R_{ij}R^{ij} = R^2 = \text{const.} \neq 0$. 4: Either $H = \text{const.} \neq 0$ or

$$n = 1 \text{ and } (ds^2 = dt^2 - \sin^2(\lambda t)dx^2 \text{ or } ds^2 = dt^2 - \sinh^2(\lambda t)dx^2). \quad (9.53)$$

For $n = 1$, the de Sitter spacetime and anti-de Sitter spacetime differ by the factor -1 in front of the metric only. For $n > 1$, only the de Sitter spacetime, having $R < 0$, is covered, because the anti-de Sitter spacetime cannot be represented as spatially flat Friedmann model. Our result shows that within the class of spatially flat Friedmann models, a characterization of the de Sitter spacetime using polynomial curvature invariants only, is possible.

Next, let us look for isometries leaving the form of the metric (9.47) invariant. Besides spatial isometries, the map $\alpha \rightarrow \tilde{\alpha}$ defined by

$$\tilde{\alpha}(t) = c + \alpha(\pm t + t_0), \quad c, t_0 = \text{const.} \quad (9.54)$$

leads to an isometric spacetime. The simplest expressions being invariant by such a transformation are H^2 and $\dot{H}$. We take α as dimensionless, then H is an inverse time and $\dot{H}$ an inverse time squared. Let $H \neq 0$ in the following. The expression

$$\varepsilon := \dot{H}H^{-2} \quad (9.55)$$

is the simplest dimensionless quantity defined for the spatially flat Friedmann models and being invariant with respect to this transformation. Let $n > 1$ in

the following, then it holds: Two metrics of type (9.47) are isometric if and only if the corresponding functions α and $\tilde{\alpha}$ are related by this transformation. It follows: Metric (9.47) with $H \neq 0$ represents the de Sitter spacetime if and only if $\varepsilon \equiv 0$.

All dimensionless invariants containing at most second order derivatives of the metric can be expressed as $f(\varepsilon)$, where f is any given function. But if one has no restriction to the order, one gets a sequence of further invariants

$$\varepsilon_2 = \ddot{H}H^{-3}, \quad \dots, \quad \varepsilon_p = \frac{d^p H}{dt^p} H^{-p-1}. \quad (9.56)$$

Let $H > 0$ in metric (9.47) with $n > 1$. We call it an asymptotic de Sitter spacetime if

$$\lim_{t \rightarrow \infty} \frac{\alpha(t)}{t} = \text{const.} \neq 0 \quad (9.57)$$

or

$$\lim_{t \rightarrow \infty} R^2 = \text{const.} > 0 \quad \text{and} \quad \lim_{t \rightarrow \infty} (n+1)R_{ij}R^{ij} - R^2 = 0 \quad (9.58)$$

or for some natural number p one has

$$\lim_{t \rightarrow \infty} \varepsilon_j = 0 \quad \text{for} \quad 1 \leq j \leq p. \quad (9.59)$$

In general, all these definitions are different. Using the identity

$$R_{ij}R^{ij} = n^2(\dot{H} + H^2)^2 + n(\dot{H} + nH^2)^2 \quad (9.60)$$

we will see that all these definitions lead to the same result if we restrict ourselves to the set of solutions of the higher-order field equations. The problem is that none of the above definitions can be generalized to inhomogeneous models. One should find a polynomial curvature invariant which equals a

positive constant if and only if the spacetime is locally the de Sitter spacetime. To our knowledge, such an invariant cannot be found in the literature, but also the non-existence of such an invariant has not been proven up to now. For the anti-de Sitter spacetime, a partial result is given in section 4.3.

This situation is quite different for the positive definite case: For signature $(++++)$ and $S_{ij} = R_{ij} - \frac{R}{4}g_{ij}$ it holds:

$$I \equiv (R - R_0)^2 + C_{ijkl}C^{ijkl} + S_{ij}S^{ij} = 0 \quad (9.61)$$

if and only if the V_4 is a space of constant curvature R_0 . So $I \rightarrow 0$ is a suitable definition of an asymptotic space of constant curvature. One possibility exists, however, for the Lorentz signature case, if one allows additional structure as follows: An ideal fluid has an energy-momentum tensor

$$T_{ij} = (\rho + p)u_i u_j - p g_{ij} \quad (9.62)$$

where u_i is a continuous vector field with $u_i u^i \equiv 1$. For matter with equation of state $\rho = -p$, the equation $T^{ij}_{;j} \equiv 0$ implies $p = \text{const.}$, and so every solution of Einstein's theory with such matter is isometric to a vacuum solution of Einstein's theory with a cosmological term. The inverse statement, however, is valid only locally:

Given a vacuum solution of Einstein's theory with a Λ -term, one has to find continuous timelike unit vector fields which need not to exist from topological reasons. And if they exist, they are not at all unique. So, it becomes possible to define an invariant J which vanishes if and only if the spacetime is de Sitter by transvecting the curvature tensor with $u^i u^j$ and/or g^{ij} and suitable linear and quadratic combinations of such terms. Then time t becomes defined by the streamlines of the vector u^i . If one defines the

asymptotic de Sitter spacetime by $J \rightarrow 0$ as $t \rightarrow \infty$, then it turns out, that this definition is not independent of the vector field u^i .

9.5.3 Lagrangian $F(R, \square R, \square^2 R, \dots, \square^k R)$

Let us consider the Lagrangian density L given by

$$L = F(R, \square R, \square^2 R, \dots, \square^k R) \sqrt{-g} \quad (9.63)$$

where R is the curvature scalar, $\square$ the D'Alembertian and g_{ij} the metric of a Pseudo-Riemannian V_D of dimension $D \geq 2$ and arbitrary signature; $g = -|\det g_{ij}|$. The main application will be $D = 4$ and metric signature $(+ - - -)$. F is supposed to be a sufficiently smooth function of its arguments, preferably a polynomial. Buchdahl [42] already dealt with such kind of Lagrangians in 1951, but then it became quiet of them for decades.

The variational derivative of L with respect to the metric yields the tensor

$$P^{ij} = -\frac{1}{\sqrt{-g}} \frac{\delta L}{\delta g_{ij}} \quad (9.64)$$

The components of this tensor read

$$P_{ij} = GR_{ij} - \frac{1}{2}Fg_{ij} - G_{;ij} + g_{ij}\square G + X_{ij} \quad (9.65)$$

where the semicolon denotes the covariant derivative, R_{ij} the Ricci tensor, and

$$X_{ij} = \sum_{A=1}^k \frac{1}{2}g_{ij}[F_A(\square^{A-1}R)^{;m}]_{;m} - F_{A(;i}[\square^{A-1}R]_{;j)} \quad (9.66)$$

having the round symmetrization brackets in its last term. For $k = 0$, i.e. $F = F(R)$, a case considered in subsection 9.5.4, the tensor X_{ij} identically vanishes. It remains to define the expressions F_A , $A = 0, \dots, k$.

$$F_k = \frac{\partial F}{\partial \square^k R} \quad (9.67)$$

and for $A = k - 1, \dots, 0$

$$F_A = \square F_{A+1} + \frac{\partial F}{\partial \square^A R} \quad (9.68)$$

and finally $G = F_0$. The brackets are essential, for any scalar Φ it holds

$$\square(\Phi_{;i}) - (\square\Phi)_{;i} = R_i^j \Phi_{;j} \quad (9.69)$$

Inserting $\Phi = \square^m R$ into this equation, one gets identities to be applied in the sequel without further notice. The covariant form of energy-momentum conservation reads

$$P_{j;i}^i \equiv 0 \quad (9.70)$$

and P_{ij} identically vanishes if and only if F is a divergence, i.e., locally there can be found a vector v^i such that $F = v^i_{;i}$ holds. Remark: Even for compact manifolds without boundary the restriction “locally” is unavoidable, for example, let $D = 2$ and V_2 be the Riemannian two-sphere S^2 with arbitrary positive definite metric. R is a divergence, but there do not exist continuous vector fields v^i fulfilling $R = v^i_{;i}$ on the whole S^2 . Example: for $m, n \geq 0$ it holds

$$\square^m R \square^n R - R \square^{m+n} R = \text{divergence} . \quad (9.71)$$

So, the terms $\square^m R \square^n R$ with naturals m and n can be restricted to the case $m = 0$ without loss of generality.

9.5.4 No hair theorems for higher-order gravity

For $n > 1$, the $n + 1$ -dimensional de Sitter spacetime is an attractor solution for the field equation derived from the Lagrangian

$$R^{(n+1)/2} . \quad (9.72)$$

It is not an attractor solution for the Lagrangian $R\Box^k R$ and $k > 0$. There exist combinations of coefficients c_i , such that the de Sitter spacetime is an attractor solution for the field equation derived from the more general Lagrangian

$$L = c_0 R^{(n+1)/2} + \sum_{k=0}^m c_k R \Box^k R. \quad (9.73)$$

Idea of Proof: Concerning fourth-order gravity this method was previously applied e.g. by Barrow 1986 [12]. The de Sitter spacetime is an exact solution for the field equation, if and only if $2RG = (n+1)F$. If we make the ansatz

$$\dot{\alpha}(t) = 1 + \beta(t) \quad (9.74)$$

we get the linearized field equation $0 = \ddot{\beta} + n\dot{\beta}$ for the Lagrangian $R^{\frac{n+1}{2}}$. For the Lagrangian $L = R\Box^k R$ we get the linearized field equation $\Box^k R = (\Box^k R)_{,0}$. For the characteristic polynomial we get a recursive formula such that the next order is received from the previous one by multiplying with $\cdot x \cdot (x+n)$. We get the roots $x_1 = -n-1$, $x_2 = -n$ (k -fold), $x_3 = 0$ (k -fold) and $x_4 = +1$. Because of the last root the de Sitter spacetime is not an attractor solution. For the Lagrangian (9.73) we get the characteristic polynomial

$$P(x) = x(x+n) \left[c_0 + \sum_{k=1}^m c_k x^{k-1} (x+n)^{k-1} (x-1)(x+n+1) \right] \quad (9.75)$$

for the linearized field equation. The transformation $z = x^2 + nx + \frac{n^2}{4}$ gives a polynomial $Q(z)$ which can be solved explicitly. So one can find those combinations of the coefficients such that the de Sitter spacetime is an attractor solution.

It turned out that all the variants of the definition of an asymptotic de Sitter solution given in subsection 9.5.2 lead to the same class of solutions,

i.e., the validity of the theorems written below does not depend on which of the variants of definition of an asymptotic de Sitter spacetime listed is applied. For the 6th-order case we can summarize as follows: Let

$$L = R^2 + c_1 R \square R \quad (9.76)$$

and

$$L_E = R - \frac{l^2}{6} L \quad (9.77)$$

with length $l > 0$. Then the following statements are equivalent.

1. The Newtonian limit of L_E is well-behaved, and the potential ϕ consists of terms $e^{-\alpha r}/r$ with $\alpha \geq 0$ only.
2. The de Sitter spacetime with $H = 1/l$ is an attractor solution for L in the set of spatially flat Friedmann models, and this can already be seen from the linearized field equation.
3. The coefficient $c_1 \geq 0$, and the graceful exit problem is solved for the quasi de Sitter phase $H \leq 1/l$ of L_E .
4. $l^2 = l_0^2 + l_1^2$ such that $l^2 c_1 = l_0^2 l_1^2$ has a solution with $0 \leq l_0 < l_1$.
5. $0 \leq c_1 < l^2/4$.

A formulation which includes also the marginally well-behaved cases reads as follows: Let L and L_E as in the previous result, then the following statements are equivalent:

1. The Newtonian limit of L_E is well-behaved, for the potential ϕ we allow $1/r$ and terms like

$$\frac{P(r)}{r} e^{-\alpha r} \quad \text{with} \quad \alpha > 0 \quad (9.78)$$

and a polynomial P .

2. The de Sitter spacetime with $H = 1/l$ cannot be ruled out to be an attractor solution for L in the set of spatially flat Friedmann models if one considers the linearized field equation only.

3. L_E is tachyonic-free.

4. $l^2 = l_0^2 + l_1^2$ with $l^2 c_1 = l_0^2 l_1^2$ has a solution with $0 \leq l_0 \leq l_1$.

5. For the coefficients we have $0 \leq c_1 \leq \frac{l^2}{4}$.

Of course, it would be interesting what happens in the region where the linearized equation does not suffice to decide.

9.5.5 Diagonalizability of Bianchi models

A Bianchi model can always be written as

$$ds^2 = dt^2 - g_{\alpha\beta}(t)\sigma^\alpha\sigma^\beta \quad (9.79)$$

where $g_{\alpha\beta}$ is positive definite and σ^α are the characterizing one-forms. It holds

$$d\sigma^\gamma = -\frac{1}{2}C_{\alpha\beta}^\gamma\sigma^\alpha \wedge \sigma^\beta \quad (9.80)$$

with structure constants $C_{\alpha\beta}^\gamma$ of the corresponding Bianchi type. It belongs to class A if $C_{\alpha\beta}^\beta = 0$. The Abelian group, Bianchi type I, and the rotation group, Bianchi type IX, both belong to class A. In most cases, the $g_{\alpha\beta}$ are written in diagonal form; it is a non-trivial problem to decide under which circumstances this can be done without loss of generality. For Einstein's theory, this problem is solved. One of its results reads: If a Bianchi model of class A, except Bianchi types I and II, has a diagonal energy-momentum tensor, then the metric $g_{\alpha\beta}(t)$ can be chosen in diagonal form. Here, the

energy-momentum tensor is called diagonal, if it is diagonal in the basis $(dt, \sigma^1, \sigma^2, \sigma^3)$.

This result rests of course on Einstein's theory and cannot be directly applied to higher-order gravity. For fourth-order gravity following from a Lagrangian $L = f(R)$ considered in an interval of R -values where

$$\frac{df}{dR} \cdot \frac{d^2f}{dR^2} \neq 0 \quad (9.81)$$

one can do the following: The application of the conformal equivalence theorem, cf. e.g. [29] or [218], is possible, the conformal factor depends on t only, so the diagonal form of metric does not change. The conformal picture gives Einstein's theory with a minimally coupled scalar field as source; the energy-momentum tensor is automatically diagonal. So, in this class of fourth-order theories of gravity, we can apply the above cited theorem of MacCallum et al. [146], [147], where the initial-value formulation of General Relativity is applied. As example we formulate: All solutions of Bianchi type IX of fourth-order gravity following from $L = R^2$ considered in a region where $R \neq 0$ can be written in diagonal form. Consequently, the ansatz used by Barrow and Sirousse-Zia [16] for this problem is already the most general one, cf. Spindel [241].

For fourth-order gravity of a more complicated structure, however, things are more involved; example: Let

$$L = R + aR^2 + bC_{ijkl}C^{ijkl} \quad (9.82)$$

with $ab \neq 0$. Then there exist Bianchi type IX models which cannot be written in diagonal form. This is a non-trivial statement because its proof needs a careful analysis of the correspondingly allowed initial values in the

Cauchy problem. To understand the difference between the cases $b = 0$ and $b \neq 0$ it proves useful to perform the analysis independently of the above cited papers [46]. For simplicity, we restrict to Bianchi type I. Then the internal metric of the hypersurface $[t = 0]$ is flat and we can choose as initial value $g_{\alpha\beta}(0) = \delta_{\alpha\beta}$. Spatial rotations do not change this equation, and we can take advantage of them to diagonalize the second fundamental form $\frac{d}{dt}g_{\alpha\beta}(0)$.

First case: $b = 0$. As additional initial conditions one has only $R(0)$ and $\frac{d}{dt}R(0)$. The field equation ensures $g_{\alpha\beta}(t)$ to remain diagonal for all times.

Second case: $b \neq 0$. Then one has further initial data

$$\frac{d^2}{dt^2}g_{\alpha\beta}(0). \quad (9.83)$$

In the generic case, they cannot be brought to diagonal form simultaneously with $\frac{d}{dt}g_{\alpha\beta}(0)$. This excludes a diagonal form of the whole solution. To complete the proof, one has of course to check that these initial data are not in contradiction to the constraint equations. This case has the following relation to the Bicknell theorem: Just for this case $b \neq 0$, the conformal relation to Einstein's theory breaks down, and if one tries to re-interpret the variational derivative of $C_{ijkl}C^{ijkl}$ as energy-momentum tensor then it turns out to be non-diagonal generically, and the theorem cannot be applied.

For higher-order gravity, the situation becomes even more involved. For a special class of theories, however, the diagonalizability condition is exactly the same as in Einstein's theory: If

$$L = R + \sum_{k=0}^m a_k R \square^k R, \quad a_m \neq 0, \quad (9.84)$$

then in a region where $2L \neq R$ the Cauchy data are the data of General Relativity, $R(0)$, and the first $2m + 1$ temporal derivatives of R at $t = 0$. All

terms with the higher derivatives behave as an energy-momentum tensor in diagonal form, and so the classical theorem applies. Let us comment on the restriction $2L \neq R$ supposed above: $F_0 = G = 0$ represents a singular point of the differential equation; and for the Lagrangian given here $G = 2L/R - 1$. For fourth-order gravity defined by a non-linear Lagrangian $L(R)$ one has

$$G = \frac{dL}{dR},$$

and $G = 0$ defines the critical value of the curvature scalar.

9.5.6 Discussion of no hair theorems

In subsection 9.5.4 we have shown: The results of the Starobinsky model are structurally stable with respect to the addition of a sixth-order term $\sim R\Box R$, if the coefficients fulfil certain inequalities. For the eighth-order case we got: For

$$L = R^2 + c_1 R\Box R + c_2 R\Box\Box R, \quad c_2 \neq 0 \quad (9.85)$$

and the case $n > 1$ the de Sitter spacetime with $H = 1$ is an attractor solution in the set of spatially flat Friedmann models if and only if the following inequalities are fulfilled:

$$0 < c_1 < \frac{1}{n+1}, \quad 0 < c_2 < \frac{1}{(n+1)^2} \quad (9.86)$$

and

$$c_1 > -(n^2 + n + 1)c_2 + \sqrt{(n^4 + 4n^3 + 4n^2)c_2^2 + 4c_2} \quad (9.87)$$

These inequalities define an open region in the $c_1 - c_2$ -plane whose boundary contains the origin; and for the other boundary points the linearized equation does not suffice to decide the attractor property.

This situation shall be called “semi-attractor” for simplicity. In a general context this notion is used to describe a situation where all Lyapunov coefficients have non-positive real parts and at least one of them is purely imaginary.

In contrary to the 6th-order case, here we do not have a one-to-one correspondence, but a non-void open intersection with that parameter set having the Newtonian limit for L_E well-behaved.

To find out, whether another de Sitter spacetime with an arbitrary Hubble parameter $H > 0$ is an attractor solution for the eighth-order field equation following from the above Lagrangian, one should remember that H has the physical dimension of an inverted time, c_1 is a time squared, c_2 is a time to power 4. So, we have to replace c_1 by $c_1 H^2$ and c_2 by $c_2 H^4$ in the above dimensionless inequalities to get the correct conditions. Example:

$$0 < c_1 H^2 < \frac{1}{4}.$$

Let us summarize: Here for a theory of gravity of order higher than four the Newtonian limit and the attractor property of the de Sitter spacetime are systematically compared. It should be noted that the details of the theory sensibly depend on the numerical values of the corresponding coefficients. So, no general overall result about this class of theories is ever to be expected.

The facts, contradicting each other from the first glance, field equations of higher than second order follow from quantum gravity considerations Mirzabekian et al. [156] on the one hand but such equations are known to be unstable in general and are therefore unphysical on the other hand, have now

considered in more details: It depends on the special circumstances whether such theories are unstable.

We have found out that for the class of theories considered here, one of the typical indicators of instability - cosmological runaway-solutions - need not to exist, even for an arbitrarily high order of the field equation. It is an additional satisfactory result that this takes place in the same range of parameters where the Newtonian limit is well behaved.

10 Solutions of the Bach-Einstein equation

For field equations of 4th order, following from a Lagrangian “Ricci scalar plus Weyl scalar”, it is shown using methods of non-standard analysis that in a neighbourhood of Minkowski spacetime there do not exist regular static spherically symmetric solutions. With that, besides the known local expansions about $r = 0$ and $r = \infty$ respectively, a global statement on the existence of such solutions is given, see [205]. Then, this result will be discussed in connection with Einstein’s particle programme. Finally, in section 10.6 it will be shown, that cosmological Bianchi type I solutions of the Bach equation exist which fail to be conformally related to an Einstein space.

10.1 Introduction to the Bach equation

General Relativity Theory starts from the Einstein-Hilbert Lagrangian

$$L_{\text{EH}} = R/2\kappa \quad (10.1)$$

with the Ricci scalar R which leads to the Einstein vacuum field equation $R_i^k = 0$ being of second order in the metrical tensor g_{ik} . Its validity is proven with high accuracy in spacetime regions where the curvature is small only. Therefore, the additional presence of a term being quadratic in the curvature is not excluded by the standard weak field experiments.

Lagrangians with squared curvature have already been discussed by Weyl in 1919 [268], by Bach in 1921 [8] and Einstein in 1921. They were guided by ideas about conformal invariance, and Einstein [66] proposed to look seriously to such alternatives. In Bach [8], the Lagrangian L_{W} with the the Weyl scalar C ,

$$L_{\text{W}} = C/2\kappa, \quad \text{with} \quad C = \frac{1}{2} \cdot C_{ijkl} C^{ijkl} \quad (10.2)$$

was of special interest. Then it became quiet of them for a couple of decades because of the brilliant results of General Relativity but not at least because of mathematical difficulties.

Later, new interest in such equations arose with arguments coming from quantum gravity, cf. e.g. [257], Borzeszkowski, Treder, Yourgrau [37], Stelle [249], Fiedler, Schimming [78]. In the large set of possible quadratic modifications a linear combination

$$L = L_{\text{EH}} + l^2 L_{\text{W}} \quad (10.3)$$

enjoyed a special interest, cf. e.g. Treder 1977 [257]. The coupling constant l has to be a length for dimensional reasons and it has to be a small one to avoid conflicts with observations: One often takes it to be the Heisenberg length, which is the Compton wave length of the proton, $= 1.3 \cdot 10^{-18}$ cm or the Planck length $= 1.6 \cdot 10^{-33}$ cm.

In this chapter we consider the Lagrangian L eq. (10.3) in connection with Einstein's particle programme, see Einstein, Pauli 1943 [69]: One asks for spherically symmetric singularity-free asymptotically flat solutions of the vacuum field equations which shall be interpreted as particles, but this cannot be fulfilled within General Relativity itself; it is still a hope, cf. [35], to realize it in such 4th order field equations.

Two partial answers have already been given: Stelle [249] has shown the gravitational potential of the linearized equations to be

$$\Phi(r) = -m/r + \exp(-r/l)c/r \quad \text{for} \quad r \ll l; \quad (10.4)$$

he also gave an expansion series about $r = 0$, and Fiedler, Schimming [78] proved its convergence and smoothness in a certain neighbourhood of $r = 0$, but said nothing about the convergence radius of this expansion.

Now we want to join these two local expansions. To this end we write down the field equations in different but equivalent versions section 10.2, calculate some linearizations section 10.3 and prove the statement of the summary in section 10.4 which will be followed by a discussion of Einstein's particle programme in section 10.5.

10.2 Notations and field equations

We start from a spacetime metric with signature $(+---)$, the Riemann and Ricci tensor being defined by $R^i_{jkl} = \Gamma^i_{jl,k} - \dots$, $R_{ik} = R^j_{ijk}$ respectively. The Weyl tensor C^i_{jkl} is the traceless part of the Riemann tensor and the Weyl scalar is given by $2C = C_{ijkl}C^{ijkl}$. Light velocity is taken to be 1 and $G = \kappa/8\pi$ is Newton's constant; $g = \det g_{ij}$.

Then we consider the Lagrangian

$$\mathcal{L} = \sqrt{-g}L = \sqrt{-g}(R + l^2C)/2\kappa + \mathcal{L}_{\text{mat}}, \quad (10.5)$$

where $\mathcal{L}_{\text{mat}}$ is the matter Lagrangian. For writing down the corresponding field equation it is convenient to introduce the Bach tensor B_{ij} , cf. Bach [8] and Wunsch [271], beforehand.

$$\frac{1}{2}B_{ij} = \frac{\kappa\delta(\sqrt{-g}L_W)}{\sqrt{-g}\delta g^{ij}} = C^a_{ij}{}^b{}_{;ba} + \frac{1}{2}C^a_{ij}{}^b{}^b R_{ba}. \quad (10.6)$$

It holds $B^i_i = 0$, $B^j_{i;j} = 0$, $B_{ij} = B_{ji}$, and B_{ij} is conformally invariant of weight -1. Variation of eq. (10.5) with respect to g_{ij} leads to the field equation

$$R_{ij} - \frac{1}{2}g_{ij}R + l^2B_{ij} = \kappa T_{ij}. \quad (10.7)$$

Now let us consider the vacuum case $T_{ij} = 0$. The trace of eq. (10.7) then simply reads $R = 0$, and eq. (10.7) becomes equivalent to the simpler one

$$R_{ij} + l^2 B_{ij} = 0. \quad (10.8)$$

Writing $k = l^{-2}$ and $\square \cdot = g^{ij}(\cdot)_{;ij}$ one gets also the equivalent system

$$R = 0; \quad kR_{ij} + \square R_{ij} = 2R_{iabj}R^{ab} + \frac{1}{2}g_{ij}R_{ab}R^{ab}. \quad (10.9)$$

The static spherically symmetric line element in Schwarzschild coordinates reads

$$ds^2 = (1 + \beta)e^{-2\lambda}dt^2 - (1 + \beta)^{-1}dr^2 - r^2d\Omega^2, \quad (10.10)$$

where β and λ depend on r only. The dot means differentiation with respect to r , and defining

$$\alpha = 2\beta - 2r\dot{\beta} + 2r\dot{\lambda}(1 + \beta) + 2r^2(\dot{\lambda}^2 - \ddot{\lambda})(1 + \beta) + r^2(\ddot{\beta} - 3\dot{\lambda}\dot{\beta}), \quad (10.11)$$

$$\zeta = r\dot{\alpha}, \quad \eta_1 = \alpha r^{-3}, \quad \eta_2 = 3\beta r^{-1}, \quad \eta_3 = \zeta r^{-3}, \quad (10.12)$$

the field equations (10.9) are just eqs. (2.15.a-c) and (2.16) of the paper [78]. To avoid the products $r\dot{\eta}_i$ we define a new independent variable x by $r = le^x$. Then $r\dot{\eta}_i = \eta'_i = d\eta_i/dx$ and we obtain

$$0 = \eta'_1 + 3\eta_1 - \eta_3, \quad (10.13)$$

$$0 = (k + r\eta_3/6)\eta'_2 + \eta_1 + \eta_3 + r^2k\eta_1 + r^3\eta_1\eta_3/6 + r\eta_2\eta_3/2 + r^3\eta_1^2/4, \quad (10.14)$$

$$0 = (1 + r\eta_2/3)\eta'_3 - 2\eta_1 + 2k\eta_2 + 2\eta_3 - r^2k\eta_1 - r^3\eta_1^2/2 - r^3\eta_1\eta_3/6 + 2r\eta_2\eta_3/3. \quad (10.15)$$

Conversely, if the system eqs. (10.13), (10.14), (10.15) is solved, the metric can be obtained by

$$\beta = r\eta_2/3, \quad \dot{\lambda} = (2r\dot{\eta}_2 + 2\eta_2 + r^2\eta_1)/(6 + 2r\eta_2), \quad \lambda(0) = 0. \quad (10.16)$$

10.3 Linearizations

Now we make some approximations to obtain the rough behaviour of the solutions. First, for $r \rightarrow \infty$, Stelle [249] has shown the following: Linearization about Minkowski spacetime leads, using our notations eqs. (10.4) and (10.10), to $\lambda = 0$ and $\beta = 2\Phi$, where powers of Φ are neglected and the term $\exp(r/l)/r$ in Φ must be suppressed because of asymptotical flatness.

That means, loosely speaking, the right hand side of eq. (10.9) will be neglected. Connected with this one may doubt the relevance of the term $cr^{-1}e^{-r/l}$ in eq. (10.4) for $m \neq 0$ and $r \rightarrow \infty$, because it is small compared with the neglected terms. Further, a finite $\Phi(0)$ requires $c = m$ in eq. (10.4). Then Φ is just the Bopp-Podolsky potential stemming from 4th order electrodynamics, and it were Pechlaner and Sexl [180] who proposed this form as representing a gravitational potential. Now we have

$$\Phi(r) = m(\exp(-r/l) - 1)/r = -m/l + rm/(2l^2) + \dots, \quad (10.17)$$

but also this finite potential gives rise to a singularity: For one of the curvature invariants we get $R_{ij}R^{ij} \approx m^2/r^4$ as $r \rightarrow 0$, that means, the linearization makes no sense for this region. From this one can already expect that in a neighbourhood of Minkowski spacetime no regular solutions exist.

Second, for $r \rightarrow 0$, in [78] it is shown that there exists a one-parameter family of solutions being singularity-free and analytical in a neighbourhood of $r = 0$. The parameter will be called ε and can be defined as follows: neglecting the terms with r in the system eqs. (10.13), (10.14), (10.15) one obtains a linear system with constant coefficients possessing just a one-parameter family of solutions being regular at $r = 0$; it reads

$$\eta_1 = \varepsilon k^2 r, \quad \eta_2 = -5\varepsilon k r, \quad \eta_3 = 4\varepsilon k^2 r. \quad (10.18)$$

The factors k^n are chosen such that ε becomes dimensionless. Now one can take eq. (10.18) as the first term of a power series $\eta_i = \sum_n a_i^{(n)} r^n$, and inserting this ansatz into that system, one iteratively obtains the coefficients $a_i^{(n)}$. For n even, $a_i^{(n)}$ vanishes: The r^3 -terms are $k^3(\varepsilon/14 + 10\varepsilon^2/21)$, $-k^2(\varepsilon/2 - 10\varepsilon^2/3)$ and $k^3(3\varepsilon/7 + 20\varepsilon^2/7)$ respectively. Furthermore, the r^{2n-1} -term is always a suitable power of k times a polynomial in ε of the order $\leq n$.

Up to the r^2 -terms the corresponding metric (10.10) reads

$$ds^2 = (1 + 5k\varepsilon r^2/3)dt^2 - (1 + 5k\varepsilon r^2/3)dr^2 - r^2 d\Omega^2. \quad (10.19)$$

Third, we look for a linearization which holds uniformly for $0 \leq r < \infty$. A glance at eqs. (10.16) and (10.10) shows that $\eta_i = 0$ gives Minkowski spacetime, and therefore we neglect terms containing products of η_i in the system. In other words, instead of the linearization used before we additionally retain the term $r^2\eta_1$. Then we proceed as follows: from eqs. (10.13) and (10.15) we obtain

$$\eta_3 = \eta_1' + 3\eta_1 \quad \text{and} \quad (10.20)$$

$$\eta_2 = \frac{-\eta_1' - 5\eta_1 + (e^{2x} - 4)\eta_1 + l^3 e^{3x} \eta_1 (\eta_1 + \eta_1'/6)}{2k + l e^x (\eta_1' + 5\eta_1 + 6\eta_1)/3}. \quad (10.21)$$

Inserting eqs. (10.20), (10.21) into eq. (10.14) one obtains a third order equation for η_1 only whose linearization reads

$$0 = \eta_1''' + 5\eta_1'' + \eta_1'(2 - e^{2x}) - \eta_1(8 + 4e^{2x}). \quad (10.22)$$

In Schmidt and Müller [231] the same vacuum field equations were discussed for axially symmetric Bianchi type I models. They possess a four-parameter group of isometries, too, and the essential field equation is also a third order

equation for one function. The difference is that in the present case spherical symmetry implies an explicit coordinate dependence.

The solution of eq. (10.22) which is bounded for $x \rightarrow -\infty$ reads

$$\eta_1 = \gamma \cdot \left[(3e^{-4x} + e^{-2x}) \sinh e^x - 3e^{-3x} \cosh e^x \right]. \quad (10.23)$$

A second solution is $\eta_1 = -12mr^{-4}$ leading to the Schwarzschild solution, this solution solves both the full and the linearized equations in accordance with the fact that it makes zero both sides of eq. (10.9), and the third one can be obtained from them up to quadrature by usual methods. A comparison of eq. (10.23) with eq. (10.18) gives $\gamma = 15\epsilon l^{-3}$.

Now we insert this η_1 into eqs. (10.21), (10.16) and neglect again powers of η_1 ; then the metric reads

$$\begin{aligned} \beta &= 5\epsilon \left[lr^{-1} \sinh(rl^{-1}) - \cosh(rl^{-1}) \right], \\ \lambda &= \frac{5\epsilon}{2l} \cdot \int_0^r \left[(l^2 z^{-2} - 1) \sinh(zl^{-1}) - lz^{-1} \cosh(zl^{-1}) \right] dz. \end{aligned} \quad (10.24)$$

This linearization has, in contrary to eqs. (10.4) and (10.18), the following property: to each $r_0 > 0$ and $\Delta > 0$ there exists an $\delta > 0$ such that for $-\delta < \epsilon < \delta$ the relative error of the linearized solution (10.24) does not exceed Δ uniformly on the interval $0 \leq r \leq r_0$.

10.4 Non-standard analysis

In this section we will prove that in a neighbourhood of Minkowski spacetime there do not exist any solutions. “Neighbourhood of Minkowski spacetime” is in general a concept requiring additional explanations because of the large number of different topologies discussed in literature. To make the above

statement sufficiently strong we apply a quite weak topology here: given a $\delta > 0$ then all spacetimes being diffeomorphic to Minkowski spacetime and fulfilling $|R_{ij}R^{ij}| < \delta k^2$ form a neighbourhood about Minkowski spacetime. In eq. (10.19) we have at $r = 0$, independently of higher order terms in r ,

$$R_{ij}R^{ij} = 100k^2\varepsilon^2/3 \quad \text{and} \quad R_{00} = 5k\varepsilon. \quad (10.25)$$

Therefore it holds: the one-parameter family of solutions being regular at $r = 0$ is invariantly characterized by the real parameter ε , and a necessary condition for it to be within a neighbourhood of Minkowski spacetime is that ε lies in a neighbourhood of 0.

Now we suppose ε to be an infinitesimal number, i.e., a positive number which is smaller than any positive real number. The mathematical theory dealing with such infinitesimals is called non-standard analysis, cf. Robinson [191]. The clue is that one can handle non-standard numbers like real numbers. Further we need the so-called Permanence principle, which is also called the Robinson lemma: Let $A(\varepsilon)$ be an internal statement holding for all infinitesimals ε . Then there exists a positive standard real $\delta > 0$ such that $A(\varepsilon)$ holds for all ε with $0 < \varepsilon < \delta$. The presumption “internal” says, roughly speaking, that in the formulation of $A(\cdot)$ the words “standard” and “infinitesimal” do not appear.

This permanence principle shall be applied as follows: $A(\varepsilon)$ is the statement: “Take eq. (10.18) as initial condition for eqs. (10.13), (10.14), (10.15) and calculate the corresponding metric (10.10), (10.16). Then there exists an $r_0 \leq l$ such that $R_{ij}R^{ij} \geq k^2$ at $r = r_0$.” Remark: At this point it is not essential whether r_0 is a standard or an infinitely large non-standard number.

Proof of $A(\varepsilon)$ for infinitesimals ε with $\varepsilon \neq 0$: the difference of eq. (10.24) in relation to the exact solution is of the order ε^2 , i.e. the relative error is infinitesimally small. For increasing r , $|\beta|$ becomes arbitrary large, i.e. $(1 + \beta)^{-1}$ becomes small and $R_{ij}R^{ij}$ increases to arbitrarily large values, cf. eq. (10.24). Now take r_0 such that with metric (10.10), (10.24) $R_{ij}R^{ij} \geq 2k^2$ holds. Then, for the exact solution, $R_{ij}R^{ij} \geq k^2$ holds at $r = r_0$, because their difference was shown to be infinitesimally small.

Now the permanence principle tells us that there exists a positive standard real $\delta > 0$ such that for all ε with $0 < |\varepsilon| < \delta$ the corresponding exact solution has a point r_0 such that at $r = r_0$ the inequality $R_{ij}R^{ij} \geq k^2$ holds. But “ $R_{ij}R^{ij} < k^2$ ” is another necessary condition for a solution to lie within a neighbourhood of Minkowski spacetime. Remark: Supposed this r_0 is an infinitely large non-standard number, then by continuity arguments also a standard finite number with the same property exists.

The statement is proved, but we have learned nothing about the actual value of the number δ . Here, numerical calculations may help. They were performed as follows: the power series for the functions η_i were calculated up to the r^6 -term, then these functions taken at $x = -4$ i.e. $r = 0.018 l$ were used as initial conditions for a Runge–Kutta integration. We got the following result: firstly, for $\varepsilon = \pm 10^{-5}$ and $r \leq 10 l$, the relative difference between the linearized solution eq. (10.23) and the numerical one is less than 2 per thousand. Secondly, for $0 < |\varepsilon| \leq 1$ the behaviour “ $\beta \rightarrow -\infty \cdot \text{sgn} \varepsilon$ ” is confirmed. That means, the statement made above keeps valid at least for $\delta = 1$. Remark: For large values ε the power series for the η_i converge very slowly, and therefore other methods would be necessary to decide about asymptotical flatness.

10.5 Discussion – Einstein’s particle programme

Fourth-order gravitational field equations could be taken as a field theoretical model of ordinary matter, the energy-momentum tensor of which is defined by

$$\kappa T_{ij}^* = R_{ij} - \frac{1}{2}g_{ij}R. \quad (10.26)$$

For our case one obtains at $r = 0$ that T_{ij}^* represents an ideal fluid with the equation of state $p^* = \mu^*/3$ and, cf. eq. (10.25),

$$\mu^*(0) = 5k\varepsilon\kappa^{-1} = 5\varepsilon/8\pi Gl^2. \quad (10.27)$$

Inserting $|\varepsilon| \geq 1$ and $l \leq 1.3 \cdot 10^{-13}$ cm into eq. (10.27), we obtain

$$|\mu^*(0)| \geq 1.5 \cdot 10^{53} \text{ g cm}^{-3}. \quad (10.28)$$

Therefore it holds: if there exists a non-trivial static spherically symmetric asymptotically flat singularity-free solution of eq. (10.8) at all, then the corresponding particle would be a very massive one: its phenomenological energy density exceeds that of a neutron star by at least 40 orders of magnitude.

The resulting statement can be understood as follows: For a small curvature the 4th order corrections to Einstein’s equations are small, too, and the situation should not be very different from that one we know from Einstein’s theory. Now we want to refer to a problem concerning Schwarzschild coordinates: the transition from a general static spherically symmetric line element to Schwarzschild coordinates is possible only in the case that the function “invariant surface of the sphere $r = \text{const.}$ in dependence on its invariant radius” has not any stationary point. Here two standpoints are possible: either one takes this as a natural condition for a reasonable particle model or one allows coordinate singularities in eq. (10.10) like “ $\beta \geq -1$,

and $\beta = -1$ at single points” which require a special care. For $\beta < -1$ one would obtain a cosmological model of Kantowski-Sachs type. Eqs. (10.5) till (10.16) remain unchanged for this case. The discussion made above is not influenced by this.

The statement on the existence of solutions can be strengthened as follows: Fiedler and Schimming [78] proved that the solutions are analytical in a neighbourhood of $r = 0$. Further, the differential equation is an analytical one and, therefore, in the subspace of singularity-free solutions they remain so in the limit $r \rightarrow \infty$. Then, there exists only a finite or countably infinite set of values ε_n such that the corresponding solution becomes asymptotically flat; the question, whether this set is empty or not, shall be subject of further investigation. Furthermore, the ε_n have no finite accumulation point. That means, there exists at most a discrete spectrum of solutions.

With respect to this fact, we remark the following: as one knows, Einstein’s theory is a covariant one. But besides this symmetry, it is homothetically invariant, too. That means, if ds^2 is changed to $e^{2\chi}ds^2$ with constant χ , then the tensor $R_{ij} - \frac{1}{2}g_{ij}R$ remains unchanged, whereas the scalars R and C will be divided by $e^{2\chi}$ and $e^{4\chi}$ respectively. From this it follows: with one solution of Einstein’s vacuum equation one obtains by homothetic invariance just a one-parameter class of solutions. On the other hand, the sum $R + l^2C$ has not such a symmetry and, therefore, one should not expect that a one-parameter family of solutions globally exists, and this is just in the scope of the particle programme where a definite particle’s mass is wanted.

10.6 Non-trivial solutions of the Bach equation exist

Following [204], we show that solutions of the Bach equation exist which are not conformal Einstein spaces. In connection with fourth-order gravitational field equations, cf. e.g. Weyl 1921 [268], [37] where the breaking of conformal invariance was discussed, the original Bach equation,

$$B_{ij} = 0, \quad (10.29)$$

enjoys current interest, see e.g. [64], [71] and [91]. Eq. (10.29) stems from a Lagrangian

$$L = \frac{1}{2} \sqrt{-g} C_{ijkl} C^{ijkl}, \quad (10.30)$$

and variation gives, cf. Bach 1921 [8],

$$\frac{1}{\sqrt{-g}} \delta L / \delta g^{ij} = B_{ij} = 2 C^a{}_{ij}{}^b{}_{;ba} + C^a{}_{ij}{}^b R_{ba}. \quad (10.31)$$

An Einstein space, defined by

$$R_{ij} = \lambda g_{ij}, \quad (10.32)$$

is always a solution of the Bach equation (10.29). But eq. (10.29) is conformally invariant, and therefore, each metric, which is conformally related to an Einstein space, fulfils eq. (10.29), too. We call such solutions trivial ones.

Now the question arises whether non-trivial solutions of the Bach equation eq. (10.29) do or do not exist. In this section we will give an affirmative answer. As a by-product, some conditions will be given under which only trivial solutions exist. Observe that eq. (10.29) is conformally invariant whereas eq. (10.32) is not. Therefore, a simple counting of degrees of freedom does not suffice.

Because the full set of solutions of eq. (10.29) is not easy to describe, let us consider some homogeneous cosmological models. Of course, we have to consider anisotropic ones, because all Friedmann models are trivial solutions of eq. (10.29). Here, we concentrate on the diagonal Bianchi type I models

$$ds^2 = dt^2 - a_i^2 (dx^i)^2 \quad (10.33)$$

with Hubble parameters $h_i = a_i^{-1} da_i/dt$, $h = \Sigma h_i$ and anisotropy parameters $m_i = h_i - h/3$. The Einstein spaces of this kind are described in [134] eq. (11.52), for $\lambda = 0$ it is just the Kasner metric $a_i = t^{p_i}$, $\Sigma p_i = \Sigma p_i^2 = 1$. All these solutions have the property that the quotient of two anisotropy parameters, m_i/m_j , which equals $(3p_i - 1)/(3p_j - 1)$ for the Kasner metric, is independent of t , and this property is a conformally invariant one. Furthermore, it holds: A solution of eqs. (10.29), (10.33) is a trivial one, if and only if the quotients m_i/m_j are constants.

Restricting now to axially symmetric Bianchi type I models, i.e., metric (10.33) with $h_1 = h_2$, the identity $\Sigma m_i = 0$ implies $m_1/m_2 = 1$, $m_3/m_1 = m_3/m_2 = -2$, i.e., each axially symmetric Bianchi type I solution of eq. (10.29) is conformally related to an Einstein space. Analogously, all static spherically symmetric solutions of eq. (10.29) are trivial ones, cf. [77].

Finally, the existence of a solution of eqs. (10.29), (10.33) with a non-constant m_1/m_2 will be shown. For the sake of simplicity we use the gauge condition $h = 0$, which is possible because of the conformal invariance of eq. (10.29). Then the 00 component and the 11 component of eq. (10.29) are sufficient to determine the unknown functions h_1 and h_2 ; $h_3 = -h_1 - h_2$ follows from the gauge condition. Defining $r = (h_1^2 + h_1 h_2 + h_2^2)^{1/2}$ and

$p = h_1/r$, eq. (10.29) is equivalent to the system

$$3d^2(pr)/dt^2 = 8pr^3 + 4c, \quad c = \text{const.}, \quad p^2 \leq 4/3, \quad (10.34)$$

$$9(dp/dt)^2 r^4 = [2rd^2r/dt^2 - (dr/dt)^2 - 4r^4](4r^2 - 3p^2r^2). \quad (10.35)$$

As one can see, solutions with a non-constant p exist, i.e., m_1/m_2 is not constant for this case.

Result: Each solution of the system eqs. (10.33), (10.34), (10.35) with $dp/dt \neq 0$ represents a non-trivial solution of the Bach equation (10.29).

11 The Newtonian limit of non-linear gravity

In this chapter, the weak-field slow-motion limit of fourth and higher-order gravity will be deduced. Here we follow [208] for the fourth-order case and [187] for the general case. More explicitly: we consider the Newtonian limit of the theory based on the Lagrangian

$$\mathcal{L} = \left(R + \sum_{k=0}^p a_k R \square^k R \right) \sqrt{-g}. \quad (11.1)$$

The gravitational potential of a point mass turns out to be a combination of Newtonian and Yukawa terms. For fourth-order and sixth-order gravity, $p = 0$ and $p = 1$ respectively, the coefficients are calculated explicitly. For general p one gets the potential to be

$$\Phi = m/r \left(1 + \sum_{i=0}^p c_i \exp(-r/l_i) \right) \quad (11.2)$$

with certain coefficients c_i fulfilling the relation

$$\sum_{i=0}^p c_i = 1/3. \quad (11.3)$$

Therefore, the potential is always unbounded near the origin, see also [10].

11.1 The Newtonian limit of 4th-order gravity

Let us consider the gravitational theory defined by the Lagrangian

$$L_g = (8\pi G)^{-1} \left(R/2 + (\alpha R_{ij} R^{ij} + \beta R^2) l^2 \right). \quad (11.4)$$

G is Newton's constant, l a coupling length and α and β numerical parameters. R_{ij} and R are the Ricci tensor and its trace. Introducing the matter Lagrangian L_m and varying $L_g + L_m$ one obtains the field equation

$$E_{ij} + \alpha H_{ij} + \beta G_{ij} = 8\pi G T_{ij}. \quad (11.5)$$

For $\alpha = \beta = 0$ this reduces to General Relativity Theory. The explicit expressions H_{ij} and G_{ij} can be found in sections 8.4 and 10.2, see also Stelle [249].

In a well-defined sense, the weak-field slow-motion limit of Einstein's theory is just Newton's theory, cf. Dautcourt 1964 [60]. In the following, we consider the analogous problem for fourth-order gravity eqs. (11.4), (11.5). In some cases, the Newtonian limit of the theory defined by eq. (11.5) has already been deduced in the literature: For the special case $\alpha = 0$ see Pechlaner and Sexl [180] or Polijevktov-Nikoladze [183]. For the case $\alpha + 2\beta = 0$ see Havas [105] or Jankiewicz [121], and for $\alpha + 3\beta = 0$ see Borzeszkowski, Treder and Yourgrau [37]. Cf. also Anandan [5], where torsion has been taken into account.

The slow-motion limit can be equivalently described as the limit $c \rightarrow \infty$, where c is the velocity of light. In this sense we have to take the limit $G \rightarrow 0$ while $G \cdot c$ and l remain constants. Then the energy-momentum tensor T_{ij} reduces to the rest mass density ρ :

$$T_{ij} = \delta_i^0 \delta_j^0 \rho, \quad (11.6)$$

$x^0 = t$ being the time coordinate. The metric can be written as

$$ds^2 = (1 - 2\phi)dt^2 - (1 + 2\psi)(dx^2 + dy^2 + dz^2). \quad (11.7)$$

Now eqs. (11.6) and (11.7) will be inserted into eq. (11.5). In our approach, products and time derivatives of ϕ and ψ can be neglected, i.e.,

$$R = 4\Delta\psi - 2\Delta\phi, \quad \text{where} \quad \Delta f = f_{,xx} + f_{,yy} + f_{,zz}. \quad (11.8)$$

Further $R_{00} = -\Delta\phi$, $H_{00} = -2\Delta R_{00} - \Delta R$ and $G_{00} = -4\Delta R$, where $l = 1$.

Then it holds: The validity of the 00-component and of the trace of eq. (11.5),

$$R_{00} - R/2 + \alpha H_{00} + \beta G_{00} = 8\pi G\rho \quad (11.9)$$

and

$$-R - 4(\alpha + 3\beta)\Delta R = 8\pi G\rho, \quad (11.10)$$

imply the validity of the full eq. (11.5).

Now, let us discuss eqs. (11.9) and (11.10) in more details: Eq. (11.9) reads

$$-\Delta\phi - R/2 + \alpha(2\Delta\Delta\phi - \Delta R) - 4\beta\Delta R = 8\pi G\rho. \quad (11.11)$$

Subtracting one half of eq. (11.10) yields

$$-\Delta\phi + 2\alpha\Delta\Delta\phi + (\alpha + 2\beta)\Delta R = 4\pi G\rho. \quad (11.12)$$

For $\alpha + 2\beta = 0$ one obtains

$$-(1 - 2\alpha\Delta)\Delta\phi = 4\pi G\rho \quad (11.13)$$

and then $\psi = \phi$ is a solution of eqs. (11.9), (11.10). For all other cases the equations for ϕ and ψ do not decouple immediately, but, to get equations comparable with Poisson's equation we apply Δ to eq. (11.10) and continue as follows.

For $\alpha + 3\beta = 0$ one gets from eq. (11.12)

$$-(1 - 2\alpha\Delta)\Delta\phi = 4\pi G(1 + 2\alpha\Delta/3)\rho. \quad (11.14)$$

The Δ -operator applied to the source term in eq. (11.14) is only due to the application of Δ to the trace, the original equations (11.9), (11.10) contain only ρ itself.

For $\alpha = 0$ one obtains similarly the equation

$$-(1 + 12\beta\Delta)\Delta\phi = 4\pi G(1 + 16\beta\Delta)\rho. \quad (11.15)$$

For all other cases - just the cases not yet covered by the literature - the elimination of ψ from the system (11.9), (11.10) gives rise to a sixth-order equation

$$-(1 + 4(\alpha + 3\beta)\Delta)(1 - 2\alpha\Delta)\Delta\phi = 4\pi G(1 + 2(3\alpha + 8\beta)\Delta)\rho. \quad (11.16)$$

Fourth-order gravity is motivated by quantum-gravity considerations and therefore, its long-range behaviour should be the same as in Newton's theory. Therefore, the signs of the parameters α , β should be chosen to guarantee an exponentially vanishing and not an oscillating behaviour of the fourth-order terms:

$$\alpha \geq 0, \quad \alpha + 3\beta \leq 0. \quad (11.17)$$

On the other hand, comparing parts of eq. (11.16) with the Proca equation it makes sense to define the masses

$$m_2 = (2\alpha)^{-1/2} \quad \text{and} \quad m_0 = (-4(\alpha + 3\beta))^{-1/2}. \quad (11.18)$$

Then eq. (11.17) prevents the masses of the spin 2 and spin 0 gravitons to become imaginary.

Now, inserting a delta source $\rho = m\delta$ into eq. (11.16) one obtains for ϕ the same result as Stelle [249],

$$\phi = mr^{-1}(1 + \exp(-m_0r)/3 - 4\exp(-m_2r)/3). \quad (11.19)$$

To obtain the metric completely one has also to calculate ψ . It reads

$$\psi = mr^{-1}(1 - \exp(-m_0 r)/3 - 2 \exp(-m_2 r)/3). \quad (11.20)$$

For finite values m_0 and m_2 these are both bounded functions, also for $r \rightarrow 0$. In the limits $\alpha \rightarrow 0$, i.e. $m_2 \rightarrow \infty$, and $\alpha + 3\beta \rightarrow 0$, i.e. $m_0 \rightarrow \infty$, the terms with m_0 and m_2 in eqs. (11.19) and (11.20) simply vanish. For these cases ϕ and ψ become unbounded as $r \rightarrow 0$.

Inserting eqs. (11.19) and (11.20) into the metric (11.7), the behaviour of the geodesics shall be studied. First, for an estimation of the sign of the gravitational force we take a test particle at rest and look whether it starts falling towards the centre or not. The result is: for $m_0 \leq 2m_2$, gravitation is always attractive, and for $m_0 > 2m_2$ it is attractive for large but repelling for small distances. The intermediate case $m_0 = 2m_2$, i.e., $3\alpha + 8\beta = 0$, is already known to be a special one from eq. (11.16).

Next, let us study the perihelion advance for distorted circle-like orbits. Besides the general relativistic perihelion advance, which vanishes in the Newtonian limit, we have an additional one of the following behaviour: for $r \rightarrow 0$ and $r \rightarrow \infty$ it vanishes and for $r \approx 1/m_0$ and $r \approx 1/m_2$ it has local maxima, i.e., resonances.

Finally, it should be mentioned that the gravitational field of an extended body can be obtained by integrating eqs. (11.19) and (11.20). For a spherically symmetric body the far field is also of the type

$$mr^{-1}(1 + a \exp(-m_0 r) + b \exp(-m_2 r)), \quad (11.21)$$

and the factors a and b carry information about the mass distribution inside the body.

11.2 Introduction to higher-order gravity

One-loop quantum corrections to the Einstein equation can be described by curvature-squared terms and lead to fourth-order gravitational field equations; their Newtonian limit is described by a potential “Newton + one Yukawa term”, cf. e.g. Stelle [249] and Teyssandier [253]. A Yukawa potential has the form $\exp(-r/l)/r$ and was originally used by Yukawa [274] to describe the meson field.

Higher-loop quantum corrections to the Einstein equation are expected to contain terms of the type $R\Box^k R$ in the Lagrangian, which leads to a gravitational field equation of order $2k + 4$, cf. [97]. Some preliminary results to this type of equations are already due to Buchdahl [42]. For $k = 1$, the cosmological consequences of the corresponding sixth-order field equations are discussed by Berkin and Maeda [25], and by Gottlöber, Müller and Schmidt [94].

In the present chapter we deduce the Newtonian limit following from this higher order field equation. The Newtonian limit of General Relativity Theory is the usual Newtonian theory, cf. e.g. Dautcourt 1964 [60] or Stephani [250]. From the general structure of the linearized higher-order field equation, cf. [218], one can expect that for this higher-order gravity the far field of the point mass in the Newtonian limit is the Newtonian potential plus a sum of different Yukawa terms. And just this form is that one discussed in connection with the fifth force, cf. [85], [242] and [197]. Here we are interested in the details of this connection between higher-order gravity and the lengths and coefficients in the corresponding Yukawa terms.

11.3 Lagrangian and field equation

Let us start with the Lagrangian

$$\mathcal{L} = \left(R + \sum_{k=0}^p a_k R \square^k R \right) \cdot \sqrt{-g}, \quad a_p \neq 0. \quad (11.22)$$

In our considerations we will assume that for the gravitational constant G and for the speed of light c it holds $G = c = 1$. This only means a special choice of units. In eq. (11.22), R denotes the curvature scalar, $\square$ the D'Alembertian, and g the determinant of the metric. Consequently, the coefficient a_k has the dimension “length to the power $2k + 2$ ”.

The starting point for the deduction of the field equation is the principle of minimal action. A necessary condition for it is the stationarity of the action:

$$-\frac{\delta \mathcal{L}}{\delta g_{ij}} = 8\pi T^{ij} \sqrt{-g}, \quad (11.23)$$

where T^{ij} denotes the energy-momentum tensor. The explicit equations for

$$P^{ij} \sqrt{-g} = -\frac{\delta \mathcal{L}}{\delta g_{ij}} \quad (11.24)$$

are given in [218]. Here we only need the linearized field equation. It reads, cf. [97]

$$P^{ij} \equiv R^{ij} - \frac{R}{2} g^{ij} + 2 \sum_{k=0}^p a_k [g^{ij} \square^{k+1} R - \square^k R^{;ij}] = 8\pi T^{ij}, \quad (11.25)$$

and for the trace it holds:

$$g_{ij} \cdot P^{ij} = -\frac{n-1}{2} R + 2n \sum_{k=0}^p a_k [g^{ij} \square^{k+1} R] = 8\pi T. \quad (11.26)$$

n is the number of spatial dimensions; the most important application is of cause $n = 3$. From now on we put $n = 3$.

11.4 The Newtonian limit in higher-order gravity

The Newtonian limit is the weak-field static approximation. So we use the linearized field equation and insert a static metric and an energy-momentum tensor

$$T_{ij} = \delta_i^0 \delta_j^0 \rho, \quad \rho \geq 0 \quad (11.27)$$

into eq. (11.25).

Without proof we mention that the metric can be brought into spatially conformally flat form, and so we may use

$$\begin{aligned} g_{ij} &= \eta_{ij} + f_{ij}, \\ \eta_{ij} &= \text{diag}(1, -1, -1, -1) \quad \text{and} \\ f_{ij} &= \text{diag}(-2\Phi, -2\Psi, -2\Psi, -2\Psi). \end{aligned} \quad (11.28)$$

Then the metric equals

$$ds^2 = (1 - 2\Phi)dt^2 - (1 + 2\Psi)(dx^2 + dy^2 + dz^2), \quad (11.29)$$

where Φ and Ψ depend on x, y and z . Linearization means that the metric g_{ij} has only a small difference to η_{ij} ; quadratic expressions in f_{ij} and its partial derivatives are neglected. We especially consider the case of a point mass. In this case it holds: $\Phi = \Phi(r)$, $\Psi = \Psi(r)$, with

$$r = \sqrt{x^2 + y^2 + z^2}, \quad (11.30)$$

because of spherical symmetry and $\rho = m \delta$. Using these properties, we deduce the field equation and discuss the existence of solutions of the above mentioned type.

At first we make some helpful general considerations: The functions Φ and Ψ are determined by eq. (11.25) for $i = j = 0$ and the trace of eq.

(11.25). If these two equations hold, then all other component-equations are automatically satisfied. For the 00-equation we need R_{00} :

$$R_{00} = -\Delta\Phi. \quad (11.31)$$

Here, the Laplacian is given as usual by

$$\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}. \quad (11.32)$$

For the inverse metric we get

$$g^{ij} = \text{diag}(1/(1-2\Phi), -1/(1+2\Psi), -1/(1+2\Psi), -1/(1+2\Psi)) \quad (11.33)$$

and $1/(1-2\Phi) = 1 + 2\Phi + h(\Phi)$, where $h(\Phi)$ is quadratic in Φ and vanishes after linearization. So we get

$$g^{ij} = \eta^{ij} - f^{ij}. \quad (11.34)$$

In our coordinate system, f^{ij} equals f_{ij} for all i, j . For the curvature scalar we get

$$R = 2(2\Delta\Psi - \Delta\Phi). \quad (11.35)$$

Moreover, we need expressions of the type $\square^k R$. $\square R$ is defined by $\square R = R_{;ij} g^{ij}$, where “;” denotes the covariant derivative. Remarks: Because of linearization we may replace the covariant derivative with the partial one. So we get

$$\square^k R = (-1)^k 2(-\Delta^{k+1}\Phi + 2\Delta^{k+1}\Psi) \quad (11.36)$$

and after some calculus

$$-8\pi\rho = \Delta\Phi + \Delta\Psi. \quad (11.37)$$

We use eq. (11.37) to eliminate Ψ from the system. So we get an equation relating Φ and $\rho = m\delta$.

$$-4\pi \left(\rho + 8 \sum_{k=0}^p a_k (-1)^k \Delta^{k+1} \rho \right) = \Delta\Phi + 6 \sum_{k=0}^p a_k (-1)^k \Delta^{k+2} \Phi. \quad (11.38)$$

In spherical coordinates it holds

$$\Delta\Phi = \frac{2}{r}\Phi_{,r} + \Phi_{,rr}, \quad (11.39)$$

because Φ depends on the radial coordinate r only.

We apply the following lemma: In the sense of distributions it holds

$$\Delta\left(\frac{1}{r}e^{-r/l}\right) = \frac{1}{rl^2}e^{-r/l} - 4\pi\delta. \quad (11.40)$$

Now we are ready to solve the whole problem. We assume

$$\Phi = \frac{m}{r} \left(1 + \sum_{i=0}^q c_i \exp(-r/l_i)\right), \quad l_i > 0. \quad (11.41)$$

Without loss of generality we may assume $l_i \neq l_j$ for $i \neq j$. Then eq. (11.38) together with that lemma yield

$$\begin{aligned} 8\pi \sum_{k=0}^p a_k (-1)^k \Delta^{k+1} \delta &= \sum_{i=0}^q \left(\frac{c_i}{t_i} + 6 \sum_{k=0}^p a_k (-1)^k \frac{c_i}{t_i^{k+2}} \right) \frac{1}{rl^2} e^{-r/l_i} \\ &\quad - 4\pi \sum_{i=0}^q \left(c_i + 6 \sum_{k=0}^p a_k (-1)^k \frac{c_i}{t_i^{k+1}} \right) \delta \\ &\quad + 24\pi \sum_{k=0}^p \sum_{j=k}^p \sum_{i=0}^p c_i a_j (-1)^{j+1} \frac{1}{t_i^{j-k}} \Delta^{k+1} \delta \end{aligned} \quad (11.42)$$

where $t_i = l_i^2$; therefore also $t_i \neq t_j$ for $i \neq j$. This equation is equivalent to the system

$$\sum_{i=0}^q c_i = 1/3, \quad (11.43)$$

$$\sum_{i=0}^q \frac{c_i}{t_i^s} = 0, \quad s = 1, \dots, p \quad (11.44)$$

$$t_i^{p+1} + 6 \sum_{k=0}^p a_k (-1)^k t_i^{p-k} = 0, \quad i = 0, \dots, q. \quad (11.45)$$

From eq. (11.45) we see that the values t_i represent $q + 1$ different solutions of one polynomial. This polynomial has the degree $p + 1$. Therefore $q \leq p$.

Now we use eqs. (11.43) and (11.44). They can be written in matrix form as

$$\begin{pmatrix} 1 & \dots & 1 \\ 1/t_0 & \dots & 1/t_q \\ \dots & & \\ 1/t_0^p & \dots & 1/t_q^p \end{pmatrix} \cdot \begin{pmatrix} c_0 \\ c_1 \\ \dots \\ c_q \end{pmatrix} = \begin{pmatrix} 1/3 \\ 0 \\ \dots \\ 0 \end{pmatrix} \quad (11.46)$$

Here, the first $q + 1$ rows form a regular matrix, the Vandermonde matrix. Therefore, we get

$$1/t_i^j = \sum_{k=0}^q \lambda_{jk} / t_i^k \quad j = q + 1, \dots, p \quad (11.47)$$

with certain coefficients λ_{jk} i.e., the remaining rows depend on the first $q + 1$ ones. If $\lambda_{j0} \neq 0$ then the system has no solution. So $\lambda_{j0} = 0$ for all $q + 1 \leq j \leq p$. But for $q < p$ we would get

$$1/t_i^q = \sum_{k=1}^q \lambda_{q+1,k} / t_i^{k-1} \quad (11.48)$$

and this is a contradiction to the above stated regularity. Therefore p equals q . The polynomial in (14) may be written as

$$6 \cdot \begin{pmatrix} 1 & 1/t_0 & \dots & 1/t_0^p \\ \dots & & & \\ 1 & 1/t_p & \dots & 1/t_p^p \end{pmatrix} \cdot \begin{pmatrix} a_0 \\ \dots \\ (-1)^p a_p \end{pmatrix} = \begin{pmatrix} -t_0 \\ \dots \\ -\tau_p \end{pmatrix} \quad (11.49)$$

This matrix is again a Vandermonde one, i.e., there exists always a unique solution $(a_0, \dots, a_p)$, which are the coefficients of the quantum corrections to the Einstein equation, such that the Newtonian limit of the corresponding

gravitational field equation is a sum of Newtonian and Yukawa potential with prescribed lengths l_i . A more explicit form of the solution is given in section 11.6.

11.5 Discussion of the weak-field limit

Let us give some special examples of the deduced formulas of the Newtonian limit of the theory described by the Lagrangian (11.22). If all the a_i vanish we get of course the usual Newton theory

$$\Phi = \frac{m}{r}, \quad \Delta\Phi = -4\pi\delta. \quad (11.50)$$

Φ and Ψ refer to the metric according to eq. (11.29). For $p = 0$ we get for $a_0 < 0$

$$\Phi = \frac{m}{r} \left[1 + \frac{1}{3} e^{-r/\sqrt{-6a_0}} \right] \quad (11.51)$$

cf. [249] and

$$\Psi = \frac{m}{r} \left[1 - \frac{1}{3} e^{-r/\sqrt{-6a_0}} \right] \quad (11.52)$$

cf. [208]. For $a_0 > 0$ no Newtonian limit exists.

For $p = 1$, i.e., the theory following from sixth-order gravity

$$\mathcal{L} = \left(R + a_0 R^2 + a_1 R \square R \right) \sqrt{-g}, \quad (11.53)$$

we get, see [187]

$$\Phi = \frac{m}{r} \left[1 + c_0 e^{-r/l_0} + c_1 e^{-r/l_1} \right] \quad (11.54)$$

and

$$\Psi = \frac{m}{r} \left[1 - c_0 e^{-r/l_0} - c_1 e^{-r/l_1} \right] \quad (11.55)$$

where

$$c_{0,1} = \frac{1}{6} \mp \frac{a_0}{2\sqrt{9a_0^2 + 6a_1}} \quad (11.56)$$

and

$$l_{0,1} = \sqrt{-3a_0 \pm \sqrt{9a_0^2 + 6a_1}}. \quad (11.57)$$

This result is similar in structure but has different coefficients as in fourth-order gravity with included square of the Weyl tensor in the Lagrangian.

The Newtonian limit for the degenerated case $l_0 = l_1$ can be obtained by a limiting procedure as follows: As we already know $a_0 < 0$, we can choose the length unit such that $a_0 = -1/3$. The limiting case $9a_0^2 + 6a_1 \rightarrow 0$ may be expressed by $a_1 = -1/6 + \varepsilon^2$. After linearization in ε we get:

$$l_i = 1 \pm \sqrt{3/2} \varepsilon c_i = 1/6 \pm 1/(6\sqrt{6}\varepsilon) \quad (11.58)$$

and applying the limit $\varepsilon \rightarrow 0$ to the corresponding fields Φ and Ψ we get

$$\begin{aligned} \Phi &= m/r \{1 + (1/3 + r/6)e^{-r}\} \\ \Psi &= m/r \{1 - (1/3 + r/6)e^{-r}\}. \end{aligned} \quad (11.59)$$

For the general case $p > 1$, the potential is a complicated expression, but some properties are explicitly known, these hold also for $p = 0, 1$. One gets

$$\Phi = m/r \left(1 + \sum_{i=0}^p c_i \exp(-r/l_i) \right) \quad (11.60)$$

and

$$\Psi = m/r \left(1 - \sum_{i=0}^p c_i \exp(-r/l_i) \right) \quad (11.61)$$

where $\sum c_i = 1/3$; $\sum$ means $\sum_{i=0}^p$ and l_i and c_i are, up to permutation of indices, uniquely determined by the Lagrangian.

There exist some inequalities between the coefficients a_i , which must be fulfilled in order to get a physically acceptable Newtonian limit. By this phrase we mean that besides the above conditions, additionally the fields Φ

and Ψ vanish for $r \rightarrow \infty$ and that the derivatives $d\Phi/dr$ and $d\Psi/dr$ behave like $O(1/r^2)$. These inequalities express essentially the fact that the l_i are real, positive, and different from each other. The last of these three conditions can be weakened by allowing the c_i to be polynomials in r instead of being constants, cf. the example with $p = 1$ calculated above.

The equality $\sum c_i = 1/3$ means that the gravitational potential is unbounded and behaves, up to a factor $4/3$, like the Newtonian potential for $r \approx 0$. The equation $\Phi + \Psi = 2m/r$ enables us to rewrite the metric as

$$ds^2 = (1 - 2\theta) \left[(1 - 2m/r) dt^2 - (1 + 2m/r)(dx^2 + dy^2 + dz^2) \right], \quad (11.62)$$

which is the conformally transformed linearized Schwarzschild metric with the conformal factor $1 - 2\theta$, where

$$\theta = \frac{m}{r} \sum c_i e^{-r/l_i} \quad (11.63)$$

can be expressed as functional of the curvature scalar, this is the linearized version of the conformal transformation theorem, cf. [218]. For an arbitrary matter configuration the gravitational potential can be obtained by the usual integration procedure.

11.6 A homogeneous sphere

For general p and characteristic lengths l_i fulfilling $0 < l_0 < l_1 < \dots < l_p$ we write the Lagrangian as

$$\begin{aligned} \mathcal{L} = R - \frac{R}{6} \Big[& (l_0^2 + \dots + l_p^2)R + (l_0^2 l_1^2 + l_0^2 l_2^2 \dots + l_{p-1}^2 l_p^2) \square R + \\ & (l_0^2 l_1^2 l_2^2 + \dots + l_{p-2}^2 l_{p-1}^2 l_p^2) \square^2 R + \dots + l_0^2 \cdot l_1^2 \cdot \dots \cdot l_p^2 \square^p R \Big] \end{aligned} \quad (11.64)$$

the coefficients in front of $\square^i R$ in this formula read

$$\sum_{0 \leq j_0 < j_1 < \dots < j_i \leq p} \prod_{m=0}^i l_{j_m}^2 . \quad (11.65)$$

Using this form of the Lagrangian, the gravitational potential of a point mass reads

$$\Phi = \frac{m}{r} \left[1 + \frac{1}{3} \sum_{i=0}^p (-1)^{i+1} \prod_{j \neq i} \left| \frac{l_j^2}{l_i^2} - 1 \right|^{-1} e^{-r/l_i} \right] , \quad (11.66)$$

$$\Psi = \frac{m}{r} \left[1 - \frac{1}{3} \sum_{i=0}^p (-1)^{i+1} \prod_{j \neq i} \left| \frac{l_j^2}{l_i^2} - 1 \right|^{-1} e^{-r/l_i} \right] . \quad (11.67)$$

For a homogeneous sphere of radius r_0 and mass m we get

$$\Phi = \frac{m}{r} \left[1 + \frac{1}{r_0^3} \sum_{i=0}^p e^{-r/l_i} l_i^2 \tilde{c}_i (r_0 \cosh(r_0/l_i) - l_i \sinh(r_0/l_i)) \right] , \quad (11.68)$$

where

$$\tilde{c}_i = (-1)^{i+1} \prod_{j \neq i} \left| \frac{l_j^2}{l_i^2} - 1 \right|^{-1} . \quad (11.69)$$

12 Cosmic strings and gravitational waves

We consider strings with the Nambu action as extremal surfaces in a given spacetime, thus, we ignore their back reaction. Especially, we look for strings sharing one symmetry with the underlying spacetime. If this is a non-null symmetry the problem of determining the motion of the string can be dimensionally reduced. Following [232], we get exact solutions for the following cases: straight and circle-like strings in a Friedmann background, straight strings in an anisotropic Kasner background, different types of strings in the metric of a gravitational wave. The solutions will be discussed.

12.1 Introduction to cosmic strings

To give detailed arguments for considering strings means carrying coals to Newcastle. So we only list the main points: A string is, generally speaking, an object possessing a two-dimensional world surface, which is also called a world sheet, in contrast to a point particle possessing a one-dimensional world line, cf. the review article by Vilenkin [261]. In details we have

1. One considers strings with the Nambu action in a D -dimensional flat spacetime. The theory can be consistently quantized for $D = 26$ only: otherwise the light cone quantization leads to a breakdown of Lorentz covariance, cf. Green, Schwarz and Witten [98]. But we consider a classical non-quantized theory only and require $D = 4$ henceforth.
2. Cosmic strings are one-dimensional topological defects in gauge field theories. They have a large mass and can be seeds for larger objects by accretion processes, cf. Zeldovich [276].
3. One looks for solutions of the Einstein equation with distribution-

valued energy-momentum tensor whose support is a two-dimensional sub-manifold of indefinite signature. The equation of state is $p_z = -\mu$ and the solution is called cosmic string. These are good candidates for the origin of structure, especially for the formation of galaxies, in the early universe.

4. If one looks for strings according to 2. or 3. in a given spacetime, i.e., with negligible back reaction of the string onto spacetime geometry, one arrives at the Nambu action, too, see Nielsen and Olesen [171] for 2. and Geroch and Traschen [87] for 3. In other words, cosmic strings are extremal, maximal or minimal in dependence of the boundary conditions, surfaces of indefinite signature in a given spacetime. This approach is justified if the diameter of the string is small compared with the curvature radius of the underlying manifold and if for the string tension $\alpha' \ll 1/G$ holds; we use units with $c = 1$. Normally, one thinks in orders of magnitude $G\alpha' = 10^{-6\pm 2}$, see Brandenberger [39]. In other words, the string is supposed to possess a mass per unit length of $10^{22\pm 2}$ g/cm, if the phase transition is supposed to be at the GUT-scale, see Frolov and Serebriany [80].

In [81], a stationary string in the Kerr-Newman metric has been considered. We use the method developed there and apply it to other cases. In this chapter we shall adopt the 4. approach and try to give some geometric insights into the motion of a string. To this end we give a sample of closed-form solutions for a special class of string solutions: strings which share a spacelike or timelike isometry with the underlying background metric.

12.2 The main formulae

The string is a two-dimensional extremal surface of indefinite signature. Let us take coordinates $(\tau, \sigma) = (y^A)$, $A = 0, 1$ within the string and coordinates

(x^i) , $i = 0, 1, 2, 3$ for the spacetime V_4 with the metric g_{ij} . The string is given by specifying the four functions $x^i(y^A)$. The signature for the metric g_{ij} is $(+ - - -)$. The tangents to the string are

$$e_A^i = \partial x^i / \partial y^A \quad (12.1)$$

and induce the metric

$$h_{AB} = e_A^i e_B^j g_{ij} \quad (12.2)$$

at the string world sheet. The signature of the string is required to be $(+ -)$, thus

$$h \equiv \det h_{AB} < 0. \quad (12.3)$$

The action to be varied is

$$I = \frac{1}{2\alpha'} \int \int \sqrt{-h} d\sigma d\tau. \quad (12.4)$$

Instead of writing down the full equations we specialize to the case we are interested here: we require that an isometry for both the underlying spacetime and the string exists. Let k_i be a non-null hypersurface-orthogonal Killing vector field, i.e.,

$$k_i k^i \neq 0, \quad k_{[i} k_{j;k]} = 0, \quad k_{(i;j)} = 0. \quad (12.5)$$

Then there exist coordinates x^i ($x^0 = t$) such

$$ds^2 = g_{00} dt^2 + g_{\alpha\beta} dx^\alpha dx^\beta. \quad (12.6)$$

$\alpha, \beta = 1, 2, 3$, the g_{ij} do not depend on t , $k_i = \partial/\partial t$. The sign of g_{00} is determined by the condition $g_{00} k_i k^i > 0$: for timelike k_i we have $g_{00} > 0$ and for spacelike k_i we have $g_{00} < 0$. The requirement that k_i is also an isometry of the string gives us the possibility to specify the functions $x^i(y^A)$ to be $t = \tau$, x^α depends on σ only. Then we get

$$e_0^i = (1, 0, 0, 0), \quad e_1^i = (0, dx^\alpha/d\sigma) \quad (12.7)$$

and $h_{01} = 0$, $h_{00} = g_{00}$,

$$I = \frac{1}{2\alpha'} \int \int \sqrt{-g_{00} g_{\alpha\beta} dx^\alpha/d\sigma dx^\beta/d\sigma} d\sigma d\tau. \quad (12.8)$$

The integrand does not depend on τ , so we can omit the τ integration. Therefore, extremizing eq. (12.8) is the same as solving the geodesic equation for the auxiliary metric $f_{\alpha\beta}$ of a V_3 defined by

$$f_{\alpha\beta} = -g_{00}g_{\alpha\beta}. \quad (12.9)$$

Remarks: 1. If k_i is not hypersurface-orthogonal then we get contains terms with $g_{0\alpha}$, and one has to add $g_{0\alpha}g_{0\beta}$ to the r.h.s. of the definition of $f_{\alpha\beta}$.

2. The dimensional reduction is not fully trivial: the geodesic equation for metric $f_{\alpha\beta}$ corresponds to compare the strings in eq. (12.8) with other strings sharing the same isometry induced by k_i , whereas the Nambu action has to be compared with all other strings, too. But writing down all full equations or counting the degrees of freedom one can see that at our circumstances no difference appears.

3. For a timelike k_i , $f_{\alpha\beta}$ is positive definite, and the condition $h < 0$ is automatically fulfilled. On the other hand, for a spacelike k_i , $f_{\alpha\beta}$ is of signature $(+ - -)$, and the validity of $h < 0$ requires the vector $dx^\alpha/d\sigma$ to be timelike, i.e., the root in eq. (12.8) has to be real.

12.3 The string in a Friedmann model

Now we specify the underlying V_4 to be a spatially flat Friedmann model

$$ds^2 = dt^2 - a^2(t)(dx^2 + dy^2 + dz^2). \quad (12.10)$$

12.3.1 The open string

First we use the spacelike Killing vector $\partial/\partial x$ of metric (12.10). We have in mind an infinitely long straight string moving through the expanding universe. We apply the formalism deduced in the previous section and get the following result: we write metric (12.10) in the form

$$ds^2 = -a^2 dx^2 + dt^2 - a^2(dy^2 + dz^2). \quad (12.11)$$

The metric $f_{\alpha\beta}$ then reads

$$ds_{(3)}^2 = a^2 dt^2 - a^4(dy^2 + dz^2). \quad (12.12)$$

Without loss of generality the string is situated at $z = 0$ and moves into the y -direction according to

$$\begin{aligned} ds_{(3)}^2 &= a^2 dt^2 - a^4 dy^2 \\ \ddot{t} + \frac{1}{a} \frac{da}{dt} \dot{t}^2 + 2 \frac{da}{dt} a \dot{y}^2 &= 0 \quad \text{with} \quad \cdot = d/d\lambda \\ a^4 \dot{y} &= M \\ a^2 \dot{t}^2 - a^4 \dot{y}^2 &= 1 \\ y(t) &= M \int \frac{dt}{a(t) \sqrt{a^4(t) + M^2}} \end{aligned} \quad (12.13)$$

where M is an integration constant. The natural distance of the string from the origin is

$$s(t) = a(t) \cdot y(t). \quad (12.14)$$

1. Example. Let $a(t) = t^n$, $n \geq 0$ then for $t \gg 1$, cf. Stein-Schabes and Burd [247],

$$y(t) \approx Mt^{1-3n}, \quad s(t) \approx Mt^{1-2n} \quad \text{for} \quad n \neq 1/3 \quad (12.15)$$

and $y(t) \approx M \ln t$ for $n = 1/3$. Interpretation: $n = 0$, i.e., the absence of gravity, implies a linear motion as it must be the case. We have

$$\lim_{t \rightarrow \infty} y(t) = \infty \quad (12.16)$$

if and only if $n \leq 1/3$, i.e., for $n > 1/3$ the string comes to rest with respect to the cosmic background after a finite time. A more stringent condition to be discussed is that the string comes to rest in a natural frame of a suitably chosen reference galaxy. This means

$$\lim_{t \rightarrow \infty} s(t) < \infty \quad (12.17)$$

and is fulfilled for $n \geq 1/2$. Therefore, the most interesting cases $n = 1/2$, the radiation model and $n = 2/3$, the Einstein-de Sitter dust model, have the property that straight open strings come to rest after a sufficiently long time independently of the initial conditions.

2. Example. Let $a(t) = e^{Ht}$, the inflationary model, $H > 0$. Then $y(t) = e^{-3Ht}$, $s(t) = e^{-2Ht}$. We have the same result as in the first example with $n \gg 1$.

3. Example. Let

$$a(t) = -t^{2/3}(1 + t^{-2} \cos(mt)). \quad (12.18)$$

This background metric is from damped oscillations of a massive scalar field or, equivalently, from fourth-order gravity $L = R + m^{-2} R^2$. We get with eq. (12.13)

$$y(t) \approx -1/t - ct^{-4} \sin mt, \quad c = \text{const.} \quad \text{for } t \rightarrow \infty, \quad (12.19)$$

which is only a minor modification of the result of the first example with $n = 2/3$, therefore, one should not expect a kind of resonance effect between the open string and a massive scalar field.

12.3.2 The closed string

We insert $dy^2 + dz^2 = dr^2 + r^2 d\Phi^2$ into metric (12.10) and use the spacelike Killing vector $\partial/\partial\Phi$: its trajectories are circles, so we have in mind a closed string with radius r moving and oscillating in the expanding universe. The corresponding geodesic equation leads to

$$\frac{d}{d\lambda}(-a^4 r^2 \dot{r}) - a^2 r \dot{t}^2 + a^4 r(\dot{x}^2 + \dot{r}^2) = 0, \quad (12.20)$$

$$\frac{d}{d\lambda}(a^4 r^2 \dot{x}) = 0, \quad (12.21)$$

$$\dot{t}^2 - a^2(\dot{x}^2 + \dot{r}^2) = \frac{1}{a^2 r^2}, \quad (12.22)$$

where the dot denotes $d/d\lambda$, λ is the natural parameter along the geodesic. We are mainly interested in solutions not moving into the x -direction, to understand the oscillating behaviour. Inserting $x = 0$ into this system of 3 equations and using the fact that always $\dot{t} \neq 0$ holds, this system can be brought into the form

$$r \cdot r'' a^3 - 2rr' a^4 a' - r'^2 a^3 + 3rr' a^2 a' + a = 0, \quad (12.23)$$

where the dash denotes d/dt .

4. Example. Let $a = 1$, i.e., we have the flat Minkowski spacetime. Then this equation reduces to

$$rr'' = r'^2 - 1. \quad (12.24)$$

The solution reads

$$r(t) = r_0 \cos((t - t_0)/r_0), \quad r' = -\sin((t - t_0)/r_0). \quad (12.25)$$

This is the oscillating solution for closed strings. At points t where $r = 0$, we have $|r'| = 1$. These points are the often discussed cusp points of the

string, where the interior string metric becomes singular and the motion approximates the velocity of light, cf. Thompson [254].

Let us compare eq. (12.24) with the analogous equation for a positive definite back-ground metric. It reads

$$rr'' = r'^2 + 1 \quad (12.26)$$

and has solutions with $\cosh$ instead of $\cos$. This is the usual minimal surface taken up e.g. by a soap-bubble spanned between two circles.

5. Example. Let $a(t) = t^n$, then we have to solve

$$rr''t^{2n} - 2nrr'3t^{4n-1} - r'^2t^{2n} + 3nrr't^{2n-1} + 1 = 0. \quad (12.27)$$

This equation governs the radial motion of a circle-like closed string in a Friedmann background. The solutions can be hardly obtained by analytic methods.

12.4 The string in an anisotropic Kasner background

Now we take as background metric

$$ds^2 = dt^2 - a^2(t)dx^2 - b^2(t)dy^2 - c^2(t)dz^2 \quad (12.28)$$

and $\partial/\partial x$ as Killing vector. Astonishingly, the anisotropy has only a minor influence on the motion of the string, so we get mainly the same formulae as in the isotropic case: the geodesic equations are

$$\frac{d}{d\lambda}(a^2b^2\dot{y}) = 0 \quad \text{hence} \quad a^2b^2\dot{y} = M_1 \quad (12.29)$$

$$\frac{d}{d\lambda}(a^2c^2\dot{z}) = 0 \quad \text{hence} \quad a^2c^2\dot{z} = M_2 \quad (12.30)$$

$$a^2\dot{t}^2 - a^2c^2\dot{z}^2 = 1 \quad (12.31)$$

and can be integrated to yield for $M_1 M_2 \neq 0$

$$y(r) = \int \left(\frac{a^2 b^2}{M_1^2} + \frac{M_2^2}{M_1^2} \cdot \frac{b^2}{c^2} + 1 \right)^{-1/2} \frac{dt}{b(t)}. \quad (12.32)$$

The equation for $z(t)$ can be obtained from this equation by interchanging $b \leftrightarrow c$ and $M_1 \leftrightarrow M_2$.

For the Kasner metric we have $a = t^p$, $b = t^q$, $c = t^r$, $p + q + r = p^2 + q^2 + r^2 = 1$. We get

$$y(r) = \int \frac{dt}{t^q \sqrt{t^{2(p+q)} + t^{2(q-r)} + 1}}. \quad (12.33)$$

The behaviour for $t \rightarrow \infty$ in dependence of the values p , q , r can be seen from this equation.

12.5 The string in a gravitational wave

As background metric we use the plane-wave ansatz

$$ds^2 = 2dudv + p^2(u)dy^2 + q^2(u)dz^2. \quad (12.34)$$

This metric represents a solution of Einstein's vacuum equation if

$$q \, d^2 p / du^2 + p \, d^2 q / du^2 = 0. \quad (12.35)$$

Let us take $\partial/\partial y$ as Killing vector. Then the geodesic equations for the auxiliary metric are

$$\frac{d}{d\lambda}(p^2 \dot{u}) = 0 \quad \text{hence} \quad p^2 \dot{u} = M_1 \neq 0, \quad (12.36)$$

$$\frac{d}{d\lambda}(p^2 q^2 \dot{z}) = 0 \quad \text{hence} \quad p^2 q^2 \dot{z} = M_2, \quad (12.37)$$

$$-2p^2 \dot{u} \dot{v} - p^2 q^2 \dot{z}^2 = 1. \quad (12.38)$$

If we take u as new independent variable, which is possible because of $\dot{u} \neq 0$, we get the solutions

$$\begin{aligned} z(u) &= \frac{M_2}{M_1} \int \frac{du'}{q^2(u')} , \\ v(u) &= \frac{-M_2^2}{2M_1^2} \int \left(\frac{1}{q^2(u')} - p^2(u') \right) du' . \end{aligned} \quad (12.39)$$

As an example let us take

$$p(u) = \sin u , \quad q(u) = \sinh u \quad (12.40)$$

which produces a vacuum solution. With this choice of p and q we get

$$z(u) = M \coth u , \quad v(u) = -M^2(4 \coth u + \sin 2u - 2u)/8 . \quad (12.41)$$

Another Killing vector of the plane wave is $\partial/\partial v$, but it is a null Killing vector and so the reduction used above does not work. But there exists a further non-null Killing vector of this plane wave: It reads

$$k_i = (0, y, H, 0), \quad \text{where} \quad H = - \int p^{-2} du . \quad (12.42)$$

By a coordinate transformation

$$\begin{aligned} u &= t & y &= w \cdot e^{-G} & G(u) &= \int \frac{du}{p^2 H} \\ z &= k & v &= x + \frac{1}{2H} e^{-2G} (w^2 - 1) \end{aligned} \quad (12.43)$$

we get the form

$$ds^2 = e^{-2G} p^2 dw^2 + 2dxdt + \frac{e^{-2G}}{p^2 H^2} dt^2 + q^2 dk^2 \quad (12.44)$$

and have to solve the equations

$$\begin{aligned} \frac{d}{d\lambda} (p^2 q^2 e^{-2G} \dot{k}) &= 0 , & \frac{d}{d\lambda} (p^2 e^{-2G} \dot{t}) &= 0 , \\ -p^2 q^2 e^{-2G} \dot{k}^2 - \frac{1}{H^2} e^{-4G} \dot{t}^2 - 2p^2 e^{-2G} \dot{x} \dot{t} &= 1 . \end{aligned} \quad (12.45)$$

For the special case $\dot{k} = 0$ one has finally

$$x(t) = -\frac{1}{2} \int e^{-2G} \left(\frac{p^2}{M_2^2} + \frac{1}{H^2 p^2} \right) dt. \quad (12.46)$$

Let us suppose a spatially flat Friedmann model eq. (12.10) with scale factor $a(t) = t^n$, and $n = 2/3$, representing the Einstein-de Sitter dust model, or $n = 1/2$, representing the radiation model. There we consider an open string which is only a little bit curved such that the approximation of a straight string is applicable. At time $t = t_0 > 0$ we can prescribe place and initial velocity v_0 of the string arbitrarily and get for $t \rightarrow \infty$ the behaviour

$$s(t) \approx \text{const} + M(v_0)t^{1-2n} \quad (12.47)$$

where s denotes the natural distance from the origin. That means, for the cases $n = 2/3$ and $n = 1/2$ we are interested in, the string comes to rest for large values t at a finite distance from the origin.

We compare this result with the analogous motion of a point-particle in the same background: in the same approximation we get

$$s(t) \approx \text{const} + M(v_0)t^{1-n}, \quad (12.48)$$

i.e., $|s| \rightarrow \infty$ as $t \rightarrow \infty$, a totally other type of motion. It should be mentioned that in the absence of gravity, i.e., $n = 0$, both motions are of the same type, but otherwise not.

The remaining calculations above indicate that the interaction of the motion of the string with scalar field oscillations, with anisotropy, and with gravitational waves is quite weak, we did not find any type of resonance effects.

13 Bibliography

- [1] Altshuler, B.: 1990, *Class. Quant. Grav.* **7**, 189. *123*
- [2] Amendola, L.: 1993, *Phys. Lett. B* **301**, 175. 1999, *Phys. Rev. D* **60**, 043501. *129*
- [3] Amendola, L., Battaglia Mayer, A., Capozziello, S., Gottlöber, S., Müller, V., Occhionero, F., Schmidt, H.-J.: 1993, *Class. Quant. Grav.* **10**, L43. *126*
- [4] Amendola, L., Capozziello, S., Litterio, M., Occhionero, F.: 1992, *Phys. Rev. D* **45**, 417. *126*
- [5] Anandan, J.: 1983, in: Hu, N., ed., *Proc. 3. Marcel Grossmann Meeting B*, Amsterdam NHPC, 941. *160*
- [6] Anderson, P.: 1986, *Phys. Lett. B* **169**, 31. *111*
- [7] Arnowitt, R., Deser, S., Misner, C.: 1962, in: Witten, E., ed., *Gravitation, an introduction to current research*, Wiley New York. *123*
- [8] Bach, R.: 1921, *Math. Zeitschr.* **9**, 110. *145, 147, 156*
- [9] Bachmann, M., Schmidt, H.-J.: 2000, *Phys. Rev. D* **62**, 043515; gr-qc/9912068. *83*
- [10] Barabash, O., Shtanov, Y.: 1999, *Phys. Rev. D* **60**, 064008; astro-ph/9904144. *159*
- [11] Barraco, D., Hamity, V.: 2000, *Phys. Rev. D* **62**, 044027. 1999, *Gen. Relat. Grav.* **31**, 213. *112*
- [12] Barrow, J.: 1986, *Phys. Lett. B* **180**, 335. 1987, *Phys. Lett. B* **187**, 12. *123, 125, 126, 136*
- [13] Barrow, J., Götz, G.: 1989, *Class. Quant. Grav.* **6**, 1253. *122, 123*
- [14] Barrow, J., Matzner, R.: 1980, *Phys. Rev. D* **21**, 336. *59, 74*
- [15] Barrow, J., Saich, P.: 1990, *Phys. Lett. B* **249**, 406. *126*

- [16] Barrow, J., Sirousse-Zia, H.: 1989, Phys. Rev. D **39**, 2187. Erratum 1990, D **41**, 1362. 61, 125, 127, 139
- [17] Barrow, J., Tipler, F.: 1988, *The anthropic cosmological principle*, Cambridge University Press. 75
- [18] Battaglia Mayer, A., Schmidt, H.-J.: 1993, Class. Quant. Grav. **10**, 2441. 128
- [19] Belinsky, V., Grishchuk, L., Khalatnikov, I., Zeldovich, Y.: 1985, Phys. Lett. B **155**, 232. 89
- [20] Belinsky, V., Grishchuk, L., Zeldovich, Y., Khalatnikov, I.: 1985, Zh. Eksp. Teor. Fiz. **89**, 346. 71, 74
- [21] Belinsky, V., Khalatnikov, I.: 1987, Zh. Eksp. Teor. Fiz. **93**, 784. 59, 74
- [22] Belinsky, V., Lifshitz, E., Khalatnikov, I.: 1972, Sov. Phys. JETP **35**, 838. 59, 125⁷
- [23] Bergmann, O.: 1981, Phys. Lett. A **82**, 383. 9
- [24] Berkin, A.: 1990, Phys. Rev. D **42**, 1016. 1991, Phys. Rev. D **44**, 1020. 127, 129
- [25] Berkin, A., Maeda, K.: 1990, Phys. Lett. B **245**, 348. 1991, Phys. Rev. D **44**, 1691. 129, 164
- [26] Bičák, J.: 2000, Lect. Notes Phys. **540**, 1-126; gr-qc/0004016. 19
- [27] Bičák, J., Podolsky, J.: 1992, in: Abstracts Conf. Gen. Relat. 13, Cordoba, 12. 1995, Phys. Rev. D **52**, 887. 127
- [28] Bičák, J., Pravda, V.: 1998, Class. Quant. Grav. **15**, 1539. 44
- [29] Bicknell, G.: 1974, J. Phys. A **7**, 341. 1061. 61, 87, 103, 139
- [30] Bleyer, U., Liebscher, D.-E., Schmidt, H.-J., Zhuk, A.: 1990, Wiss. Zeitschr. d. Univ. Jena, Naturwiss. Reihe **39**, 20. 85

⁷The authors are in alphabetical order of the original Russian language version.

- [31] Bleyer, U., Schmidt, H.-J.: 1990, Int. J. Mod. Phys. A **5** (1990) 4671. *121*
- [32] Bojowald, M., Kastrup, H., Schramm, F., Strobl, T.: 2000, Phys. Rev. D **62** (2000) 044026. *111*
- [33] Borde, A.: 1994, Phys. Rev. D **50**, 3692. *124*
- [34] Borde, A., Vilenkin, A.: 1994, Phys. Rev. Lett. **72**, 3305. *124*
- [35] Borzeszkowski, H. v.: 1981, Ann. Phys. (Leipz.) **38**, 239. *146*
- [36] Borzeszkowski, H. v., Treder, H.-J.: 1994, Found. Phys. **24**, 949. *127*
- [37] Borzeszkowski, H. v., Treder, H.-J., Yourgrau, W.: 1978, Ann. Phys. (Leipz.) **35**, 471. *146, 156, 160*
- [38] Boulware, D.: 1973, Phys. Rev. D **8**, 2363. *45*
- [39] Brandenberger, R.: 1987, Int. J. Mod. Phys. A **2**, 77. *176*
- [40] Brauer, U., Rendall, A., Reula, O.: 1994, Class. Quant. Grav. **11**, 2283. *123*
- [41] Breizman, B., Gurovich, V., Sokolov, V.: 1970, Zh. Eksp. Teor. Fiz. **59**, 288. *125*
- [42] Buchdahl, H.: 1951, Acta Math. **85**, 63. *128, 134, 164*
- [43] Buchdahl, H.: 1962, Nuovo Cim. **23**, 141. *92*
- [44] Burd, A.: 1993, Class. Quant. Grav. **10**, 1495. *123*
- [45] Burd, A., Barrow, J.: 1988, Nucl. Phys. B **308**, 929. *122*
- [46] Calzetta, E.: 1989, Class. Quant. Grav. **6**, L227. *76, 140*
- [47] Calzetta, E., Sakellariadou, M.: 1992, Phys. Rev. D **45**, 2802. *127*
- [48] Campanelli, M., Lousto, C.: 1996, Phys. Rev. D **54**, 3854. *112*
- [49] Capozziello, S., Lambiase, G.: 2000, Gen. Relat. Grav. **32**, 295. 673. *112*
- [50] Carfora, M., Marzuoli, A.: 1984, Phys. Rev. Lett. **53**, 2445. *106*

- [51] Carmeli, M., Charach, C. : 1980, Phys. Lett. A **75**, 333. *9*
- [52] Chiba, T., Maeda, K.: 1994, Phys. Rev. D **50**, 4903. *124*
- [53] Cohen, M., Cohen, J.: 1971, in: Kuper, C., ed., *Relativity and Gravitation*, Gordon and Breach New York, 99. *51*
- [54] Coley, A., Tavakol, R.: 1992, Gen. Relat. Grav. **24**, 835. *125*
- [55] Collins, C., Szafron, D.: 1981, J. Math. Phys. **22**, 543. *9*
- [56] Contreras, C., Herrera, R., del Campo, S.: 1995, Phys. Rev. D **52**, 4349. *127*
- [57] Cotsakis, S., Demaret, J., De Rop, Y., Querella, L.: 1993, Phys. Rev. D **48**, 4595. *125*
- [58] Cotsakis, S., Flessas, G.: 1993, Phys. Lett. B **319**, 69. *127*
- [59] Coule, D.: 2000, Phys. Rev. D **62**, 124010; gr-qc/0007037. 1995, Class. Quant. Grav. **12**, 455; gr-qc/9408026. *83*
- [60] Dautcourt, G. : 1963, Math. Nachr. **27**, 277. 1964, Acta Phys. Polon. **25**, 637. *45, 54, 164*
- [61] Demaret, J., Querella, L., Scheen, C.: 1999, Class. Quant. Grav. **16**, 749. Demaret, J., Querella, L.: 1995, Class. Quant. Grav. **12**, 3085. *112*
- [62] Demianski, M.: 1984, Nature **307**, 140. *123*
- [63] Denisov, V. I.: 1972, J. Exp. Teor. Phys. **62**, 1990. *45*
- [64] Dzhunushaliev, V., Schmidt, H.-J.: 2000, J. Math. Phys. **41**, 3007; gr-qc/9908049. *156*
- [65] Ehlers, J., Kundt, W.: 1962, in: Witten, E., ed., *Gravitation, an introduction to current research*, Wiley New York, 49. *36*
- [66] Einstein, A.: 1914, Sitzungsber. Akad. d. Wiss. Berlin, 1030. 1916, Sitzungsber. Akad. d. Wiss. Berlin, 688. 1918, Sitzungsber. Akad. d. Wiss. Berlin, 154. 1921, Sitzungsber. Akad. d. Wiss. Berlin, 261. *35, 145*

- [67] Einstein, A.: 1939, Ann. Math. **40**, 922. *46*
- [68] Einstein, A., De Sitter, W.: 1932, Proc. Nat. Acad. Sci. **18**, 213. *19, 24*
- [69] Einstein, A., Pauli, W.: 1943, Ann. Math. **44**, 131. *146*
- [70] Ellis, G.: 1984, Ann. Rev. Astron. Astrophys. **22**, 157. Ellis, G.: 1967, J. Math. Phys. **8**, 1171. *13, 104*
- [71] Esteban, E., Kazanas, D.: 2001, Gen. Relat. Grav. **33**, 1281. *156*
- [72] Fabris, J., Pelinson, A., Shapiro, I.: 2001, Nucl. Phys. B **597**, 539; hep-th/0009197. *39*
- [73] Fabris, J., Reuter, S.: 2000, Gen. Relat. Grav. **32**, 1345. *83*
- [74] Faraoni, V., Gunzig, E., Nardone, P.: 1999, Fund. Cosmic Physics **20**, 121; gr-qc/9811047. *112*
- [75] Feldman, H.: 1990, Phys. Lett. B **249**, 200. *127*
- [76] Ferraris, M.: 1986, in: Fabbri, R., Modugno, M., eds., *VI Conv. Naz. Rel. Gen. e Fisica della Grav.*, Pitagora Edit. Bologna, 127. *110*
- [77] Fiedler, B., Schimming, R.: 1980, Rep. Math. Phys. **17**, 15. *39*
- [78] Fiedler, B., Schimming, R.: 1983, Astron. Nachr. **304**, 221. *146, 148, 149, 155*
- [79] Frehland, E.: 1972, Ann. Phys. (Leipz.) **28**, 91. *45*
- [80] Frolov, V., Serebriany, E.: 1987, Phys. Rev. D **35**, 3779. *176*
- [81] Frolov, V., Skarzhinsky, V., Zelnikov, A., Heinrich, O.: 1988, Preprint Potsdam-Babelsberg, PRE-ZIAP 88-14. *176*
- [82] Fulling, S.: 1973, Phys. Rev. D **7**, 2850. *58*
- [83] Fulling, S., Parker, L.: 1974, Ann. Phys. NY **87**, 176. *58*
- [84] Gemelli, G.: 1997, Gen. Relat. Grav. **29**, 1163. *39*
- [85] Gerbal, D., Sirousse-Zia, H.: 1989, C. R. Acad. Sci. Paris **309**, 353. *165*

- [86] Geroch, R.: 1969, Commun. Math. Phys. **13**, 180. *42, 64, 73, 115*
- [87] Geroch, R., Traschen, J.: 1987, Phys. Rev. D **36**, 1017. *176*
- [88] Gibbons, G. and Hawking, S.: 1977, Phys. Rev. D **15**, 2738. *123*
- [89] Gibbons, G., Hawking, S., Stewart, J.: 1987, Nucl. Phys. B **281**, 736. *85*
- [90] Goenner, H.: 1987, in: MacCallum, M., ed., *Proc. 11th Int. Conf. Gen. Rel. Grav.*, Cambridge University Press, 262. *110*
- [91] Gorbatenko, M., Pushkin, A., Schmidt, H.-J.: 2002, Gen. Relat. Grav. **34**, in press; gr-qc/0106025. *156*
- [92] Gottlöber, S.: 1984, Ann. Phys. (Leipz.) **41**, 45. 1984, Astron. Nachr. **305**, 1. 1987, Astrophys. Space Sc. **132**, 191. *59, 107, 109*
- [93] Gottlöber, S., Müller, V.: 1987, Class. Quant. Grav. **4**, 1427. *59*
- [94] Gottlöber, S., Müller, V., Schmidt, H.-J.: 1991, Astron. Nachr. **312**, 291. *129, 164*
- [95] Gottlöber, S., Müller, V., Schmidt, H.-J., Starobinsky, A.: 1992, Int. J. Mod. Phys. D **1**, 257. *126*
- [96] Gottlöber, S., Schmidt, H.-J.: 1984, Potsdamer Forschungen B **43**, 117. *59*
- [97] Gottlöber, S., Schmidt, H.-J., Starobinsky, A.: 1990, Class. Quantum Grav. **7**, 893. *129, 164, 165*
- [98] Green, M., Schwarz, J., Witten, E.: 1987, *Superstring theory*, Cambridge University Press. *175*
- [99] Griffiths, J.: 1995, Gen. Relat. Grav. **27**, 905. *13, 26*
- [100] Grishchuk, L., Sidorov, J.: 1988, in: Markov, M., ed., *Proc. 4. Sem. Quantum Gravity Moscow*, WSPC Singapore, 700. *85*
- [101] Gurovich, V., Schmidt, H.-J., Tokareva, I.: 2001, Gen. Relat. Grav. **33**, 591; gr-qc/0007046. *83*

- [102] Hall, G.: 1988, *Gen. Relat. Grav.* **20**, 671. 1990, *J. Math. Phys.* **31**, 1198. 1991, *Diff. Geom. Appl.* **1**, 35. *40*
- [103] Halliwell, J., Hawking, S.: 1985, in: Markov, M., ed., *Proc. 3. Sem. Quantum Gravity Moscow*, WSPC Singapore, 509. *85*
- [104] Harvey, A.: 1990, *Class. Quant. Grav.* **7**, 715. *41*
- [105] Havas, P.: 1977, *Gen. Relat. Grav.* **8**, 631. *160*
- [106] Hawking, S.: 1984, *Nucl. Phys. B* **239**, 257. 1984, in: De Witt, B., Stora, R., eds., *Relativity, Groups, and Topology II*, North Holland PC Amsterdam. *59*
- [107] Hawking, S., Ellis, G.: 1973, *The large scale structure of space-time*, Cambridge University Press. *36*
- [108] Hawking, S., Luttrell, J.: 1984, *Nucl. Phys. B* **247**, 250. *59*
- [109] Hawking, S., Moss, I.: 1982, *Phys. Lett. B* **110**, 35. *123*
- [110] Hehl, F., McCrea, J., Mielke, E., Ne'eman, Y.: 1995, *Phys. Repts.* **258**, 1. *112*
- [111] Hindawi, A., Ovrut, B., Waldram, D.: 1996, *Phys. Rev. D* **53**, 5583. *112*
- [112] Hoyle, F., Narlikar, J.: 1963, *Proc. R. Soc. Lond. A* **273**, 1. *123*
- [113] Hübner, P., Ehlers, J.: 1991, *Class. Quant. Grav.* **8**, 333. *123*
- [114] Hwang, J.: 1991, *Class. Quant. Grav.* **8**, L133. *126*
- [115] Hwang, J., Noh, H.: 2001, *Phys. Lett. B* **506**, 13. *112*
- [116] Ishihara, R.: 1986, *Phys. Lett. B* **179**, 217. *112*
- [117] Israel, W.: 1966, *Nuovo Cim. B* **44**, 1. *45, 54*
- [118] Israel, W., Stewart, J.: 1980, in: Held, A., ed., *General Relativity and Gravitation II*, Plenum New York, 491. *46*
- [119] Ivashchuk, V., Melnikov, V.: 1995, *Grav. Cosmol.* **1**, 133; hep-th/9503223. *111*

- [120] Jakubiec, A., Kijowski, J.: 1988, Phys. Rev. D **37**, 1406. 1989, J. Math. Phys. **30**, 1073. *110*
- [121] Jankiewicz, C.: 1981, Acta Phys. Polon. **13**, 859. *160*
- [122] Jensen, L., Stein-Schabes, J.: 1987, Phys. Rev. D **35**, 1146. *127*
- [123] Jordan, P., Ehlers, J., Kundt, W.: 1960, Abh. Akad. Wiss. Mainz, Math./Nat. Kl. **2**, 21. *36*
- [124] Kao, W.: 2000, Phys. Rev. D **61**, 047501. *112*
- [125] Kerner, R.: 1982, Gen. Relat. Grav. **14**, 453. *94*
- [126] Kiefer, C.: 2001, in: Janke, W., Pelster, A., Schmidt, H.-J., Bachmann, M., eds., *Fluctuating paths and fields*, WSPC Singapore, 729. *83*
- [127] Kirsten, K., Cognola, G., Vanzo, L.: 1993, Phys. Rev. D **48**, 2813. *129*
- [128] Kleinert, H., Schmidt, H.-J.: 2000, *Cosmology with curvature-saturated gravitational lagrangian $R/\sqrt{1+l^4R^2}$* , Preprint gr-qc/0006074. *112*
- [129] Kluske, S., Schmidt, H.-J.: 1996, Astron. Nachr. **317**, 337; gr-qc/9503021. *121*
- [130] Kofman, L., Linde, A., Starobinsky, A.: 1996, Phys. Rev. Lett. **76**, 1011. *123*
- [131] Kottler, F.: 1918, Ann. Phys. (Leipz.) **56**, 410. *128*
- [132] Kowalski, O.: 1993, Nagoya Math. J. **132**, 1. *31*
- [133] Kowalski, O., Nikcevic, S.: 1995, *On Ricci eigenvalues of locally homogeneous Riemannian 3-manifolds*, preprint. *32*
- [134] Kramer, D., Stephani, H., Herlt, E., MacCallum, M.: 1980, *Exact solutions of Einstein's field equations*, Verl. d. Wiss. Berlin. *11*
- [135] Krasilnikov, V. A., Pavlov, V. I.: 1972, Vestnik Univ. Moskau **27**, 235. *46*
- [136] Krasinski, A.: 1980, Proc. Conf. General Relativity GR 9 Jena, 44. *9*

- [137] Krasinski, A.: 1997, *Inhomogeneous Cosmological Models*, Cambridge University Press. 9, 19
- [138] Kuchař, K.: 1968, Czech. J. Phys. B **18**, 435. 45
- [139] Kung, J.: 1996, Phys. Rev. D **53**, 3017. 112
- [140] Lake, K.: 1979, Phys. Rev. D **19**, 2847. 1981, Lett. Nuovo Cim. **30**, 186. 1993, J. Math. Phys. **34**, 5900. 41, 42, 45, 46
- [141] Lanczos, K.: 1924, Ann. Phys. (Leipz.) **74**, 518. 45
- [142] Lee, C. W.: 1978, Proc. R. Soc. Lond. A **364**, 295. 10
- [143] López, C.: 1981, Nuovo Cim. B **66**, 17. 45
- [144] Lu, M. and Wise, M.: 1993, Phys. Rev. D **47**, R3095. 129
- [145] Lukash, V. N., Schmidt, H.-J.: 1988, Astron. Nachr. **309**, 25. 71, 72, 107, 109
- [146] MacCallum, M.: 1971, Commun. Math. Phys. **20**, 57. 1972, Phys. Lett. A **40**, 385. 15, 127, 139
- [147] MacCallum, M., Stewart, J., Schmidt, B.: 1970, Commun. Math. Phys. **17**, 343. 127, 139
- [148] Macias, A.: 1999, Gen. Relat. Grav. **31**, 653. 83
- [149] Madsen, M., Ellis, G.: 1988, Mon. Not. R. Astron. Soc. **234**, 67. 76
- [150] Maeda, K.: 1988, Phys. Rev. D **37**, 858. 1989, Phys. Rev. D **39**, 3159. 126
- [151] Maeda, K., Sato, H.: 1983, Progr. Theor. Phys. **70**, 772. 46, 54
- [152] Maeda, K., Stein-Schabes, J., Futamase, T.: 1989, Phys. Rev. D **39**, 2848. 126
- [153] Martin, G., Mazzitelli, F.: 1994, Phys. Rev. D **50**, R613. 129
- [154] Mignemi, S.: 1996, Ann. Phys. NY 245 (1996) 23. 111
- [155] Mijic, M., Stein-Schabes, J.: 1988, Phys. Lett. B **203**, 353. 126

- [156] Mirzabekian, A., Vilkovisky, G., Zhytnikov, V.: 1996, Phys. Lett. B **369**, 215. *142*
- [157] Mondaini, R., Vilar, L.: 1993, Int. J. Mod. Phys. D **2**, 477. *124*
- [158] Mongan, T.: 2001, Gen. Relat. Grav. **33**, 1415. *83*
- [159] Moniz, P.: 1993, Phys. Rev. D **47**, 4315. *128*
- [160] Morgan, L., Morgan, T.: 1970, Phys. Rev. D **2**, 2756. *45*
- [161] Morris, M.: 1989, Phys. Rev. D **39**, 1511. *127*
- [162] Müller, V.: 1986, Ann. Phys. (Leipz.) **43**, 67. *125*
- [163] Müller, V., Schmidt, H.-J.: 1985, Gen. Relat. Grav. **17**, 769. 1989, Gen. Relat. Grav. **21**, 489. *28, 71, 98, 99, 105*
- [164] Müller, V., Schmidt, H.-J., Starobinsky, A.: 1988, Phys. Lett. B **202**, 198. *125*
- [165] Murphy, G.: 1973, Phys. Rev. D **8**, 4231. *123*
- [166] Nakao, K., Nakamura, T., Oohara, K., Maeda, K.: 1991, Phys. Rev. D **43**, 1788. *127*
- [167] Nakao, K., Maeda, K., Nakamura, T., Oohara, K.: 1993, Phys. Rev. D **47**, 3194 (1993). *127*
- [168] Nakao, K., Shiromizu, T., Maeda, K.: 1994, Class. Quant. Grav. **11**, 2059. *123*
- [169] Nariai, H.: 1973, Progr. Theor. Phys. **49**, 165. 1974, Progr. Theor. Phys. **51**, 613. *87*
- [170] Neugebauer, G.: 1980, *Relativistische Thermodynamik*, Akad.-Verl. Berlin. *46*
- [171] Nielsen, H., Olesen, P.: 1973, Nucl. Phys. B **61**, 45. *176*
- [172] Novotny, J.: 1984, Coll. J. Bolyai Math. Soc. Diff. Geom. Debrecen **46**, 959. *94*

- [173] Oleak, H.: 1974, *Astron. Nachr.* **295**, 107. 1987, *Ann. Phys. (Leipz.)* **44**, 74. *124*
- [174] Ono, S., Kondo, S.: 1960, in: Flügge, S., ed., *Handbuch der Physik* **10**, Springer Berlin, 134. *46, 47*
- [175] Page, D.: 1984, *Class. Quant. Grav.* **1**, 417. 1987, *Phys. Rev. D* **36**, 1607. *57, 59, 61, 67, 74*
- [176] Paiva, F., Reboucas, M., Hall, G., MacCallum, M.: 1998, *Class. Quant. Grav.* **15**, 1031. *44*
- [177] Papapetrou, A., Hamoui, A.: 1968, *Ann. Inst. H. Poincaré A* **9**, 179. *45*
- [178] Papapetrou, A., Treder, H.-J.: 1959, *Math. Nachr.* **20**, 53. *45*
- [179] Parker, L., Raval, A.: 1999, *Phys. Rev. D* **60** 063512. *83*
- [180] Pechlaner, E., Sexl, R.: 1966, *Commun. Math. Phys.* **2**, 165. *149, 160*
- [181] Peter, P., Polarski, D., Starobinsky, A.: 1994, *Phys. Rev. D* **50**, 4827. *123*
- [182] Pimentel, L.: 2001, *Gen. Relat. Grav.* **33**, 781; gr-qc/0009046. *83*
- [183] Polijevktov-Nikoladze, N.: 1967, *J. Exp. Teor. Fiziki* **52**, 1360. *160*
- [184] Pollney, D., Skea, J., D’Inverno, R.: 2000, *Class. Quant. Grav.* **17**, 2885. *44*
- [185] Pollock, M.: 1987, *Phys. Lett. B* **192**, 59. *125, 126*
- [186] Prigogine, I., Geheniau, J., Gunzig, E., Nardone, P.: 1988, *Proc. Nat. Acad. Sci.* **85**, 7428. 1989, *Gen. Relat. Grav.* **21**, 767. 1989, *Int. J. Theor. Phys.* **28**, 927. *124*
- [187] Quandt, I., Schmidt, H.-J.: 1991, *Astron. Nachr.* **312**, 97; gr-qc/0109005. *128, 159, 171*
- [188] Rainer, M.: 1995, *Grav. Cosmol.* **1**, 121. 1995, *Int. J. Mod. Phys. D* **4**, 397. *44*

- [189] Rainer, M., Schmidt, H.-J.: 1995, Gen. Relat. Grav. **27**, 1265; gr-qc/9507013. *14, 32, 33, 34*
- [190] Rainer, M., Zhuk, A.: 2000, Gen. Relat. Grav. **32**, 79. *111*
- [191] Robinson, A.: 1966, *Non-standard analysis*, North Holland Amsterdam. *152*
- [192] Rogers, B., Isaacson, J.: 1991, Nucl. Phys. B **364**, 381. 1992, Nucl. Phys. B **368**, 415. *127*
- [193] Romero, C., Tavakol, R., Zalaletdinov, R.: 1996, Gen. Relat. Grav. **28**, 365. *112*
- [194] Saa, A.: 1996, J. Math. Phys. **37**, 2346. *122*
- [195] Sakharov, A.: 2000, Gen. Relat. Grav. **32**, 365; 1967, Dokl. Akad. Nauk SSSR **177**, 70. *128*
- [196] Sanchez, M.: 1998, Gen. Relat. Grav. **30**, 915. *29*
- [197] Sanders, R. : 1984, Astron. Astrophys. **136**, L21. 1986, Astrophys. J. **154**, 135. *165*
- [198] Sanyal, A., Modak, B.: 2001, Phys. Rev. D **63**, 064021; gr-qc/0107001. *83*
- [199] Schimming, R., Schmidt, H.-J.: 1990, NTM-Schriftenr. Geschichte der Naturwiss., Technik, Medizin **27**, 41. *128*
- [200] Schmidt, H.-J.: 1982, Astron. Nachr. **303**, 227; gr-qc/0105104. *9, 15*
- [201] Schmidt, H.-J.: 1982, Astron. Nachr. **303**, 283; gr-qc/0105105. *19*
- [202] Schmidt, H.-J.: 1983, in: Bertotti, B., ed., *Contributed Papers Conf. General Relativity GR10 Padova* **1**, 339. *45, 46*
- [203] Schmidt, H.-J.: 1984, Gen. Relat. Grav. **16**, 1053; gr-qc/0105106. *46*
- [204] Schmidt, H.-J.: 1984, Ann. Phys. (Leipz.) **41**, 435; gr-qc/0105108. *156*
- [205] Schmidt, H.-J.: 1985, Astron. Nachr. **306**, 67; gr-qc/0105107. *145*

- [206] Schmidt, H.-J.: 1986, Conf. General Relativity GR 11 Stockholm, 117. 87
- [207] Schmidt, H.-J.: 1986, *Physikalische Konsequenzen von Feldgleichungen vierter Ordnung bei der Beschreibung des frühen Universums und isolierter Gravitationsfelder*, Thesis B, Akad. d. Wiss. Berlin; 2nd ed.: 1994, Tectum-Verlag Marburg. 87, 88, 89, 106
- [208] Schmidt, H.-J.: 1986, Astron. Nachr. **307**, 339; gr-qc/0106037. 159, 170
- [209] Schmidt, H.-J.: 1987, Astron. Nachr. **308**, 183; gr-qc/0106035. 87, 103, 110
- [210] Schmidt, H.-J.: 1987, J. Math. Phys. **28**, 1928. Addendum: 1988, J. Math. Phys. **29**, 1264. 14, 42, 73
- [211] Schmidt, H.-J.: 1988, Class. Quant. Grav. **5**, 233. 110
- [212] Schmidt, H.-J.: 1988, Astron. Nachr. **309**, 307; gr-qc/0106036. 87
- [213] Schmidt, H.-J.: 1988, Phys. Lett. B **214**, 519. 75, 121
- [214] Schmidt, H.-J.: 1989, Class. Quant. Grav. **6**, 557. 1994, Phys. Rev. D **50**, 5452; gr-qc/0109006. 1995, Phys. Rev. D **52**, 6198; gr-qc/0106034. 112
- [215] Schmidt, H.-J.: 1990, Astron. Nachr. **311**, 99; gr-qc/0108087. 58
- [216] Schmidt, H.-J.: 1990, Astron. Nachr. **311**, 165; gr-qc/0109004. 113
- [217] Schmidt, H.-J.: 1990, in: Janyska, J., Krupka, D., eds., *Proc. Conf. Brno, Diff. Geom. Appl.*, WSPC Singapore, 405; gr-qc/0109001. 77
- [218] Schmidt, H.-J.: 1990, Class. Quantum Grav. **7**, 1023. 128, 139, 164, 165, 172
- [219] Schmidt, H.-J.: 1992, in: Sato, H., Nakamura, T., eds., *Proc. Sixth Marcel Grossmann Meeting on General Relativity Kyoto*, WSPC Singapore, 92. 129
- [220] Schmidt, H.-J.: 1993, Gen. Relat. Grav. **25**, 87. Erratum **25**, 863. 127

- [221] Schmidt, H.-J.: 1994, Phys. Rev. D **49**, 6354. Erratum: 1996, Phys. Rev. D **54**, 7906; gr-qc/9404038. *27*
- [222] Schmidt, H.-J.: 1995, Abstr. Conf. General Relativity GR 14 Florence; gr-qc/0109007. *31*
- [223] Schmidt, H.-J.: 1996, in: Janyska, J., Kolar, I., Slovak, J., eds., *Proc. Conf. Brno Diff. Geom. Appl.*, 119; gr-qc/0109008. *31*
- [224] Schmidt, H.-J.: 1996, in: Sardanashvily, G., ed., *New frontiers in gravitation*, Hadronic Press Palm Harbor, 337; gr-qc/9404037. *31, 44*
- [225] Schmidt, H.-J.: 1996, Gen. Relat. Grav. **28**, 899; gr-qc/9512006. *25*
- [226] Schmidt, H.-J.: 1997, Grav. Cosmol. **3**, 185; gr-qc/9709071. *19*
- [227] Schmidt, H.-J.: 1998, Int. J. Theor. Phys. **37**, 691; gr-qc/9512007. *29*
- [228] Schmidt, H.-J.: 1998, in: Rainer, M., Schmidt, H.-J., eds., *Current topics in mathematical cosmology*, WSPC Singapore, 288; gr-qc/9808060. 1993, Fortschr. Phys. **41**, 179. *27*
- [229] Schmidt, H.-J.: 1999, Gen. Relat. Grav. **31**, 1187; gr-qc/9905051. *111*
- [230] Schmidt, H.-J.: 2000, Ann. Phys. (Leipz.) **9**, SI-158. gr-qc/9905103. *44*
- [231] Schmidt, H.-J., Müller, V.: 1985, Gen. Relat. Grav. **17**, 971. *150*
- [232] Schmidt, H.-J., Semmelmann, U.: 1989, Astron. Nachr. **310**, 103; gr-qc/0106089. *175*
- [233] Senovilla, J.: 1998, Gen. Relat. Grav. **30**, 701. *27*
- [234] Schrödinger, E.: 1938, Comment. Vatican. Acad. **2**, 321. 1956, *Expanding Universes*, Cambridge University Press. *58*
- [235] Shibata, M., Nakao, K., Nakamura, T., Maeda, K.: 1994, Phys. Rev. D **50**, 708. *124*
- [236] Shinkai, H., Maeda, K.: 1993, Phys. Rev. D **48**, 3910. 1994, Phys. Rev. D **49**, 6367. *127*
- [237] Shiromizu, T., Nakao, K., Kodama, H., Maeda, K.: 1993, Phys. Rev. D **47**, R3099. *124, 127*

- [238] Singh, T., Padmanabhan, T.: 1988, Int. J. Mod. Phys. A **3**, 1593. 75
- [239] Sirousse-Zia, H.: 1982, Gen. Relat. Grav. **14**, 751. 125
- [240] Spero, A., Baierlein, R.: 1978, J. Math. Phys. **19**, 1324. 9
- [241] Spindel, P.: 1994, Int. J. Mod. Phys. D **3**, 273. 125, 127, 139
- [242] Stacey, F., Tuck, G., Holding, S., Maher, A., Morris, D.: 1981, Phys. Rev. D **23**, 1683. 165
- [243] Starobinsky, A.: 1978, Pis'ma Astron. Zh. **4**, 155. 58, 74, 107
- [244] Starobinsky, A.: 1983, Sov. Phys. JETP Lett. **37**, 66. 126, 127
- [245] Starobinsky, A., Schmidt, H.-J.: 1987, Class. Quant. Grav. **4**, 695. 61, 103, 106, 127
- [246] Starobinsky, A., Tsujikawa, S., Yokoyama, J: 2001, *Cosmological perturbations from multi-field inflation in generalized Einstein theories*, astro-ph/0107555; Nucl. Phys. B in press. 112
- [247] Stein-Schabes, J., Burd, A.: 1988, Phys. Rev. D **37**, 1401. 179
- [248] Stelle, K.: 1977, Phys. Rev. D **16**, 953. 90, 122, 128
- [249] Stelle, K.: 1978, Gen. Relat. Grav. **9**, 353. 146, 149, 160, 163, 164, 170
- [250] Stephani, H.: 1982, *General Relativity*, Cambridge University Press. 7, 36, 164
- [251] Szekeres, P.: 1975, Commun. Math. Phys. **41**, 55. 9, 21, 25
- [252] Taub, A.: 1980, J. Math. Phys. **21**, 1423. 49
- [253] Teyssandier, P.: 1989, Class. Quant. Grav. **6**, 219. 164
- [254] Thompson, J.: 1988, Phys. Rev. D **37**, 283. 182
- [255] Thomson, W. (Lord Kelvin): 1858, Proc. R. Soc. London **9**, 255. 47
- [256] Tolman, R. C.: 1934, *Relativity Thermodynamics and Cosmology*, Clarendon Press Oxford. 19

- [257] Treder, H.-J.: 1975, Ann. Phys. (Leipz.) **32**, 383. 1977, in: Brauer, W., ed., *75 Jahre Quantentheorie*, Akad.-Verl. Berlin, 279. 1991, Found. Phys. **21**, 283. *129, 146*
- [258] Turner, M.: 1983, Phys. Rev. D **28**, 1243. *59*
- [259] Turner, M., Widrow, L.: 1988, Phys. Rev. D **37**, 3428. *75*
- [260] Urbandtke, H.: 1972, Acta Physica Austriaca **35**, 1. *45*
- [261] Vilenkin, A.: 1985, Phys. Rep. **121**, 263. *175*
- [262] Vilenkin, A.: 1992, Phys. Rev. D **46**, 2355. *124*
- [263] Vilkovisky, G.: 1992, Class. Quant. Grav. **9**, 895. *128, 129*
- [264] Voorhees, B.: 1972, Phys. Rev. D **5**, 2413. *45*
- [265] Wainwright, J.: 1981, J. Phys. A **14**, 1131. *9, 19*
- [266] Wands, D.: 1994, Class. Quant. Grav. **11**, 269. *129*
- [267] Weinberg, S.: 1979, in: Hawking, S., Israel, W., eds., *General Relativity*, Cambridge University Press., 790. *105*
- [268] Weyl, H.: 1919, Ann. Phys. (Leipz.) **59**, 101. 1921, *Raum, Zeit, Materie*, Springer Berlin. *145, 156*
- [269] Weyl, H.: 1927, *Handbuch der Philosophie*, chapter “Philosophie der Mathematik und Naturwissenschaft”, Oldenburg. *122*
- [270] Whitt, B.: 1984, Phys. Lett. B **145**, 176. *88*
- [271] Wünsch, V.: 1976, Math. Nachr. **73**, 37. *147*
- [272] Xu, J., Li, L., Liu, L.: 1994, Phys. Rev. D **50**, 4886. *125*
- [273] Yokoyama, J. and Maeda, K.: 1990, Phys. Rev. D **41**, 1047. *125*
- [274] Yukawa, H.: 1935, Proc. Math. Phys. Soc. Japan **17**, 48. *164*
- [275] Zalaletdinov, R.: 1996, Gen. Relat. Grav. **28**, 953. *112*
- [276] Zeldovich, Y.: 1980, Mon. Not. R. Astron. Soc. **192**, 663. *175*